

## ABOUT DIFFERENTIABILITY AND $VBG_*$ CLASS

MAŁGORZATA TUROWSKA

### ABSTRACT

Let  $X$  be a finite dimensional real Banach space. We show that if the contingent of the curve  $\Gamma: (a, b) \rightarrow X$  fulfils some conditions then each parametrization of that curve is  $VBG_*$ . Stanisław Saks proved that each  $VBG_*$  function is differentiable at a set of full Lebesgue measure. The result of this paper is a partial converse of that theorem.

### 1. INTRODUCTION

We will present a generalization of the concepts of functions of bounded variation in the restricted sense ( $VB_*$ ) and of generalized bounded variation in the restricted sense ( $VBG_*$ ) in the case of functions of a real variable that takes values in a real normed space. Let us recall first these definitions in the case of real-valued functions.

**Definition 1.** [2], [5] *If  $F: [a, b] \rightarrow \mathbb{R}$  and  $[\alpha, \beta] \subset [a, b]$ , then the value*

$$\sup \{ |F(x) - F(y)| : x \in [\alpha, \beta], y \in [\alpha, \beta] \}$$

*is called an oscillation of the mapping  $F$  on the interval  $[\alpha, \beta]$  and is denoted by  $\omega(F, [\alpha, \beta])$ .*

**Definition 2.** [2], [5] *If  $F: [a, b] \rightarrow \mathbb{R}$  and  $E \subset [a, b]$  then a mapping  $F$  is called of bounded variation in the restricted sense on the set  $E$ , or simply, is of  $VB_*$  on  $E$ , if*

$$\sup \sum_k \omega(F, [a_k, b_k]) < \infty,$$

*where  $([a_k, b_k])_{k \in \mathbb{N}}$  is any sequence of non-overlapping intervals such that  $a_k \in E$ ,  $b_k \in E$ . The number  $\sup \sum_k \omega(F, [a_k, b_k])$  is denoted by  $V_E F$ .*

Definition 2 can be generalized on the case of the mapping  $F$  with value in a real normed space  $X$ .

---

• Małgorzata Turowska — e-mail: malgorzata.turowska@apsl.edu.pl  
Pomeranian University in Słupsk.

**Definition 3.** Let  $X$  be a real normed space and  $\|\cdot\|$  be the norm in  $X$ . By the oscillation of a mapping  $F: [a, b] \rightarrow X$  on  $[\alpha, \beta] \subset [a, b]$  we call the value

$$\sup \{ \|F(x) - F(y)\| : x \in [\alpha, \beta], y \in [\alpha, \beta] \}.$$

This oscillation will be denoted by the symbol  $\omega(F, [\alpha, \beta])$ .

**Definition 4.** Let  $X$  be a real normed space and  $E \subset [a, b]$ . We say that a mapping  $F: [a, b] \rightarrow X$  is  $VB_*$  on the set  $E$ , and denote  $F \in VB_*(E)$ , if

$$\sup \sum_k \omega(F, [a_k, b_k]) < \infty,$$

where  $([a_k, b_k])_{k \in \mathbb{N}}$  is any sequence of non-overlapping intervals such that  $a_k \in E$ ,  $b_k \in E$ . The value  $\sup \sum_k \omega(F, [a_k, b_k])$  is denoted by  $V_E F$ .

Now we assume that dimension of  $X$  is finite. Observe that the fact that  $F$  is  $VB_*$  on some set is independent of the choice of a norm in  $X$ . Let  $F: [a, b] \rightarrow X$  and  $e = (e_1, \dots, e_n)$  be a base of the space  $X$ . Then

$$F = \sum_{i=1}^n F_i e_i.$$

Mappings  $F_i$  are called coordinates of the mapping  $F$  with respect to the base  $e$ . We also shall use denotation  $F = (F_1, \dots, F_n)$ . Straightforward calculations prove the next lemma.

**Lemma 1.** If  $X$  is a finite dimensional real normed space,  $F: [a, b] \rightarrow X$  and  $E \subset [a, b]$ , then:

- (1) If  $F$  is  $VB_*$  on  $E$  then for each base  $e = (e_1, \dots, e_n)$  of the space  $X$  mappings  $F_i$  are  $VB_*$  on  $E$  for each  $i \in \{1, \dots, n\}$ .
- (2) If there exists a base  $e = (e_1, \dots, e_n)$  of the space  $X$  for which mappings  $F_i$ ,  $i \in \{1, \dots, n\}$ , are  $VB_*$  on the set  $E$  then  $F \in VB_*(E)$ .

**Definition 5.** [2], [5] Let  $E \subset [a, b]$ . We say that a mapping  $F: [a, b] \rightarrow \mathbb{R}$  is of generalized bounded variation in the restricted sense on  $E$ , or simply, is  $VBG_*$  on the set  $E$ , and denote  $F \in VBG_*(E)$ , if  $E$  is a countable union of sets on each of which the mapping  $F$  is  $VB_*$ .

We can generalize this definition in the following way:

**Definition 6.** Let  $X$  be a real normed space,  $E \subset [a, b]$ . We will say that a mapping  $F: [a, b] \rightarrow X$  is  $VBG_*$  on  $E$ , and denote  $F \in VBG_*(E)$ , if  $E$  is a countable union of sets such that for each of them  $F$  is  $VB_*$ .

The proof of the next lemma is technical, we shall omit it.

**Lemma 2.** Let  $X$  be a real normed space,  $\dim X = n$ ,  $F: [a, b] \rightarrow X$  and  $E \subset [a, b]$ . Then

- (1) If  $F$  is  $VBG_*$  on  $E$  then for each base  $e = (e_1, \dots, e_n)$  of the space  $X$  mappings  $F_i$  are  $VBG_*$  on the set  $E$  for each  $i \in \{1, \dots, n\}$ .
- (2) If there exists a base  $e = (e_1, \dots, e_n)$  of the space  $X$  for which each mapping  $F_i$ ,  $i \in \{1, \dots, n\}$ , is  $VBG_*$  on  $E$  then  $F$  is  $VBG_*$  on  $E$ .

**Theorem 1.** [5] *Let  $E \subset [a, b]$ . If a function  $F: [a, b] \rightarrow \mathbb{R}$  is  $VBG_*$  on the set  $E$ , then  $F$  is differentiable at a set of full Lebesgue measure.*

The obvious corollary of this theorem for a mappings which take values in a real normed space is as follows:

**Corollary 1.** *Let  $X$  be a real normed space,  $\dim X < \infty$  and  $E \subset [a, b]$ . If a mapping  $F: [a, b] \rightarrow X$  is  $VBG_*$  on the set  $E$ , then  $F$  is differentiable (in the Fréchet sense) at almost all points of this set.*

**Definition 7.** [6] *Let  $\emptyset \neq M \subset Z$ , where  $Z$  is a real normed space. Let  $z$  belong to the closure of the set  $M$ . The set*

$$\left\{ v \in Z : \exists (z_n)_{n \in \mathbb{N}}, z_n \in M, \lim_{n \rightarrow \infty} z_n = z, \exists (\lambda_n)_{n \in \mathbb{N}}, \lambda_n > 0 : \lim_{n \rightarrow \infty} \lambda_n (z_n - z) = v \right\}$$

*is called the tangent cone to  $M$  at  $z$  and is denoted by  $\text{Tan}(M, z)$ . The elements of  $\text{Tan}(M, z)$  are called vectors tangent to  $M$  at  $z$ . The set  $\text{Tan}(M, z)$  is also called the contingent of  $M$  at  $z$  (see [1], [5]).*

The basic properties of the contingent and the connections between differentiability of a mapping  $f: X \rightarrow Y$  at a point, where  $X, Y$  are real normed spaces and the contingent of its graph one can find in [3], [4], [6], [7].

**Definition 8.** *If  $X$  is a real normed space, then a mapping  $f$  is called an embedding if it is a homeomorphism of the interval  $(a, b)$  into  $X$ , where  $f((a, b))$  is equipped with the subspace topology. A subset  $\Gamma$  of the space  $X$  is called a curve if there is an embedding  $f: (a, b) \rightarrow X$  such that  $f((a, b)) = \Gamma$ . This embedding is called a parametrization of the curve  $\Gamma$ .*

The following theorem gives a connection between the contingent of a curve and the existence of a differentiable parametrization of this curve.

**Theorem 2.** [8] *Let  $X$  be a real normed space for which  $1 < \dim X < \infty$ . Assume that for a curve  $\Gamma \subset X$  the following conditions are fulfilled:*

- (i) *for each  $p \in \Gamma$  the contingent  $\text{Tan}(\Gamma, p)$  is one-dimensional linear subspace of  $X$ ,*
- (ii) *there exists a subspace  $Y$  of  $X$  such that  $\text{codim} Y = 1$  and*

$$\text{Tan}(\Gamma, p) \not\subset Y$$

*for each  $p \in \Gamma$ .*

Then there exist an open interval  $(c, d)$  and a differentiable parametrization  $g: (c, d) \rightarrow \Gamma$  of the curve  $\Gamma$  such that

$$\inf_{t \in (c, d)} \|g'(t)\| > 0.$$

**Corollary 2.** [8] *Let  $f: (a, b) \rightarrow \Gamma$  be a parametrization of the curve  $\Gamma$ . Then under assumptions of theorem 2, for every open interval  $(c, d)$  there exists a mapping  $g: (c, d) \rightarrow \Gamma$  such that the mapping  $g^{-1} \circ f: (a, b) \rightarrow (c, d)$  is an increasing homeomorphism.*

**Corollary 3.** [8] *Under assumptions of theorem 2, each parametrization of the curve  $\Gamma$  is almost everywhere differentiable.*

**Theorem 3.** [2] *A mapping  $F: [0, 1] \rightarrow \mathbb{R}$  is continuous and  $VBG_*$  on the interval  $[0, 1]$  if and only if there exists a homeomorphism  $h: [0, 1] \rightarrow [0, 1]$  such that  $F \circ h$  is differentiable.*

We will use the following easy generalization of theorem 3.

**Theorem 4.** *Let  $F: [0, 1] \rightarrow \mathbb{R}$ . The mapping  $F$  is continuous and  $VBG_*$  on  $[0, 1]$  if and only if there exists a homeomorphism  $h: [c, d] \rightarrow [0, 1]$  such that  $F \circ h$  is differentiable.*

## 2. MAIN RESULTS

Applying theorem 2., lemma 2. and theorem 4. we will prove that each parametrization of a curve  $\Gamma$  satisfying assumptions of theorem 2 is  $VBG_*$ . The following theorem is a partial converse of the corollary 1.

**Theorem 5.** *Let  $X$  be a real normed space such that  $1 < \dim X < \infty$ . Assume that for a curve  $\Gamma \subset X$  the following conditions are fulfilled:*

- (i) *for each  $p \in \Gamma$  the contingent  $\text{Tan}(\Gamma, p)$  is one-dimensional linear subspace of  $X$ ,*
- (ii) *there exists a subspace  $Y$  of  $X$  such that  $\text{codim} Y = 1$  and*

$$\text{Tan}(\Gamma, p) \not\subset Y$$

*for each  $p \in \Gamma$ .*

*Then each parametrization  $f: (a, b) \rightarrow \Gamma$  of the curve  $\Gamma$  is  $VBG_*$  in  $(a, b)$ .*

*Proof.* Let  $f: (a, b) \rightarrow \Gamma$  be a parametrization of the curve  $\Gamma$ . By theorem 2, there exists a differentiable parametrization  $g: (c, d) \rightarrow \Gamma$  of that curve. Obviously  $f^{-1} \circ g$  is a homeomorphism of  $(c, d)$  onto  $(a, b)$ .

Fix an interval  $[c_1, d_1]$  contained in  $(c, d)$ . Then there exists an interval  $[a_1, b_1]$  in the set  $(a, b)$  such that the mapping

$$(f^{-1} \circ g)|_{[c_1, d_1]}: [c_1, d_1] \rightarrow [a_1, b_1]$$

is a homeomorphism of  $[c_1, d_1]$  onto  $[a_1, b_1]$ .

Denote  $h^* = (f^{-1} \circ g)|_{[c_1, d_1]}$ ,  $f^* = f|_{[a_1, b_1]}$  and  $g^* = g|_{[c_1, d_1]}$ . Obviously,  $f^*$  is continuous and  $g^*$  is differentiable.

Fix a base  $e = (e_1, \dots, e_n)$  of the space  $X$ . Then

$$f^*(t) = \sum_{i=1}^n f_i^*(t)e_i \quad \text{and} \quad g^*(\tau) = \sum_{i=1}^n g_i^*(\tau)e_i,$$

where  $f_i^*: [a_1, b_1] \rightarrow \mathbb{R}$ ,  $g_i^*: [c_1, d_1] \rightarrow \mathbb{R}$ ,  $i \in \{1, \dots, n\}$  and  $t \in [a_1, b_1]$ ,  $\tau \in [c_1, d_1]$ . Since  $g^* = f^* \circ h^*$ , then  $g_i^* = f_i^* \circ h^*$  for each  $i \in \{1, \dots, n\}$ . The mapping  $g^*$  is differentiable, so  $g_i^*$  is differentiable if  $i \in \{1, \dots, n\}$ .

Moreover,  $h^*$  is a homeomorphism and  $f_i^*$  is continuous if  $i \in \{1, \dots, n\}$  and by theorem 4 we have  $f_i^* \in VBG_*([a_1, b_1])$  for each  $i \in \{1, \dots, n\}$ .

By lemma 2(2) we conclude that the mapping

$$f^*: [a_1, b_1] \rightarrow X$$

is  $VBG_*$  in  $[a_1, b_1]$ . Therefore the mapping  $f$  is  $VBG_*$  on each closed subinterval of  $(a, b)$ . The interval  $(a, b)$  is a countable union of closed subintervals, so the mapping  $f$  is  $VBG_*$  on  $(a, b)$ .  $\square$

#### REFERENCES

- [1] G. Bouligand, *Introduction à la géométrie infinitésimale directe*, Librairie Vuibert, Paris 1932.
- [2] A. M. Bruckner, *Differentiation of Real Functions*, Lecture Notes in Math., 659, Springer-Verlag, Berlin 1978.
- [3] H. Federer, *Geometric Measure Theory*, Springer Verlag, Berlin, Heidelberg, New York 1969.
- [4] T. M. Flett, *Differential analysis*, Cambridge University Press, London 1980.
- [5] S. Saks, *Theory of the Integral*, G. E. Stechert & Co., Warszawa-Lwów-New York 1937.
- [6] L. Schwartz, *Analyse Mathématique*, vol 1, Hermann, Paris 1967.
- [7] M. Turowska, *A geometric condition for differentiability*, Tatra Mt. Math. Publ. 28 (2004), 179-186.
- [8] M. Turowska, *On curves possessing parametrizations whose derivatives are separated from zero*, Real functions, density topology and related topics, Łódź University Press 2011, 173-182.

Received: October 2014

Małgorzata Turowska  
 POMERANIAN UNIVERSITY IN SŁUPSK,  
 INSTITUTE OF MATHEMATICS,  
 UL. ARCISZEWSKIEGO 22D, 76-200 SŁUPSK, POLAND  
 E-mail address: malgorzata.turowska@aps1.edu.pl