



## Impact of climatic changes in the Late Pleistocene on migrations and extinctions of mammals in Europe: four case studies

Mateusz BACA<sup>1</sup>, Adam NADACHOWSKI<sup>2</sup>\*, Grzegorz LIPECKI<sup>2</sup>, Paweł MACKIEWICZ<sup>3</sup>,  
Adrian MARCISZAK<sup>4</sup>, Danijela POPOVIĆ<sup>5</sup>, Paweł SOCHA<sup>4</sup>, Krzysztof STEFANIAK<sup>4</sup> and Piotr WOJTAL<sup>2</sup>

- <sup>1</sup> University of Warsaw, Center of Precolumbian Studies, Krakowskie Przedmieście 26/28, 00-927 Warszawa, Poland
- <sup>2</sup> Institute of Systematics and Evolution of Animals, Polish Academy of Sciences, Sławkowska 17, 31-016 Kraków, Poland
- <sup>3</sup> University of Wrocław, Department of Genomics, Faculty of Biotechnology, Joliot-Curie 14a, 50-383 Wrocław, Poland
- <sup>4</sup> University of Wrocław, Department of Paleozoology, Institute of Environmental Biology, Sienkiewicza 21, 50-335 Wrocław, Poland
- <sup>5</sup> University of Warsaw, Centre of New Technologies, Banacha 2C, 02-097 Warszawa, Poland



Baca, M., Nadachowski, A., Lipecki, G., Mackiewicz, P., Marciszak, A., Popović, D., Socha, P., Stefaniak, K., Wojtal, P., 2017. Impact of climatic changes in the Late Pleistocene on migrations and extinction of mammals in Europe: four case studies. *Geological Quarterly*, 61 (2): 291–304, doi: 10.7306/gq.1319

Climate changes that occurred during the Late Pleistocene had profound effects on the distribution of many plant and animal species and influenced the formation of contemporary faunas and floras of Europe. The course and mechanisms of responses of species to past climate changes are now being intensely studied by the use of direct radiocarbon dating and genetic analyses of fossil remains. Here, we review the advances in understanding these processes by the example of four mammal species: woolly mammoth (*Mammuthus primigenius*), cave bear (*Ursus spelaeus* s.l.), saiga antelope (*Saiga tatarica*) and collared lemmings (*Dicrostonyx* ssp.). The cases discussed here as well as others show that migrations, range shifts and local extinctions were the main responses to climate changes and that the dynamics of these climate-driven processes were much more profound than was previously thought. Each species reacted in its individual manner, which depended on its biology and adaptation abilities to changing environmental and climatic conditions. The most severe changes in European ecosystems that affected the largest number of species took place around 33–31 ka BP, during the Last Glacial Maximum 22–19 ka BP and the Late Glacial warming 15–13 ka BP.

Key words: ancient DNA, radiocarbon dating, migrations, climate changes.

### INTRODUCTION

The Late Pleistocene was an interval marked by multiple climate changes of a magnitude greater than those observed today (Rasmussen et al., 2014). The impact of those fluctuations on Late Pleistocene mammal populations has been intensely studied by examination of changes in spatial distribution of fossil mammalian faunas in time (e.g., Graham et al., 1996; Sommer and Nadachowski, 2006; Lister and Stuart, 2008; Markova and van Kolfschoten, 2008). This period witnessed also mass extinction on an unprecedented scale not observed in fossil records for millions of years (Sandom et al., 2014; Stuart, 2015). By the end of the Pleistocene, most terrestrial megafaunal species (heavier than 44 kg) became extinct or their population sizes decreased substantially (Barnosky et al., 2004; Koch and Barnosky, 2006). The course and the timing of

these events differ from region to region and the causes of these extinctions are the subject of ongoing debate. Some researchers point only to climate changes, some blame human activities such as overhunting and habitat alternation (Alroy, 2001; Sandom et al., 2014) whereas others suggest a combination of these two causes as the main trigger (Barnosky et al., 2004; Prescott et al., 2012; Stuart, 2015).

In recent years two main advances accelerated research in this field. The first one was an increase in the accessibility and popularity of AMS radiocarbon dating. The growing number of directly dated fossils allowed for precise tracking of changes in species' geographic distribution, migrations and dating extinction events (e.g., MacPhee et al., 2002; Stuart, 2005; Sommer et al., 2008, 2014; Pacher and Stuart, 2009; Stuart and Lister, 2012, 2014). It enabled direct correlation between these events and climatic and environmental changes. The second advance was the emergence of ancient DNA studies. The examination of genetic diversity has added another level of complexity to research into the Late Pleistocene populations. Ancient DNA facilitated investigations of intraspecific processes, such as population replacements or changes in their diversity; events that usually do not manifest in the fossil record. These two methods

\* Corresponding author, e-mail: nadachowski@isez.pan.krakow.pl  
Received: April 18, 2016; accepted: June 13, 2016; first published online: March 6, 2017



combined proved to be the most versatile approach in reconstruction of past population histories (e.g., Campos et al., 2010a, b; Lorenzen et al., 2011; Horn et al., 2014; Palkopoulou et al., 2016).

The main aim of our study is to review the recent progress that has been made in research into four species: woolly mammoth (*Mammuthus primigenius*), cave bear (*Ursus spelaeus* s.l.), saiga antelope (*Saiga tatarica*) and collared lemming (*Dicrostonyx* ssp.) and to discuss its implications for our understanding of the impact of climate changes on migrations and extinctions of mammals in the Late Pleistocene.

### WOOLLY MAMMOTH (*MAMMUTHUS PRIMIGENIUS*)

Elephants (Elephantidae) are the largest terrestrial mammals, with a stout body, characteristic long, highly pronounced proboscis or trunk, a combination of nose and upper lip, large ears, column-like limbs, and small tail. The woolly mammoth represents elephants well-adapted to the cold and arid steppe-tundra environment (Maschenko, 2002; Lister et al., 2005). This species was widespread during the Late Pleistocene from Western Europe through the whole of northern Asia

to the northern part of North America (Kahlke, 2015). Morphological data has revealed that woolly mammoths were present in Europe from around 200 ka BP until the end of the Pleistocene (Lister and Sher, 2001; Lister et al., 2005). However, genetic studies indicated that the Late Pleistocene history of *Mammuthus primigenius* was characterized by a complex series of range expansions and contractions, demographic changes and clade replacements (Palkopoulou et al., 2013).

Phylogenetic analyses revealed that Holarctic mammoths belonged to three distinct mitochondrial (mtDNA) lineages (Fig. 1). The most widespread lineage I had a nearly Holarctic distribution, lineage II was confined to Central-East Asia while specimens from lineage III are known from Europe (Debruyne et al., 2008; Palkopoulou et al., 2013). The divergence of lineages I and II was previously estimated to ca. 1 Ma (Debruyne et al., 2008; Gilbert and Drautz, 2008), however, most recent estimations suggest a much younger date, about 300 ka BP (Palkopoulou et al., 2013). Coalescent simulations suggested that a split of three mammoth populations took place around 200 ka BP and was followed by a demographic expansion that started around 121 ka BP (Palkopoulou et al., 2013). This expansion coincides broadly with the end of Eemian Interstadial, which suggests that mammoths survived this warm period confined to refugial areas and expanded as climate became cooler at the beginning of the Weichselian Glaciation (Palkopoulou et




Fig. 1. Woolly mammoth (*Mammuthus primigenius*)

**A** – Bayesian phylogeny of Holarctic woolly mammoths based on mtDNA cytochrome b sequences. The tree is a chronogram where branch lengths denote time elapsed since divergence and the positions of tips correspond to calibrated radiocarbon ages of samples; **B** – distribution of palaeontological sites with woolly mammoth remains radiocarbon-dated to the periods indicated. Colours indicate mitochondrial DNA lineages (modified after Palkopoulou et al., 2013)



al., 2013). Surprisingly, this was not supported by the analyses of whole palaeogenomes, which indicated a much earlier expansion ca. 280 ka years ago and a maximum effective population size during the Eemian (Palkopoulou et al., 2015).

Despite these ambiguities in the early history of mammoth populations, ancient DNA revealed also two more recent population turnovers. In the Eemian Interglacial and Early Weichselian, woolly mammoths that belonged to clade I were most probably confined to North America. It was estimated that ca. 66 ka BP they started to expand westwards through Beringia and reached the area occupied by mammoths of clade II. Both populations lived in sympatry for 20 ka, when clade II suddenly disappeared from the fossil record ca. 40 ka BP (Palkopoulou et al., 2013). Woolly mammoths belonging to clade I expanded farther west and reached Europe. The earliest specimen in Europe originating from this clade came from the site in Vologda Oblast in Russia and was dated to 32 ka BP. The appearance of clade I in Europe coincides with the disappearance of the endemic European population (clade III), with the latest specimen dated to ca. 34 ka BP. There is, however, no evidence for any overlap between these populations and it seems that the extinction of clade III was not driven by the appearance of newcomers from the east. This scenario is supported by the lack of radiocarbon dating of mammoth remains in Central Europe between ca. 34 and 33 ka BP, e.g. in Poland (Nadachowski et al., 2011; Fig. 2). An important increase of mammoth population size in Europe took place between 31 and 29 ka BP, which is confirmed by dense radiocarbon dating of mammoth remains collected from almost the whole North European Plain (Nada-

chowski et al., 2011; Ukkonen et al., 2011; Markova et al., 2013). Interestingly, during almost the whole of the Last Glacial Maximum (LGM), between ca. 22 and 18 ka BP (Fig. 2), there are no dated mammoth records in northwestern, northern and central Europe sites (Stuart et al., 2004; Nadachowski et al., 2011; Ukkonen et al., 2011), which suggests a long-term contraction. Mammoths re-immigrated to Europe for the next 3–4 millennia at the end of GS-2 (Greenland Stadial 2) with the return of severe climatic conditions and open vegetation (Lister and Stuart, 2008; Nadachowski et al., 2011; Fig. 2). The next break in the mammoth record coincides roughly with the Bølling-Allerød warming (Greenland Interstadial 1; GI-1) followed by one more recolonization of Latvia and Estonia, dated to the Younger Dryas (GS-1) (Lõugas et al., 2002; Stuart et al., 2002; Ukkonen et al., 2011; see Fig. 7). The last populations of *Mammuthus primigenius* in Europe lived in northwestern Russia and disappeared around the Younger Dryas-Holocene boundary ca. 11.8–11.4 ka BP (Lõugas et al., 2002; Ukkonen et al., 2011).

### CAVE BEAR (*URSUS SPELAEUS* S.L.)

The bear family (Ursidae) are large mammals with a big head and thick neck, small eyes and short tail, muscular bodies with stout legs and large paws. Most bears are omnivores, although the diets of polar bear (*Ursus maritimus*) and giant




Fig. 2. Woolly mammoth (*Mammuthus primigenius*)

Radiocarbon dates of woolly mammoth remains from Poland alongside the NGRIP GICC05 ice core  $\delta^{18}\text{O}$  record (Svensson et al., 2008). Dates were compiled from Nadachowski et al. (2011), Pawłowska (2015), Wilczyński et al. (2015), Wojtal and Wilczyński (2015) and recalibrated in OxCal v. 4.2 (Ramsey, 2009) using the Intcal13 calibration curve (Reimer et al., 2013). MIS – Marine Isotopic Stages, GS – Greenland Stadials, GI – Greenland Interstadials; LGM – Last Glacial Maximum (after Mix et al., 2001). Grey stripes illustrate inferred periods of the absence of mammoths from Poland



panda (*Ailuropoda melanoleuca*) are very narrow and specialized. The former is a carnivore and the latter is an obligate consumer of bamboo. It is assumed that the cave bear (*Ursus spelaeus* s.l.), one of the most widespread mammals in the Late Pleistocene in Europe, was an obligate vegetarian (Baca et al., 2016). It evolved from the Middle Pleistocene *Ursus deningeri* and differentiated into several forms recognized at morphological or genetic levels (Hofreiter et al., 2002; Rabeder et al., 2004a, b). In Europe, two main forms, *U. spelaeus* and *U. ingressus* existed. They differentiated probably between 414,000 and 173,000 years ago (Knapp et al., 2009). *U. spelaeus* lived mainly in Western Europe and its remains have been found in Spain, France, Germany, Belgium, Italy and Austria, although it has also been recorded in Altai (Rabeder et al., 2004b; Knapp et al., 2009). *U. ingressus* inhabited southeastern and Central Europe and its remains have been discovered mainly in Romania, Slovenia, Ukraine, Czech Republic, Poland, Slovakia and Greece but also in Austria, Germany and Switzerland (Rabeder et al., 2004b; Baca et al., 2014). Moreover, two small, dwarf cave bear forms that are considered as subspecies, *U. spelaeus eremus* and *U. spelaeus ladinicus* have been reported in high alpine caves in Austria and Italy (Rabeder and Hofreiter, 2004; Rabeder et al., 2004a; Fig. 3A). Another major group of large bears, named *U. deningeri kudarensis*, was discovered in the Caucasus (Baryshnikov, 1998; Knapp et al., 2009). Recent analysis of mtDNA of the Middle Pleistocene *U. deningeri* from Sima de los Huesos, Atapuerca, Spain, revealed that *U. deningeri kudarensis* constitutes the most divergent cave bear lineage (Dabney et al., 2013). It was suggested that it belonged to a separate branch of cave bear evolution and hence its taxonomic status was changed to *U. kudarensis* (Dabney et al., 2013; Stiller et al., 2014).

The geographic origin of *U. ingressus* is not known, but the basal position of haplotypes from the Romanian site Peștera cu Oase in the phylogeny of this species points to southeastern Europe (Baca et al., 2012). The phylogeographic picture of *U. ingressus* is unclear as the mtDNA phylogeny lacks significant support and clear phylogeographic lineages are not differentiated (Fig. 3A). It seems, however, that the spread of *U.*

*ingressus* in Europe may have proceeded independently along the main European mountain ranges, the Alps and Carpathians (Baca et al., 2014).

It has been proposed that between 60 and 50 ka BP *U. ingressus* started westward migration along the Alps (Hofreiter et al., 2004; Rabeder and Hofreiter, 2004; Münzel et al., 2011). The earliest remains of this species from the Austrian and Swiss Alps, dated to ca. 50 ka BP, are known from Gamssulzen Cave. The westernmost record of *U. ingressus* came from the Schnurenloch site near Bern, Switzerland, but species attribution was based solely on morphology without genetic confirmation (Rabeder and Hofreiter, 2004). Recently, several cave bear specimens from the Kraków-Częstochowa Upland, Poland yielded mtDNA haplotypes identical to those found in Alpean *U. ingressus*, which suggests that this population spread also northwards beyond the Carpathian Arc (Popović et al., 2015).

In some cases, the appearance of *U. ingressus* was associated with the disappearance of other cave bear forms that inhabited the area. The most interesting evidence of such replacement came from the three cave sites in the Ach Valley in Swabian Jura, Germany: Hohle Fels, Geißenklösterle, and Sirgenstein (Hofreiter et al., 2007; Münzel et al., 2011). The latest occurrence of the native inhabitant in this area, *U. spelaeus*, was dated to ca. 31.5 ka BP while the earliest record of *U. ingressus* from Geißenklösterle was dated to 36.3 ka BP. Only a single *U. ingressus* individual yielded such a date while most of others were dated to ca. 32 ka BP, which suggests that the main immigration took place just before the local extinction of *U. spelaeus*. A similar replacement was recorded in Herdengel Cave in Austria, where *U. ingressus* replaced one of the small cave bear forms, *U. s. eremus*. All of the eremus cave bears were dated by stratigraphic context to more than 60 ka BP, whereas all *U. ingressus* were younger than 37 ka BP (Stiller et al., 2014). The appearance of *U. ingressus* did not always result in the replacement of other cave bear forms. In the two Austrian caves, Ramesch and Gamssulzen, located about 10 km from each other, *U. ingressus* lived side by side with *U. s. eremus* for at least 15,000 years (Hofreiter et al., 2004).

Less is known about the cave bear population that inhabited surroundings of the Western Carpathians and Sudetes. Radio-




Fig. 3. Cave bear (*Ursus spelaeus* s.l.)

A – Bayesian phylogeny of European cave bears based on mtDNA control region sequences. The tree is a phylogram where branch lengths are proportional to amounts of sequence differences; B – spatial distribution of cave bear remains classified as *U. s. spelaeus* and *U. ingressus* (after Popović et al., 2015). Colours indicate mitochondrial DNA lineages of cave bears





carbon dates suggest more or less continuous presence of cave bears in this area (Nadachowski et al., 2008; Wojtal et al., 2015) during the Late Pleistocene. Preliminary results of genetic investigation of specimens from multiple sites in Poland, the Czech Republic, Slovakia and Ukraine revealed that *U. ingressus* was the only form of cave bear present in this part of Europe (Popović et al., 2015; Fig. 3B). It appeared in the Carpathians and Sudetes much earlier than in the Alps. The earliest remains confirmed with morphological and ancient DNA analyses are from Niedźwiedzia Cave in the Sudetes. Bone collagen from one specimen from this cave was dated with the U-Th method to about 80 ka BP (with an accompanying  $^{14}\text{C}$  date of >50 ka  $^{14}\text{C}$  BP) and the other was stratigraphically dated to more than ca. 70 ka BP (Baca et al., 2014). The mtDNA haplotypes of specimens from Niedźwiedzia Cave formed a divergent cluster in phylogenetic trees, which confirms the early separation and expansion of this population (Baca et al., 2012, 2014). Recently, two cave bears with similar mtDNA haplotypes were recorded farther west in Zoolithen Cave (Upper Franconia, Germany) (Stiller et al., 2014). In this cave, remains of *U. spelaeus* were also discovered and interestingly, an opposite replacement was suggested as the estimated age of both *U. ingressus* specimens was older than that of nine *U. spelaeus* individuals (Stiller et al., 2014). These results are, however, based on molecular dating of the remains and should be further confirmed with direct radiocarbon dating.

Another debated issue in cave bear population history is the timing and causes of their extinction. Direct radiocarbon dates indicate that last cave bears went extinct prior to the LGM. Until recently, it was thought that they disappeared from the fossil record quite synchronously in different parts of Europe around 24  $^{14}\text{C}$  ka BP (28 ka BP) at the end of GI-3 (Hofreiter et al., 2002; Pacher and Stuart, 2009; Bocherens et al., 2014). Palaeogenetic analyses showed, however, that the demise of cave bears started ca. 50 ka  $^{14}\text{C}$  BP (Stiller et al., 2010), thus about 25,000 years before their final extinction. It has been proposed that the changing climate was one of the main causes of the cave bear extinction (Baca et al., 2016).

Cave bears are generally considered herbivorous based on their craniodental adaptations (Kurtén, 1976; Mackiewicz et al., 2010; Wiszniowska et al., 2010; van Heteren et al., 2014) and most of the studies of stable isotopes ( $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$ ) from bone and tooth collagen confirm that they were strict vegetarians (Bocherens et al., 1994, 1997; Taboada et al., 1999; Fernández-Mosquera et al., 2001; Münzel et al., 2011; Krajcarz et al., 2016). The climate change which began after GS-3 caused severe transformations in plant communities all around Europe (Helmens, 2014). Vegetation seasons shortened and the availability of high quality plant material, which seems crucial for the survival of cave bears, decreased. Dietary habits of cave bears did not change during their presence for the last 10,000 years in Europe and this ecological niche conservatism may have led to the decline of their populations (Bocherens et al., 2014).

Besides the environmental changes several other factors might have influenced the cave bear populations. There is a substantial evidence of hunting of cave bears by humans (Münzel et al., 2011; Wojtal et al., 2015), as well as competition by them for caves as a shelter (Grayson and Delpech, 2003). Possibly also large carnivores such as cave lion (*Panthera spelaea*) and cave hyena (*Crocuta crocuta spelaea*) hunted hibernating cave bears (Diedrich, 2014). Analysis of whole mitochondrial genomes of bears from several sites in northwestern Spain revealed recently that cave bears, in contrast to the brown bears, exhibited strong fidelity to the caves where they were born. It was suggested that facing growing competition with modern humans and Neanderthals such homing behav-

iour might have contributed to their extinction. Tendency to come back to the same cave for dormancy may prevented colonization of new caves and made cave bears a predictable prey for hunters (Fortes et al., 2016).

Recently several young cave bear specimens were reported in Western Europe. One specimen from the Rochedane site (French Jura), confirmed with ancient DNA as *U. spelaeus*, yielded an AMS date of about 28.5 ka BP (Bocherens et al., 2014). Two specimens from Chiostraccio Cave (Siena, Italy) were dated to ca. 28 and 27 ka BP, respectively (Martini et al., 2014). Stiller et al. (2014) suggested that cave bear populations might have declined from east to west as most of the samples younger than 30 ka BP were found in Western Europe. However, young specimens have also been reported in Eastern Europe (Fig. 4). Two specimens from the Kraków-Częstochowa Upland, from Deszczowa and Komarowa caves, were dated to 28.6 ka BP and confirmed as *U. ingressus* (Popović et al., 2015; Wojtal et al., 2015). Another one from Isabela Textorisova Cave (Velka Fatra Mts., Slovakia) (Sabol et al., 2014) was similarly dated to 28.7 ka BP. The youngest genetically confirmed cave bear specimen so far come from Stajnia Cave in Poland and was dated to around 26.1 ka BP. Baca et al. (2016) gathered 206 radiocarbon dated specimens and following eight approaches estimated the extinction time of *U. spelaeus sensu lato* to between 27.0 and 24.3 ka BP. These findings clearly indicate that the pattern recognized by Stiller et al. (2014) was a result of sampling bias and that the late cave bear survived independently in isolated populations in different parts of Europe, even into the middle of GS-3 stadial. Especially the karst regions may have provided suitable microclimate for long survival of this species (Baca et al., 2016).


The large amount of directly dated and genetically analysed cave bear remains provide insight into cave bear population dynamics in Europe. However, some aspects of the mode of life of cave bears and details of their extinction, evolution and phylogeography still await explanation.

## SAIGA ANTELOPE (SAIGA TATARICA)

Hollow-horned ruminants (Bovidae) are the most diverse family in the order Artiodactyla. More delicately-built species, including the genus Saiga, belong to the Antilopini together with gazelles and other antelopes (Groves and Leslie, 2011). The bizarre-looking saigas are not related to sheep and goats, as used to be thought, but are the sister group of the gazelles. They are non-territorial and nomadic antelopes, gathering in massive herds of many thousands of individuals before migrations. The saiga is a specialized steppe herbivore adapted to flat plains and avoiding rugged terrains. At present it inhabits dry steppes and semi-deserts but during the Late Pleistocene it was widespread in vast areas of Eurasia and North America, belonging to so-called "*Mammuthus-Coelodonta*" faunal complex (Kahlke, 1999, 2014). The number of dispersal events to Europe during the Late Pleistocene was limited by the ecological requirements of this species and by weather conditions. The large thickness of snow cover in winter was the largest restriction on long-distance migrations of saiga.

There is still hot debate whether the saiga represents two or one species. Some palaeontologists distinguish two fossil species: *Saiga borealis* and *S. tatarica* (Baryshnikov and Tikhonov, 1994), others only one species with two subspecies, *S. tatarica borealis* and *S. tatarica tatarica* in spite of some distinct differences found in the skull morphology between both forms






**Fig. 4. Cave bear (*Ursus spelaeus* s.l.)**

**A** – distribution of palaeontological sites with cave bear remains younger than 32 ka BP; **B** – cave bear remains younger than 32 ka BP alongside the NGRIP GICC05  $\delta^{18}\text{O}$  ice core record; specimens genetically determined as *U. spelaeus* are coloured in blue, and *U. ingressus* in green

(Kahlke, 1991; Ratajczak et al., 2016). The current genetic studies also suggest that both fossil and recent saiga should be classified as one species, *Saiga tatarica* (Kholodova et al., 2006; Campos et al., 2010b).

The permanent occurrence of this antelope in steppe areas in Eastern Europe north of the Black Sea, in the Crimea and Dobruja (Dobrogea) in Romania during all of MIS 3 cannot be questioned (Markova et al., 1995; Péan et al., 2013; Ridush et al., 2013). Previously published overviews suggest that the

saiga was also present in Western and Central Europe in MIS 3 (Stewart, 2007; Markova et al., 2010). However, this evidence was not recently confirmed by direct radiocarbon dating (Nadachowski et al., 2016). During MIS 2 and the early part of MIS 1 (the Late Glacial), saiga appeared in Europe in three immigration waves (Fig. 5). The oldest one was probably restricted to Central Europe (e.g., Poland and Czech Republic; Nadachowski et al., 2016) and was temporally limited to the period just before GI-2 (Fig. 5A), i.e. a short warming between



**Fig. 5. Saiga antelope (*Saiga tatarica*)**

**A** – a plot of direct radiocarbon dates of saiga antelope in Europe (Langlais et al., 2015; Nadachowski et al., 2016; Barshay-Szmidt et al., 2016) alongside the NGRIP GICC05  $\delta^{18}\text{O}$  ice core record, calibrated using program OxCal v. 4.2 (Ramsey, 2009); GS – Greenland Stadials, GI – Greenland Interstadials; **B** – spatial distribution of sites with saiga remains in Europe; white circles denote sites with remains dated by archaeological context, while coloured circles indicate sites with radiocarbon-dated remains; colours correspond to three putative waves of migration of saiga into Europe (supplemented and modified after Nadachowski et al., 2016)



23.3 and 22.9 ka BP (Rasmussen et al., 2014). The second, geographically much wider migration started just after the LGM, ca. 19.5 ka BP, and lasted for ca. 3,500–4,000 years to ca. 15.5 ka BP (Nadachowski et al., 2016; Fig. 5). During this time, *Saiga tatarica* was present in the whole of Europe north of the Carpathians, Alps and Pyrenees and reached the Aquitaine Basin and Gascony in SW France. The comparison of a still limited number of direct dates suggests rather migrations of saiga from western Europe to its eastern refugial areas because the dates from France (Langlais et al., 2015; Barshay-Szmidt et al., 2016) are older than those from Germany and Poland (Nadachowski et al., 2016; Fig. 5). This interesting observation should be, however, confirmed by more direct radiocarbon dating. The reduction in range of the saiga in Europe was continued in the Late Glacial (Fig. 5) due to the development of vegetation cover not suitable for this herbivore. However, during cooler episodes of the Late Glacial, between the Bølling and Allerød (GI-1d, former Older Dryas) and within the Allerød (GI-1c2) (Rasmussen et al., 2014) *Saiga tatarica* probably returned to Central Europe, but this re-immigration was restricted only to the area north of the Carpathians (Nadachowski et al., 2016; Fig. 5). It seems that this species was unable to extend its range to the west of Europe during the Younger Dryas (GS-1), the last cold phase of the Pleistocene, even though the climatic and environmental conditions were suitable for this herbivore.

The history of saiga in the European Late Pleistocene, especially the directions of migrations (westward or eastward), needs further investigation basing on new radiocarbon dates and on palaeogenetic studies to explain the complex migration events of this endangered ruminant.

#### COLLARED LEMMING (*DICROSTONYX* SSP.)

Morphological and molecular evidence shows that voles and lemmings create a monophyletic group of rodents placed by most authors within the Arvicolinae subfamily (Chaline et al., 1999). Collared lemmings (*Dicrostonyx* spp.) are cold-adapted animals that are restricted to dry and treeless Arctic tundra environments (Kowalski, 1995). They are a key species in trophic networks in Arctic ecosystems as they are prey for predators such as arctic fox, snow owl and stoat. Four contemporary species are recognized within the genus: *D. torquatus* that inhabits Eurasia from White Sea to Bering Strait, as well as *D. groenlandicus*, *D. richardsoni* and *D. hudsonius* that inhabit North America. These species have been recognized on the basis of phylogenetic analysis of mitochondrial DNA, karyotype diversity and hybridisation experiments (Jarrell and Fregda, 1993; Fedorov and Goropashnaya, 1999). This classification is currently the most commonly accepted one.

At present collared lemmings show nearly circumpolar distribution and do not exceed latitude 65°N, but in the Pleistocene their range was much wider and their remains are known from palaeontological sites throughout Europe and Asia (e.g., Markova et al., 2010; Ponomarev and Puzachenko, 2015). Pleistocene collared lemmings are classified according to the growing complexity of occlusal surface of molar teeth. The oldest ones are known from the Early Pleistocene and are classified as *Predicrostonyx* (Nadachowski, 1992), which was followed by *D. renidens*, *D. meridionalis* and *D. simplicior* in the Mid Pleistocene (Zazhigin, 1980; Smirnov et al., 1986) and *D.*

*gulielmi* in the Late Pleistocene. It was suggested that the transition between *D. gulielmi* and the recent *D. torquatus* took place in the Late Glacial about 15–13 ka BP. Morphometric analyses of molar teeth revealed that during this time, a substantial change in the structure of collared lemming populations took place (Smirnov, 2002). This was further supported by genetic analyses of contemporary collared lemming populations. Phylogeographic analyses revealed the existence of five allopatric populations within the present distribution of the species (Fedorov et al., 1999). Each of the populations was characterized by low genetic diversity, which was interpreted as an effect of the regional population reductions that might have taken place during the warm intervals of the Holocene. Other evidence came from the palaeogenetic analyses of collared lemming remains from the Pymva Shor site in the northern Pre-Urals that revealed the signature of a severe population bottleneck around 14.5 ka BP (Prost et al., 2010). The climate warming during the Bølling-Allerød (GI-1a-c) period in the Late Glacial was accompanied by a northward expansion of forests and supposedly this forced the contraction and isolation of collared lemming populations that resulted in their decline (Fedorov, 1999).

Recently, Palkopoulou et al. (2016) used ancient DNA and radiocarbon dating to reconstruct the Late Pleistocene evolutionary history of collared lemmings. They obtained mtDNA sequences from more than 300 collared lemming specimens from multiple palaeontological sites, and 48 direct radiocarbon AMS dates. Phylogenetic analyses revealed that five distinct populations of collared lemmings, represented by five mitochondrial lineages (EA1–EA5; Fig. 6A), existed in Europe. Each of the populations was widely distributed, ranging from Western France to the Ural Mountains. The radiocarbon dating revealed that subsequent populations succeeded each other through time during the last ca. 50 ka. This was interpreted as a series of collared lemming population extinctions and recolonizations that took place across the whole European continent. The phylogenetic pattern found indicated that subsequent recolonizations proceeded in an east to west direction from a hypothetical refugial area in northeastern Siberia. Dates for the two earliest populations (EA1 and EA2) are close to radiocarbon dating limits but it seems that both were already present in Europe prior to 50 ka BP. The EA1 population vanished around 50 ka BP and the EA2 population around 42.3 ka BP. Population EA3 appeared in Europe ca. 32 ka BP and disappeared 22.8 ka BP and was followed by population EA4, dated to a short period between 22.2 and 20.5 ka BP. Population EA5 emerged ca. 20.3 ka BP and disappeared from Europe ca. 14.5 ka BP. All Holocene and modern collared lemmings from Siberia belong to this EA5 mtDNA lineage (Fig. 6B).

It has been postulated that the population turnovers that took place across such a large geographical range had to be driven by environmental or ecological changes in the steppe-tundra ecosystem. The dating of the two earliest populations (EA1 and EA2) is not very precise and the connection between their demise and climate changes remains unclear. The spread of population EA3, which seems reliably dated, started just after GI-5 and might be associated with climate cooling that begun with the onset of GS-5. Surprisingly, the two following population turnovers, i.e. the replacement of population EA3 by EA4 and EA4 by EA5, took place within the LGM. During this period, arctic tundra environments preferred by the collared lemmings spread over large areas of Eurasia (Tarasov et al., 2000). Thus, the causes of these replace-






Fig. 6. Collared lemming (*Dicrostonyx* ssp.)

**A** – Bayesian phylogeny of Eurasian collared lemmings based on mtDNA cytochrome b sequences. The tree is a chronogram where branch lengths denote time elapsed since divergence and the positions of tips correspond to calibrated radiocarbon sample age. Colours indicate different mtDNA lineages; **B** – spatial distribution of palaeontological sites with collared lemming in different periods. Colours indicate different mtDNA lineages (modified after [Palkopoulou et al., 2016](#)) and white circles denote sites with remains dated by archaeological context

ments remain unclear; however, it has been shown that they coincide with brief warming periods recorded in the high resolution palynological record from Lago Grande di Montichio, Italy ([Palkopoulou et al., 2016](#)).

## DISCUSSION AND CONCLUSIONS

High-resolution reconstructions of migration and extinction events during the Late Pleistocene obtained in recent years show that species responded to climate changes according to their individual adaptations and there are no grounds for considering species as faunal complexes responding simultaneously and in the same manner ([Hofreiter and Stewart, 2009](#);

[Stewart et al., 2010](#)). This is well-illustrated by the comparison of three species reviewed here: woolly mammoth, saiga antelope and collared lemming. These are all considered as members of the “*Mammuthus–Coleodonta*” faunal complex ([Kahlke, 1999, 2014](#)), but their Late Pleistocene histories in Europe differ substantially. During the last 50 ka mammoths were present in Europe more or less continuously, with a possible range contraction only during the LGM. The first confirmed appearance of the saiga antelope was most probably related with the brief warming GI-2 dated to ca. 23 ka BP. However, only after the LGM was important extension of its range in Europe observed. Collared lemmings most probably occurred, as did mammoths, continuously since 50 ka, but their population turnover was much more intense and happened more often than in the case of previous species. Interestingly,








Fig. 7. Population turnovers and extinctions in Europe based on radiocarbon dating

**A** – woolly mammoth, colours indicate different mtDNA clades (only specimens with genetic information are shown); **B** – woolly mammoth from Europe, without differentiation into the clades, source of the dates is as in [Figure 2](#) supplemented with data from [Nadachowski et al. \(2011\)](#) and [Ukkonen et al. \(2011\)](#); **C** – cave bear, blue – *U. s. spelaeus*, green – *U. ingressus*; **D** – saiga antelope, colours indicate three putative expansion waves; **E** – collared lemmings, colours indicate different populations (EA1–EA5). MIS – Marine Isotope Stages, GI – Greenland Interstadials; GS – Greenland Stadials; LGM – Last Glacial Maximum (after [Mix et al., 2001](#)); vertical stripes indicate timing of population turnovers or extinctions

along with these apparent differences, some similarities are visible. It seems that most of the observed events are grouped around three periods. The first one covers the GI-5 warming and the onset of the GS-5 cooling around 33–31 ka BP, the second one partly overlaps the maximum extent of the Scandinavian Ice Sheet during the LGM at ca. 22–19 ka BP and the last one is the abrupt climate warming during the Late Glacial ca. at 15–13 ka BP ([Fig. 7](#)). In agreement with that, the climate during the LGM (as defined by [Mix et al., 2001](#)) and the warming during the Late Glacial have been previously recognized as major factors that substantially influenced the distribution of many mammal species in Europe.

During the LGM, numerous species withdrew from Europe or restricted their ranges into so-called southern glacial refugia ([Hewitt, 2004](#); [Stewart and Cooper, 2008](#)). This concerns most of temperate species such as red deer (*Cervus elaphus*; [Sommer et al., 2008](#); [Skog et al., 2009](#)), roe deer (*Capreolus*

*capreolus*; [Sommer et al., 2009](#); [Sommer and Zachos, 2009](#)), brown hare (*Lepus europaeus*; [Stamatis et al., 2009](#)) as well as cold-adapted ones, such as mammoths and saiga antelopes ([Fig. 7](#)). During the maximum extent of the Scandinavian Ice Sheet, environmental conditions in Central and northwestern Europe were extremely harsh, with continuous permafrost covering most of Poland, Germany, Belgium, Netherlands and northern France ([Huijzer and Vandenberghe, 1998](#)). [Ukkonen et al. \(2011\)](#) suggested that during that time productivity of environments was extremely low, which forced the contraction of even cold-adapted species. However, the harsh conditions during the LGM do not seem to have led to many extinction events. [Cooper et al. \(2015\)](#) pointed out that the gradual climate deterioration prior to and the relative stability during the LGM allowed mammalian populations to retreat or to adapt to the changing environment. Two Europe-wide replacements of *Dicrostonyx* populations observed between 23 and 20 ka BP suggest, how-



ever, that some short-term environmental changes must have taken place at that time.

The Late Glacial period was marked by several climate oscillations, with the warm Bølling-Allerød Interstadial (GI-1) interrupted by the Older Dryas (GI-1d) and followed by the Younger Dryas (GS-1) cold phases (Steffensen et al., 2008). These rapid changes are linked to the extinction of the cave lion (*Panthera spelaea*; Stuart and Lister, 2011) and woolly rhinoceros (*Coelodonta antiquitatis*; Stuart and Lister, 2012).

The period 33–31 ka BP has been suggested only recently from the analyses of collared lemmings. Palkopoulou et al. (2016) noticed that the appearance of one of the *Dicrostonyx* populations in Europe at the onset of GS-5, ca. 31 ka BP, coincides broadly with the replacement of the endemic European mammoth population by another one from Asia and with the replacement of *U. s. spelaeus* by *U. ingressus* in Central Europe (Fig. 7). Furthermore, recent new radiocarbon AMS data showed that the last populations of spotted hyena (*Crocuta crocuta*) also became extinct ca. 31 ka BP (Stuart and Lister, 2014). The recent worldwide survey of extinction and popula-

tion replacements revealed that the timing of these events was significantly correlated with rapid and high amplitude climate changes during interstadials (Cooper et al., 2015). The concentration of such events around GI-5 fits well to this hypothesis as this interstadial represents one of the most instantaneous climate changes during the last glaciation.

All four case studies presented here unveil the Late Pleistocene communities as complex and highly dynamic, characterized by geographic range shifts, migrations, replacements and local extinctions, which seem to be common phenomena rather than exceptions. Cooper et al. (2015) note that such processes might be essential for maintaining ecosystem stability in periods of high climate variability.

**Acknowledgements.** The work related to cave bears was supported by the Polish National Science Centre grant no. 2012/07/B/NZ8/02845. We are grateful to the reviewers L. Lõugas and T. van Kolfschoten for their constructive comments and to M. Molak for her suggestions and help with proof-reading and correction of the manuscript.

## REFERENCES

- Alroy, J., 2001. A multispecies overkill simulation of the end-Pleistocene megafaunal mass extinction. *Science*, **292**: 1893–1896.
- Baca, M., Stankovic, A., Stefaniak, K., Marciszak, A., Nadachowski, A., Węgleński, P., Mackiewicz, P., 2012. Genetic analysis of cave bear specimens from Niedźwiedzia cave. *Palaeontologica Electronica*, **15**: 16.
- Baca, M., Mackiewicz, P., Stankovic, A., Popović, D., Stefaniak, K., Czarnogórska, K., Nadachowski, A., Gąsiorowski, M., Hercman, H., Weglenski, P., 2014. Ancient DNA and dating of cave bear remains from Niedźwiedzia Cave suggest early appearance of *Ursus ingressus* in Sudetes. *Quaternary International*, **339–340**: 217–223.
- Baca, M., Popović, D., Stefaniak, K., Marciszak, A., Urbanowski, M., Nadachowski, A., Mackiewicz, P., 2016. Retreat and extinction of the Late Pleistocene cave bear (*Ursus spelaeus* sensu lato). *The Science of Nature*, **103**: 92.
- Barnosky, A.D., Koch, P.L., Feranec, R.S., Wing, S.L., Shabel, A.B., 2004. Assessing the causes of late Pleistocene extinctions on the continents. *Science*, **306**: 70–75.
- Barshay-Szmidt, C., Costamagno, S., Henry-Gambier, D., Laroulandie, V., Pétilion J.-M., Boudadi-Maligne, M., Kuntz, D., Langlais, M., Mallye, J.-B., 2016. New extensive focused AMS <sup>14</sup>C dating of the Middle and Upper Magdalenian of the western Aquitaine/Pyrenean region of France (ca. 19–14 ka cal BP): Proposing a new model for its chronological phases and for the timing of occupation. *Quaternary International*, **414**: 62–91.
- Baryshnikov, G., 1998. Cave bears from the Paleolithic of the Greater Caucasus. *Illinois State Museum Scientific Papers*, **27**: 69–118.
- Baryshnikov, G., Tikhonov, A., 1994. Notes on skulls of Pleistocene saiga of northern Eurasia. *Historical Biology*, **8**: 209–234.
- Bocherens, H., Fizet, M., Mariotti, A., 1994. Diet, physiology and ecology of fossil mammals as inferred from stable carbon and nitrogen isotope biogeochemistry: implications for Pleistocene bears. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **107**: 213–225.
- Bocherens, H., Billiou, D., Patou-Mathis, M., Bonjean, D., Otte, M., Mariotti, A., 1997. Paleobiological implications of the isotopic signatures (<sup>13</sup>C, <sup>15</sup>N) of fossil mammal collagen in Scladina Cave (Sclayn, Belgium). *Quaternary Research*, **48**: 370–380.
- Bocherens, H., Bridault, A., Drucker, D.G., Hofreiter, M., Münzel, S.C., Stiller, M., van der Plicht J., 2014. The last of its kind? Radiocarbon, ancient DNA and stable isotope evidence from a late cave bear (*Ursus spelaeus* ROSEN-MÜLLER, 1794) from Rochedane (France). *Quaternary International*, **339–340**: 179–188.
- Campos, P.F., Willerslev, E., Sher, A., Orlando, L., Axelsson, E., Tikhonov, A., Aaris-Sørensen, K., Greenwood, A.D., Kahlke R.D., Kosintsev, P., Krakhmalnaya, T., Kuznetsova, T., Lemey, P., MacPhee, R., Norris, C.A., Shepherd, K., Suchard, M.A., Zazula, G.D., Shapiro, B., Gilbert, T.P., 2010a. Ancient DNA analyses exclude humans as the driving force behind late Pleistocene musk ox (*Ovibos moschatus*) population dynamics. *Proceedings of the National Academy of Sciences*, **107**: 5675–5680.
- Campos, P.F., Kristensen, T., Orlando, L., Sher, A., Kholodova, M.V., Götherström, A., Hofreiter, M., Drucker, D.G., Kosintsev, P., Tikhonov, A., Baryshnikov, G.F., Willerslev, E., Gilbert, M.T.P., 2010b. Ancient DNA sequences point to large loss of mitochondrial genetic diversity in the saiga antelope (*Saiga tatarica*) since the Pleistocene. *Molecular Ecology*, **19**: 4863–4875.
- Chaline, J., Brunet-Lecomte, P., Montuire, S., Viriot, L., Courant, F., 1999. Anatomy of the arvicoline radiation (Rodentia): palaeogeographical, palaeoecological history and evolutionary data. *Annales Zoologici Fennici*, **36**: 239–267.
- Cooper, A., Turney, C., Hughen, K.A., Barry, W., McDonald, H.G., Bradshaw, C.J.A., 2015. Abrupt warming events drove Late Pleistocene Holarctic megafaunal turnover. *Science Express*, **349**: 1–8.
- Dabney, J., Knapp, M., Glocke, I., Gansauge, M.T., Weihmann, A., Nickel, B., Valdiosera, C., Garcia, N., Pääbo, S., Arsuaga, J.L., Meyer, M., 2013. Complete mitochondrial genome sequence of a Middle Pleistocene cave bear reconstructed from ultrashort DNA fragments. *Proceedings of the National Academy of Sciences*, **110**: 15758–15763.
- Debruyne, R., Chu, G., King, C.E., Bos, K., Kuch, M., Schwarz, C., Szpak, P., Gröcke, D.R., Matheus, P., Zazula, G., Guthrie, D., Froese, D., Buigues, B., de Marliave, C., Flemming, C.,



- Poinar, D., Fisher, D., Southon, J., Tikhonov, A.N., MacPhee, R.D.E., Poinar, H.N., 2008. Out of America: ancient DNA evidence for a new world origin of Late Quaternary woolly mammoths. *Current Biology*, **18**: 1320–1326.
- Diedrich, C.G., 2014. Palaeopopulations of Late Pleistocene top predators in Europe: Ice Age spotted hyenas and steppe lions in battle and competition about prey. *Paleontology Journal*, **2014**: 1–34.
- Fedorov, V.B., 1999. Contrasting mitochondrial DNA diversity estimates in two sympatric genera of Arctic lemmings (*Dicrostonyx*: *Lemmus*) indicate different responses to Quaternary environmental fluctuations. *Proceedings of the Royal Society of London B: Biological Sciences*, **266**: 621–626.
- Fedorov, V., Goropashnaya, A., 1999. The importance of ice ages in diversification of arctic collared lemmings (*Dicrostonyx*): evidence from the mitochondrial cytochrome b region. *Hereditas*, **130**: 301–307.
- Fedorov, V.B., Fredga, K., Jarrell, G.H., 1999. Mitochondrial DNA variation and the evolutionary history of chromosome races of collared lemmings (*Dicrostonyx*) in the Eurasian Arctic. *Journal of Evolutionary Biology*, **12**: 134–145.
- Fernández-Mosquera, D., Vila-Taboada, M., Grandal-d'Anglade, A., 2001. Stable isotopes data ( $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$ ) from the cave bear (*Ursus spelaeus*): a new approach to its palaeoenvironment and dormancy. *Proceedings of the Royal Society B: Biological Sciences*, **268**: 1159–1164.
- Fortes, G.G., Grandal-d'Anglade, A., Kolbe, B., Fernandes, D., Meleg, I.N., García-Vázquez, A., Pinto-Llona, A.C., Constantín, S., de Torres, T.J., Ortiz, J.E., Frischauf, C., Rabeder, G., Hofreiter, M., Barlow, A., 2016. Ancient DNA reveals differences in behaviour and sociality between brown bears and extinct cave bears. *Molecular Ecology*, **25**: 4907–4918.
- Gilbert, M., Drautz, D., 2008. Intraspecific phylogenetic analysis of Siberian woolly mammoths using complete mitochondrial genomes. *Proceedings of the National Academy of Sciences*, **105**: 8327–8332.
- Graham, R.W., Lundelius, E.L., Graham, M.A., Schroeder, E.K., Toomey, R.S., Anderson, E., Barnosky, A.D., Burns, J.A., Churcher, C.S., Grayson, D.K., Guthrie, D., Harington, C.R., Jefferson, G.T., Martin, L.D., McDonald, G., Morlan, R.E., Semken, Jr., H.A., Webb, S.D., Werdelin, L., Wilson, M.C., 1996. Spatial response of mammals to Late Quaternary environmental fluctuations. *Science*, **272**: 1601–1606.
- Grayson, D.K., Delpech, F., 2003. Ungulates and the Middle-to-Upper Paleolithic transition at Grotte XVI (Dordogne, France). *Journal of Archaeological Science*, **30**: 1633–1648.
- Groves, C.P., Leslie, D.E. Jr., 2011. Family Bovidae (hollow-horned ruminants). In: *Handbook of the Mammals of the World, 2 Hoofed Mammals* (eds. D.E. Wilson and R.A. Mittermeier): 444–779. Lynx Edicions, Barcelona.
- Hewitt, G.M., 2004. Genetic consequences of climatic oscillations in the Quaternary. *Philosophical transactions of the Royal Society of London. Series B, Biological Sciences*, **359**: 183–195.
- Helmens, K.F., 2014. The last interglacial-glacial cycle (MIS 5-2) re-examined based on long proxy records from central and northern Europe. *Quaternary Science Reviews*, **86**: 115–123.
- Hofreiter, M., Stewart, J., 2009. Ecological change, range fluctuations and population dynamics during the Pleistocene. *Current Biology*, **19**: R584–R594.
- Hofreiter, M., Capelli, C., Krings, M., Waits, L., Conard, N., Münzel, S., Rabeder, G., Nagel, D., Paunovic, M., Jambresić, G., Meyer, S., Weiss, G., Pääbo, S., 2002. Ancient DNA analyses reveal high mitochondrial DNA sequence diversity and parallel morphological evolution of Late Pleistocene cave bears. *Molecular Biology and Evolution*, **19**: 1244–1250.
- Hofreiter, M., Rabeder, G., Jaenicke-Després, V., Withalm, G., Nagel, D., Paunovic, M., Jambresić, G., Pääbo, S., 2004. Evidence for reproductive isolation between Cave Bear populations. *Current Biology*, **14**: 40–43.
- Hofreiter, M., Münzel, S., Conard, N.J., Pollack, J., Slatkin, M., Weiss, G., Pääbo, S., 2007. Sudden replacement of cave bear mitochondrial DNA in the late Pleistocene. *Current Biology*, **17**: R122–3.
- Horn, S., Prost, S., Stiller, M., Makowiecki, D., Kuznetsova, T., Benecke, N., Pucher, E., Hufthammer, A.K., Schouwenburg, C., Shapiro, B., Hofreiter, M., 2014. Ancient mitochondrial DNA and the genetic history of Eurasian beaver (*Castor fiber*) in Europe. *Molecular Ecology*, **23**: 1717–1729.
- Huijzer, B., Vandenberghe, J., 1998. Climatic reconstruction of the Weichselian Pleniglacial in northwestern and central Europe. *Journal of Quaternary Science*, **13**: 391–417.
- Jarrell, G., Fredga, K., 1993. How many kinds of lemmings? A taxonomic overview. In: *The Biology of Lemmings* (eds. N. Stenseth and R. Ims): 45–57. Academic Press, London.
- Kahlke, R.D., 1991. Pleistocene distributional and evolutionary history of the genus *Saiga* Gray, 1843 (Mammalia, Artiodactyla, Bovidae) in the Palaearctic. *Vertebrata Palasiatica*, **10**: 314–322.
- Kahlke, R.D., 1999. The History of the Origin, Evolution and Dispersal of the Late Pleistocene *Mammuthus-Coelodonta* Faunal Complex in Eurasia (Large Mammals). Fenske Companies: Rapid City, SD.
- Kahlke, R.D., 2014. The origin of Eurasian Mammoth faunas (*Mammuthus-Coelodonta* Faunal Complex). *Quaternary Science Reviews*, **96**: 32–49.
- Kahlke, R.D., 2015. The maximum geographic extension of Late Pleistocene *Mammuthus primigenius* (Proboscidea, Mammalia) and its limiting factors. *Quaternary International*, **379**: 147–154.
- Kholodova, M.V., Milner-Gulland, E.J., Easton, A.J., Amgalan, L., Arylov, Iu.A., Bekenov, A., Grachev, Iu.A., Lushchekina, A.A., Ryder, O., 2006. Mitochondrial DNA variation and population structure of the critically endangered saiga antelope *Saiga tatarica*. *Oryx*, **40**: 103–107.
- Knapp, M., Rohland, N., Weinstock, J., Baryshnikov, G.F., Sher, A., Nagel, D., Rabeder, G., Pinhasi, R., Schmidt, H.A., Hofreiter, M., 2009. First DNA sequences from Asian cave bear fossils reveal deep divergences and complex phylogeographic patterns. *Molecular Ecology*, **18**: 1225–1238.
- Koch, P.L., Barnosky, A.D., 2006. Late Quaternary extinctions: state of the debate. *Annual Review of Ecology, Evolution, and Systematics*, **37**: 215–250.
- Kowalski, K., 1995. Lemmings (Mammalia, Rodentia) as indicators of temperature and humidity in the European Quaternary. *Acta Zoologica Cracoviensia*, **38**: 85–94.
- Krajcarz, M., Pacher, M., Krajcarz, M.T., Laughlan, L., Rabeder, G., Sabol, M., Wojtal, P., Bocherens, H., 2016. Isotopic variability of cave bears ( $\delta^{15}\text{N}$ ,  $\delta^{13}\text{C}$ ) across Europe during MIS 3. *Quaternary Science Review*, **131**: 51–72.
- Kurtén, B., 1976. The cave bear story: life and death of a vanished animal. Columbia University Press.
- Langlais, M., Laroulandie, V., Costamagno, S., Pétilion, J.-M., Mallye, J.-B., Lacrampe-Cuyaubère, F., Boudadi-Maligne, M., Barshay-Szmidt, C., Masset, C., Pubert, E., Rendu, W., Lenoir, M., 2015. Premiers temps du Magdalénien en Gironde Réévaluation des fouilles Trécolle à Saint-Germain-le-Rivière (France). *Bulletin de la Société Préhistorique Française*, **112**: 5–58.
- Lister, A.M., Sher, A.V., 2001. The origin and evolution of the woolly mammoth. *Science*, **294**: 1094–1097.
- Lister, A.M., Sher, A.V., van Essen, H., Wei, G., 2005. The pattern and process of mammoth evolution in Eurasia. *Quaternary International*, **126–128**: 49–64.
- Lister, A.M., Stuart, A.J., 2008. The impact of climate change on large mammal distribution and extinction: evidence from the last glacial/interglacial transition. *Comptes Rendus Geoscience*, **340**: 615–620.
- Lorenzen, E.D., Nogués-Bravo, D., Orlando, L., Weinstock, J., Binladen, J., Marske, K.A., Ugan, A., Borregaard, M.K., Gilbert, M.T.P., Nielsen, R., Ho, S.Y.W., Goebel, T., Graf, K.E.,



- Byers, D., Stenderup, J.T., Rasmussen, M., Campos, P.F., Leonard, J.A., Koepfli, K.P., Froese, D., Zazula, G., Stafford, T.W.Jr., Aaris-Sørensen, K., Batra, P., Haywood, A.M., Singarayer, J.S., Valdes, P.J., Boeskorov, G., Burns, J.A., Davydov, S.P., Haile, J., Jenkins, D.L., Kosintsev, P., Kuznetsova, T., Lai, X., Martin, L.D., McDonald, G.H., Mol, D., Meldgaard, M., Munch, K., Stephan, E., Sablin, M., Sommer, R.S., Sipko, T., Scott, E., Suchard, M.A., Tikhonov, A., Willerslev, R., Wayne, R.K., Cooper, A., Hofreiter, M., Sher, A., Shapiro, B., Rahbek, C., Willerslev, E., 2011. Species-specific responses of Late Quaternary megafauna to climate and humans. *Nature*, **479**: 359–364.
- Lõugas, L., Ukkonen, P., Jungner, H., 2002. Dating the extinction of European mammoths: new evidence from Estonia. *Quaternary Science Reviews*, **21**: 1347–1354.
- Mackiewicz, P., Wiszniowska, T., Olejniczak, A.J., Stefaniak, K., Socha, P., Nadachowski, A., 2010. Analysis of dental enamel thickness in bears with special attention to *Ursus spelaeus* and *U. wenzensis* (= *minimum*) in comparison to selected representatives of mammals. In: *Morphology and Systematics of Fossil Vertebrates* (ed. D. Nowakowski): 60–77: DN Publisher, Wrocław.
- MacPhee, R.D.E., Tikhonov, A.N., Mol, D., de Marliave, C., van der Plicht, H., Greenwood, A.D., Flemming, C., Agenbroad, L., 2002. Radiocarbon chronologies and extinction dynamics of the late Quaternary mammalian megafauna of the Taimyr Peninsula, Russian Federation. *Journal of Archaeological Science*, **29**: 1017–1042.
- Markova, A.K., van Kolfschoten, T. eds., 2008. *Evolution of European Ecosystems During Pleistocene–Holocene Transition 24–8 kyr BP* (in Russian with English summary). KMK Scientific Press, Moscow.
- Markova, A.K., Smirnov, N.G., Kozharinov, A.V., Kazantseva, N.E., Simakova, A.N., Kitaev L.M., 1995. Late Pleistocene distribution and diversity of mammals in Northern Eurasia. *Paleontologia i Evolució*, **28–29**: 5–143.
- Markova, A.K., Puzachenko, A.Y., van Kolfschoten, T., 2010. The North Eurasian mammal assemblages during the end of MIS 3 (Brianskian–Late Karginian–Denekamp Interstadial). *Quaternary International*, **212**: 149–158.
- Markova, A.K., Puzachenko, A.Y., van Kolfschoten, T., van der Plicht, J., Ponomarev, D.V., 2013. New data on changes in the European distribution of the mammoth and the woolly rhinoceros during the second half of the Late Pleistocene and the early Holocene. *Quaternary International*, **292**: 4–14.
- Martini, I., Coltorti, M., Mazza, P.P., Rustioni, M., Sandrelli, F., 2014. The latest *Ursus spelaeus* in Italy, a new contribution to the extinction chronology of the cave bear. *Quaternary Research*, **81**: 117–124.
- Maschenko, E., 2002. Individual development, biology and evolution of the woolly mammoth. *Cranium*, **19**: 4–120.
- Mix, A.C., Bard, E., Schneider, R., 2001. Environmental processes of the ice age: land, oceans, glaciers (EPILOG). *Quaternary Science Reviews*, **20**: 627–657.
- Münzel, S.C., Stiller, M., Hofreiter, M., Mitnik, A., Conard, N.J., Bocherens, H., 2011. Pleistocene bears in the Swabian Jura (Germany): genetic replacement, ecological displacement, extinctions and survival. *Quaternary International*, **245**: 1–13.
- Nadachowski, A., 1992. Early Pleistocene *Predicrostonyx* (Rodentia, Mammalia) from Poland. *Acta Zoologica Cracoviensia*, **35**: 203–216.
- Nadachowski, A., Lipecki, G., Stefaniak, K., Lorenc, M., Wojtal, P., 2008. Extinction of the cave bear (*Ursus spelaeus*) in Central Europe (in Polish and English summary). In: *Archaeozoological research in Poland and Middle-East Europe, data – methods – interpretations* (eds. D. Makowiecki, R. Ablamowicz, D. Ablamowicz, K. Smiarowski and M. Makohonienko): 76–78. Bogucki Wydawnictwo Naukowe, Poznań.
- Nadachowski, A., Lipecki, G., Wojtal, P., Miękina, B., 2011. Radiocarbon chronology of woolly mammoth (*Mammuthus primigenius*) from Poland. *Quaternary International*, **245**: 186–192.
- Nadachowski, A., Lipecki, G., Ratajczak, U., Stefaniak, K., Wojtal, P., 2016. Dispersal events of the saiga antelope (*Saiga tatarica*) in Central Europe in response to the climatic fluctuations in MIS 2 and the early part of MIS 1. *Quaternary International*, **420**: 357–362.
- Pacher, M., Stuart, A.J., 2009. Extinction chronology and palaeobiology of the cave bear (*Ursus spelaeus*). *Boreas*, **38**: 189–206.
- Palkopoulou, E., Dalen, L., Lister, A.M., Vartanyan, S., Sablin, M., Sher, A., Nystrom Edmark, V., Barnes, I., Germonpre, M., Barnes, I., Thomas, J.A., 2013. Holarctic genetic structure and range dynamics in the woolly mammoth. *Proceedings of the Royal Society of London B: Biological Sciences*, **280**: 20131910.
- Palkopoulou, E., Mallick, S., Skoglund, P., Enk, J., Rohland, N., Li, H., Omrak, A., Vartanyan, S., Poinar, H., Götherström, A., Reich, D., Dalén, L., 2015. Complete genomes reveal signatures of demographic and genetic declines in the Woolly Mammoth. *Current Biology*, **25**: 1–6.
- Palkopoulou, E., Baca, M., Abramson, N.I., Sablin, M., Socha, P., Nadachowski, A., Prost, S., Germonpre, M., Kosintsev, P., Smirnov, N.G., Vartanyan, S., Ponomarev, D., Nystrom, J., Nikolskiy, P., Jass, C.N., Litvinov, Y.N., Kalthoff, D.C., Grigoriev, S., Fadeeva, T., Douka, A., Higham, T.F.G., Ersmark, E., Pitulko, V., Pavlova, E., Stewart, J.R., Weglenski, P., Stankovic, A., Dalen, L., 2016. Synchronous genetic turnovers across Western Eurasia in Late Pleistocene collared lemmings. *Global Change Biology*, **22**: 1710–1721.
- Pawłowska, K., 2015. Elephantids from Pleistocene Poland: state of knowledge. *Quaternary International*, **379**: 89–105.
- Péan, S., Puaud, S., Crépin, L., Prat, S., Quiles, A., van der Plicht, J., Valladas, H., Stuart, A.J., Drucker, D.G., Patou-Mathis, M., Lanoë, F., Yanevich, A., 2013. The Middle to Upper Paleolithic sequence of Buran-Kaya III (Crimea, Ukraine): new stratigraphic, paleoenvironmental, and chronological results. *Radiocarbon*, **55**: 1454–1469.
- Ponomarev, D., Puzachenko, A., 2015. Evolution of occlusal shape of the first and second upper molars of Middle-Pleistocene collared lemmings (*Dicrostonyx*, Arvicolinae, Rodentia) in northeast European Russia. *Boreas*, **44**: 741–759.
- Popović, D., Baca, M., Stefaniak, K., Lipecki, G., Nadachowski, A., Marciszak, A., Sabol, M., Ridush, B., Robličková, M., Káňa, V., Obada, T., Stankovic, A., Mackiewicz, P., 2015. Genetic studies of cave bear (*Ursus ingressus*) populations from Central and Eastern Europe. In: *21st International Cave Bear Symposium, Leiden, Neatherlands. Program and Abstracts* pp. 28, WPZ, Leiden.
- Prescott, G.W., Williams, D.R., Balmford, A., Green, R.E., Manica, A., 2012. Quantitative global analysis of the role of climate and people in explaining late Quaternary megafaunal extinctions. *Proceedings of the National Academy of Sciences*, **109**: 4527–4531.
- Prost, S., Smirnov, N., Fedorov, V.B., Sommer, R.S., Stiller, M., Nagel, D., Knapp, M., Hofreiter, M., 2010. Influence of climate warming on arctic mammals? New insights from ancient DNA studies of the collared lemming *Dicrostonyx torquatus*. *PLoS One*, **5**: e10447.
- Rabeder, G., Hofreiter, M., 2004. Der neue Stammbaum der alpinen Höhlenbären. *Die Höhle*, **55**: 1–19.
- Rabeder, G., Hofreiter, M., Nagel, D., Withalm, G., 2004a. New taxa of Alpine cave bears (Ursidae, Carnivora). *Cahiers Scientifiques-Muséum d'Histoire Naturelle de Lyon*, **HS 2**: 49–67.
- Rabeder, G., Hofreiter, M., Withalm, G., 2004b. The systematic position of the Cave Bear from Potočka zijalka (Slovenia). *Mitteilungen der Kommission für Quartärforschung der Österreichischen Akademie der Wissenschaften*, **13**: 197–200.





- Ramsey, C.B., 2009. Bayesian analysis of radiocarbon dates. *Radiocarbon*, **51**: 337–360.
- Rasmussen, S.O., Bigler, M., Blockley, S.P., Blunier, T., Buchardt, S.L., Clausen, H.B., Cvijanovic, I., Dahl-Jensen, D., Johnsen, S.J., Fischer, H., Gkinis, V., Guillevic, M., Hoek, W.Z., Lowe, J.J., Pedro, J.B., Popp, T., Seierstad, I.K., Steffensen, J.P., Svensson, A.M., Vallelonga, P., Vinther, B.M., Walker, M.J.C., Wheatley, J.J., Winstrup, M., 2014. A stratigraphic framework for abrupt climatic changes during the Last Glacial period based on three synchronized Greenland ice-core records: refining and extending the INTIMATE event stratigraphy. *Quaternary Science Reviews*, **106**: 14–28.
- Ratajczak, U., Shpansky, A.V., Malikov, D.G., Stefaniak, K., Nadachowski, A., Wojtal, P., Ridush, B., Krakhmalnaya, T.V., Stepanchuk, V., Mackiewicz, P., 2016. Quaternary skulls of the saiga from eastern Europe and Siberia: *Saiga borealis* versus *Saiga tatarica* – one species or two? *Quaternary International*, **420**: 329–347.
- Reimer, P.J., Bard, E., Bayliss, A., Beck, J.W., Blackwell, P.G., Ramsey, B.C., Buck, C.E., Cheng, H., Edwards, R.L., Friedrich, M., Grootes, P.M., Guilderson, T.P., Hafflidason, H., Hajdas, I., Hatté, C., Heaton, T.J., Hoffmann, D.L., Hogg, A.G., Hughen, K.A., Kaiser, K.F., Kromer, B., Manning, S.W., Niu, M., Reimer, R.W., Richards, D.A., Scott, E.M., Southon, J.R., Staff, R.A., Turney, C.S.M., van der Plicht, J., 2013. IntCal13 and Marine13 radiocarbon age calibration curves, 0–50,000 years cal BP. *Radiocarbon*, **55**: 1869–1887.
- Ridush, B., Stefaniak, K., Socha, P., Proskurnyak, Y., Marciszak, A., Vremir, M., Nadachowski, A., 2013. Emine-Bair Khosar Cave in the Crimea, a huge bone accumulation of Late Pleistocene fauna. *Quaternary International*, **284**: 151–160.
- Sabol, M., Bendík, A., Grivalský, M.M., Lizák, J., Michlík, I., 2014. Latest and highest fossil record of cave bears (*Ursus ex gr. spelaeus*) in Slovakian Western Carpathians. *Quaternary International*, **339–340**: 189–196.
- Sandom, C., Faurby, S., Sandel, B., Svenning, J.C., 2014. Global late Quaternary megafauna extinctions linked to humans, not climate change. *Proceedings of the Royal Society of London B: Biological Sciences*, **281**: 20133254.
- Skog, A., Zachos, F.E., Rueness, E.K., Feulner, P.G.D., Mysterud, A., Langvatn, R., Lorenzini, R., Hmwe, S.S., Lehoczy, I., Hartl, G.B., Stenseth, N.C., Jakobsen, K.S., 2009. Phylogeography of red deer (*Cervus elaphus*) in Europe. *Journal of Biogeography*, **36**: 66–77.
- Smirnov, N.G., 2002. The rate of morphological changes in populations as a characteristic of the chronological structure of the evolutionary process. In: *Populations, Communities, Evolution: Novoe Znanie*, Kazan: 190–210.
- Smirnov, N.G., Bolshakov, V.N., Borodin, A.V., 1986. Pleistocene rodents of northern Siberia (in Russian). Nauka, Moscow.
- Sommer, R.S., Nadachowski, A., 2006. Glacial refugia of mammals in Europe: evidence from fossil records. *Mammal Review*, **36**: 251–265.
- Sommer, R.S., Zachos, F.E., 2009. Fossil evidence and phylogeography of temperate species: “Glacial refugia” and post-glacial recolonization. *Journal of Biogeography*, **36**: 2013–2020.
- Sommer, R.S., Zachos, F.E., Street, M., Jöris, O., Skog, A., Benecke, N., 2008. Late Quaternary distribution dynamics and phylogeography of the red deer (*Cervus elaphus*) in Europe. *Quaternary Science Reviews*, **27**: 714–733.
- Sommer, R.S., Fahlke, J.M., Schmölcke, U., Benecke, N., Zachos, F.E., 2009. Quaternary history of the European roe deer *Capreolus capreolus*. *Mammal Review*, **39**: 1–16.
- Sommer, R.S., Kalbe, J., Ekström, J., Benecke, N., Ronnie, L., 2014. Range dynamics of the reindeer in Europe during the last 25,000 years. *Journal of Biogeography*, **41**: 298–306.
- Stamatis, C., Suchentrunk, F., Moutou, K.A., Giacometti, M., Haerer, G., Djan, M., Vapa, L., Vukovic, M., Tvrtković, N., Sert, H., Alves, P.C., Mamuris, Z., 2009. Phylogeography of the brown hare (*Lepus europaeus*) in Europe: a legacy of south-eastern Mediterranean refugia? *Journal of Biogeography*, **36**: 515–528.
- Steffensen, J.P., Andersen, K.K., Bigler, M., Clausen, H.B., Dahl-Jensen, D., Fischer, H., Goto-Azuma, K., Hansson, M., Johnsen, S.J., Jouzel, J., Masson-Delmotte, V., Popp, T., Rasmussen, S.O., Röthlisberger, R., Ruth, U., Stauffer, B., Siggaard-Andersen, M.L., Sveinbjörnsdóttir, A.E., Svensson, A., White, J.W.C., 2008. High-resolution Greenland ice core data show abrupt climate change happens in few years. *Science*, **321**: 680–684.
- Stewart, J.R., 2007. Neanderthal extinction as part of the faunal change in Europe during Oxygen Isotope Stage 3. *Acta Zoologica Cracoviensia*, **50A**: 93–124.
- Stewart, J., Cooper, A., 2008. Ice Age refugia and Quaternary extinctions: an issue of Quaternary evolutionary palaeoecology. *Quaternary Science Reviews*, **27**: 2443–2448.
- Stewart, J.R., Lister, A.M., Barnes, I., Dalén, L., 2010. Refugia revisited: individualistic responses of species in space and time. *Proceedings of the Royal Society B*, **277**: 661–671.
- Stiller, M., Baryshnikov, G.F., Bocherens, H., Grandal d’Anglade, A., Hilpert, B., Münzel, S.C., Pinhasi, R., Rabeder, G., Rosendahl, W., Trinkaus, E., Hofreiter, M., Knapp, M., 2010. Withering away – 25,000 years of genetic decline preceded cave bear extinction. *Molecular Biology and Evolution*, **27**: 975–978.
- Stiller, M., Molak, M., Prost, S., Rabeder, G., Baryshnikov, G.F., Rosendahl, W., Münzel, S., Bocherens, H., Grandal-d’Anglade, A., Hilpert, B., Germonpré, M., Stasyk, O., Pinhasi, R., Tintori, A., Rohland, N., Mohandesan, E., Ho, S.Y.W., Hofreiter, M., Knapp, M., 2014. Mitochondrial DNA diversity and evolution of the Pleistocene cave bear complex. *Quaternary International*, **339–340**: 224–231.
- Stuart, A.J., 2005. The extinction of woolly mammoth (*Mammuthus primigenius*) and straight-tusked elephant (*Palaeoloxodon antiquus*) in Europe. *Quaternary International*, **126–128**: 171–177.
- Stuart, A.J., 2015. Late Quaternary megafaunal extinctions on the continents: a short review. *Geological Journal*, **50**: 338–363.
- Stuart, A.J., Lister, A.M., 2011. Extinction chronology of the cave lion *Panthera spelaea*. *Quaternary Science Reviews*, **30**: 2329–2340.
- Stuart, A.J., Lister, A.M., 2012. Extinction chronology of the woolly rhinoceros *Coelodonta antiquitatis* in the context of late Quaternary megafaunal extinctions in northern Eurasia. *Quaternary Science Reviews*, **51**: 1–17.
- Stuart, A.J., Lister, A.M., 2014. New radiocarbon evidence on the extirpation of the spotted hyaena (*Crocuta crocuta* (Erxl.) in northern Eurasia. *Quaternary Science Reviews*, **96**: 108–116.
- Stuart, A.J., Sulerzhitsky, L.D., Orlova, L.A., Kuzmin, Y.V., Lister, A.M., 2002. The latest woolly mammoths (*Mammuthus primigenius* Blumenbach) in Europe and Asia: a review of the current evidence. *Quaternary Science Reviews*, **21**: 1559–1569.
- Stuart, A.J., Kosintsev, P.A., Higham, T.F.G., Lister, A.M., 2004. Pleistocene to Holocene extinction dynamics in giant deer and woolly mammoth. *Nature*, **431**: 684–689.
- Svensson, A., Andersen, K.K., Bigler, M., Clausen, H.B., Dahl-Jensen, D., Davies, S.M., Johnsen, S.J., Muscheler, R., Parrenin, F., Röthlisberger, R., Seierstad, I., Steffensen, J.P., Vinther, B.M., 2008. A 60 000 year Greenland stratigraphic ice core chronology. *Climate of the Past*, **4**: 47–57.
- Tarasov, P.E., Volkova, V.S.V., Webb, T., Guiot, J., Andreev, A.A., Bezusko, L.G., Bezusko, T.V., Bykova, G.V., Dorofeyuk, N.I., Kvavadze, E.V., Osipova, I.M., Panova, N.K., Sevastyanov, D.V., 2000. Last glacial maximum biomes reconstructed from pollen and plant macrofossil data from northern Eurasia. *Journal of Biogeography*, **27**: 609–620.
- Ukkonen, P., Aaris-Sørensen, K., Arppe, L., Clark, P.U., Daugnor, L., Lister, A.M., Löugas, L., Seppä, H., Sommer, R.S., Stuart, A.J., Wojtal, P., Zupinš, I., 2011. Woolly mammoth (*Mammuthus primigenius* Blum.) and its environment in



- northern Europe during the last glaciation. *Quaternary Science Reviews*, **30**: 693–712.
- Van Heteren, A.H., MacLarnon, A., Soligo, C., Rae, T.C., 2014.** Functional morphology of the cave bear (*Ursus spelaeus*) cranium: a three-dimensional geometric morphometric analysis. *Quaternary International*, **339–340**: 209–216.
- Taboada, V.M., Mosquera, F.D., Gonzalez, L.F., d'Anglade, G.A., Romani, V.J.R., 1999.** Paleoecological implications inferred from stable isotopic signatures ( $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$ ) in bone collagen of "*Ursus spelaeus*" ROS.-HEIN. *Cadernos do Laboratorio Xeolóxico de Laxe: Revista de Xeoloxía Galega e do Hercínico Peninsular*, **24**: 73–87.
- Wilczyński, J., Wojtal, P., Sobieraj, D., Sobczyk, K., 2015.** Kraków Spadzista trench C2: new research and interpretations of Gravettian settlement. *Quaternary International*, **359–360**: 96–113.
- Wiszniewska, T., Mackiewicz, P., Stefaniak, K., Socha, P., Nowakowski, D., Nadachowski, A., 2010.** Dental enamel structure in fossil bears *Ursus spelaeus* and *U. wenzensis* (= *minimum*) in comparison to selected representatives of other Carnivora. In: *Morphology and Systematics of Fossil Vertebrates* (ed. D. Nowakowski): 125–142: DN Publisher, Wrocław.
- Wojtal, P., Wilczyński, J., 2015.** Zooarchaeological studies of large mammal remains from Kraków Spadzista site – trench C2 and trench E1 (2011–2012 excavations). In: *A Gravettian site in Southern Poland: Kraków Spadzista* (eds. P. Wojtal, J. Wilczyński and G. Haynes): 93–111. ISEA PAS.
- Wojtal, P., Wilczyński, J., Nadachowski, A., Münzel, S.C., 2015.** Gravettian hunting and exploitation of bears in Central Europe. *Quaternary International*, **359–360**: 58–71.
- Zazhigin, V.S., 1980.** Late Pliocene and Anthropogene Rodents of the South of Western Siberia (in Russian). Nauka, Moscow.

