

Reflections on subglacial megafloods: their possible cause, occurrence, and consequence for the global climate

A.J. (TOM) VAN LOON

Geological Institute, Adam Mickiewicz University, Maków Polnych 16, 61-606 Poznań, Poland;
e-mail: tvanloon@amu.du.pl

Abstract

Huge water reservoirs exist subglacially, for instance in the form of lakes. Failure of a wall around such a lake underneath the central part of a large ice mass may result in huge water outbursts. The resulting megafloods will rarely be traceable along the ice front, because much power is lost by friction and because the flood spreads subglacially over a large area. Truly giant meltwater outbursts might, however, still have an enormous power when reaching the marginal area of an ice cap. Such a megaflood may both affect the integrity of the ice mass, and help create a sliding layer over which huge ice masses can be easily transported towards the ocean, thus triggering a Heinrich-like event. This would have great impact on the global climate. Some of the Heinrich events that occurred during the past hundreds of thousands of years may well have been due to giant subglacial water outbursts, and such a situation may equally well occur in the time to come.

Keywords: jökulhlaups, megafloods, Heinrich events, sliding layer, subglacial lakes, climate change

Introduction

Some of the potentially more catastrophic natural phenomena are the sudden outbursts of glacial meltwater (Fig. 1), commonly called 'jökulhlaups' (from the Icelandic 'jökul' = glacier; and 'hlaup' = run-off). Most of the 'classical' Icelandic jökulhlaups seem directly related to geothermal heating or subglacial volcanism (see, among others, Kjartansson, 1951; Tryggvason, 1960; Björnsson and Einarsson, 1991; Larsen et al., 1998; Björnsson et al., 2000; Larsen, 2000; Maria et al., 2000). Volcanism-induced subglacial melting outside Iceland has been described from the Azas Plateau in the Tuva Republic of the Russian Federation (Komatsu

et al., 2007). The term 'jökulhlaup' is also used for sudden meltwater floods (with a discharge many times larger than normally) that have a different origin, for instance the failure of an ice or rock mass damming off a supraglacial or proglacial lake (Kochel et al., 2009).

Although the term 'jökulhlaup' is commonly used for the floods that are released in front of the ice (and thus can be observed directly), the term has also been used for supraglacial examples (Russell et al., 2001b). Although referring to a principally identical feature, the term will – in order to avoid confusion – not be used in the present contribution for the uncommonly large floods inside or underneath an ice cap that might originate due to the sud-



Fig. 1. The Skeiðarárjökull jökulhlaup following the Gyggja river, which is here ~100 m wide. The picture was taken on 5 November 1996 at 12.00 h, about one hour after water started to emerge from this part of the Vatnajökull glacier.

den release of huge volumes of meltwater from an englacial or subglacial water mass. Instead, the terms 'subglacial outburst' and 'subglacial megaflood' will be used.

Most Icelandic jökulhlaups cause little damage or casualties, because they involve a moderate mass of water, and occur in areas with little population. Their economic consequences can nevertheless be large (Rist, 1983; Jónsson et al., 1998), as the floods may destroy roads and bridges (Fig. 2). The damaging nature is partly due to the torrential character of the floods, which commonly carry a huge amount of cobbles and boulders. These may be transported through relatively narrow, but deep channels



Fig. 2. The destructive power of the 1996 Vatnajökull jökulhlaup. **A:** On 5 November 1996 at 14.40 h. The 900 m long Skeiðará bridge and protective dike system over the river are still intact, in spite of the high discharge of $15,000 \text{ m}^3 \text{ s}^{-1}$. **B:** The discharge (about $5,000 \text{ m}^3 \text{ s}^{-1}$ at this time, a few hours after the situation shown in Figure 2-A) has diminished, but the eastern end of the bridge crossing the streamway over the sandur plain has been washed away in the meantime.




Fig. 3. Deep channels within the snout of the Vatnajökull glacier during the 1996 jökulhlaup. **A:** During the main phase (photo Magnus Haldorsson). **B:** During the waning stage.

(Marren et al., 2009), which can be considered as the proglacial succession of the subglacial channels that become visible in the distal parts of the ice mass as a result of hydraulic fracturing of the snout (Fig. 3). Jökulhlaups can change both proglacial landforms (Smith et al., 2000; Russell et al., 2001a; Fay, 2002b) and proglacial sedimentation patterns (Maizels, 1991; Russell, 1993; Russell et al., 1999a; Fay, 2002a; Fleisher et al., 2003).

It is commonly accepted that jökulhlaups greatly contribute to the accumulation of sandurs (Tómasson, 1974; Maizels, 1983, 1989, 1993, 1997; Russell & Knudsen, 1999b, 2002b; Cassidy et al., 2003) because huge volumes of sediment are transported (Tómasson et al., 1980) – and the large sediment load influences, in turn, the jökulhlaup mechanism (Fowler & Ng, 1996). Yet, the deposits that are formed by a meltwater outburst inside an ice mass, i.e. by the failure of an ice dam blocking an englacial lake (Roberts et al., 2000a), as well as those formed underneath the ice by the outburst from a subglacial lake or by the sudden melting of large ice masses due to volcanic activity underneath the ice cap (Russell et al., 2003; Stevenson et al., 2009) are rarely preserved, because they are often destroyed by later meltwater outbursts. Even from outwash areas in front of the ice, descriptions of jökulhlaup deposits are commonly brief (Calkin, 2002). Only few descriptions of catastrophic floods of this specific type regard the Pleistocene (among others by O'Connor & Baker, 1992; Smith & Fisher, 1993; Rudoy & Baker, 1993; Geirsdóttir

et al., 2000; Mokhtari Fard & Ringberg, 2001; Rudoy, 2002). Large floods of meltwater from other sources (e.g. those from Lake Agassiz, released by the retreat of the Laurentide Ice Sheet that had blocked the outlet) have been reconstructed in fairly much detail (see, for instance, Teller et al., 2002; Clarke et al., 2004; Clayton & Knox, 2008). Such supraglacial and proglacial meltwater floods are out of scope in the present contribution.

Large meltwater outburst derived from the subglacial release of huge water masses are known in fairly much detail only from the 20th century (Guðmundsson et al., 1995), but reconstructions have also been made for Pleistocene occurrences. It was found, for instance, that broad, deep tunnel valleys along the margins of some lobes of the Laurentide Ice Sheet were formed by headward erosion of conduits through which catastrophic releases of water from subglacial reservoirs could take place, and that these reservoirs needed only a few decades to become refilled (Hooke & Jennings, 2006).

Well known jökulhlaups

The best known proglacial (subaerial) jökulhlaups are the small- and medium-scale events that occur on average every 1–2 years on Iceland. Much information is available with respect to their discharge and sediment load (Tómasson, 1974; Tómasson et al., 1980). These small-scale events commonly take place more




Fig. 4. Water emerging from underneath the front of the glacier on November 5, 1996, after the eruption of the Gjalp. Main portal, formed by fracturing of the snout of the Skeiðarárjökull, during the waning stage, one day after the jökulhlaup started. The ice wall is almost a hundred metres high.




Fig. 5. Large ice blocks (some tens of metres in diameter) were carried along with the water loaded also with smaller debris, during the first day of the Skeiðarárjökull outburst.

or less periodically (Mathews & Clague, 1993). The periodical occurrence can be due to the seasonal shift of glacier fronts, damming off glacial lakes during advance, and providing an outlet again during retreat (Tweed & Russell, 1999).

In contrast, large-scale jökulhlaups tend to occur with irregular intervals or as isolated phenomena (Brodzikowski & Van Loon, 1987, 1991), due to the destruction of lake-ice masses damming off a meltwater lake; the destruction of such ice dams is commonly ascribed to melting due to a sudden increase of geothermal heat (cf. Björnsson & Guðmundsson, 1993), related to volcanic eruptions. In other cases, it turns out that subglacial tunnels are formed that house water flows which undermine – by physical or thermal erosion – an ice mass that dams off a lake (Thórarinnsson, 1953; Liestöl, 1956; Björnsson, 1974; Nye, 1976), thus initiating a sudden outburst (Liu-Jinshi, 1992; Zhang-Xiangsong, 1992).

The characteristics of these large-scale jökulhlaups are much less well known than those of their smaller counterparts because the frequency of such events diminishes roughly exponentially with increasing size, as do most natural events. The best known examples of large-scale jökulhlaups (floods with discharges of at least tens of thousands of cubic meters per second) are those of 1918 (Katla) (Thórarinnsson, 1957; Björnsson, 1974; Björnsson et al., 2001) and 1996 (Vatnajökull) (Jónsson et al., 1998; Rus-

sell and Knudsen, 1999a, 2002a; Gomez et al., 2000, 2002). The latter, which lasted 33 hours, was caused by the rapid melting of some 3.8 km³ of ice (Guðmundsson et al., 1997; Björnsson, 2002) due to a volcanic eruption on 30 September 1996 underneath the Vatnajökull ice cap (Guðmundsson et al., 1997). It resulted in a sudden flood of some 45,000–53,000 m³ of water per second that was issued from underneath the glacier (Fig. 4) at velocities up to 6 m s⁻¹, and transported ice blocks (Fig. 5) of at least 25 m long over the Skeiðararsandur. This flood was nevertheless small compared to the 1918 Katla jökulhlaup, a torrential flood of some 300,000 m³ s⁻¹ of water, transporting 25,000 tons of ice per second and an equal amount of sediment, which resulted in a density of 1.170 g cm⁻³ (Tómasson, 1996). Krüger (1994) mentions the transport of ice blocks of 40–50 m (most of which were left behind on the sandur area between the ice front and the sea) by this flood. The largest Holocene jökulhlaup known from Iceland had a peak discharge of 900,000 m³ s⁻¹ (Alho et al., 2005). This catastrophic event was still of restricted size, in its turn, in comparison with a jökulhlaup that took place in front of Lake Missoula during the late Pleistocene, estimated to have reached a peak discharge of at least 17 (±3) million m³ s⁻¹ (O'Connor & Baker, 1992).

Objectives

The above can be considered as a simplified overview of some aspects of jökulhlaups that are relevant in the framework of questions such as: (1) is it possible that only part of the subglacial outbursts can be observed directly, because most of the meltwater remains hidden due to an entirely subglacial development (water running from one subglacial reservoir to another); (2) could such subglacial meltwater outbursts have extreme characteristics; and (3) if so, might the effects spread to beyond the ice front, and what could be the effects for the non-glaciated world (for instance, for the global climate)?

It seems worth while to consider these questions, as a good understanding of large meltwater floods may be beneficial for economic, safety and climate-policy reasons. Modelling on the basis of a better insight may help to predict such floods (Clarke, 1982; Björnsson, 1992; Maizels and Russell, 1992) and thus to reduce the risk of damage and loss of lives.

It seems also worth while to mention here explicitly that the main objective of the present contribution is to discuss the possibility of the occurrence of huge subglacial water outbursts, the possible sequence of events, and their possible consequences. It is not intended to analyse the possible pertinent processes themselves in any detail; nor is it intended to analyse the hydrological conditions or the physics that play a role just before, during and immediately after such an event, as no measurements of the physical conditions during such an event have ever been made under the central part of a huge ice cap; such an analysis could therefore now not be based on measurement data. Rather it is the intention to stimulate other researchers to think about methods that might help to gather such data, and to discuss the possible causes and consequences of huge subglacial water outbursts.

Considerations

If extremely large subglacial outbursts might occur, huge bodies of subglacial water

are a prerequisite. Large ice bodies tend to have – at least locally – a temperate character, which means that the combination of the geothermal heat flux and the insulation capacity of the ice results in the presence of water at the base of the ice mass. The volume of water can, indeed, be extremely large: Lake Vostok, under 3750–4150 m of ice of the Antarctic ice sheet (Souchez et al., 2003), has a surface area of some 14,000 km² (roughly 200×70 km). With a maximum depth of 670 m (Priscu et al., 1999), the water content of the lake is some 5,000 km³. This does not necessarily imply, however, that all this water – or a significant part of it – may be released in the form of a giant subglacial flood, but even a minor percentage of such a water volume could easily result in a sudden flood that may be comparable with those described from glacial times. Such an event requires, obviously, specific conditions, among which changes in the configuration of the ice masses that surround the lake, or susceptibility of the surrounding hard-rock walls to erosion.

Lake Vostok (Fig. 6) was, for several reasons, the object of much research during the past decade. It was found, for instance, that melting and freezing of the roof above the lake involve considerable quantities of ice (Jouzel et al., 1999; Siegert et al., 2000), which may be related to the heat flux from the lake into the ice (Thoma et al., 2008). Keeping in mind that there are more than 150 subglacial lakes in the Antarctic region (Kohler, 2007), although not of the same size, one must conclude that the total mass of material that undergoes phase




Fig. 6. Subglacial Lake Vostok (~200 km×70 km) on Antarctica, made visible through RADARSAT. Photo NASA/Goddard Space Flight Center Scientific Visualization Studio.

changes between water and ice must be much larger than the figures for Lake Vostok alone indicate. This is important because it implies that there are giant amounts of water available under the Antarctic ice sheet, which might be the source of sudden subglacial outbursts, for instance when the ice mass damming off such a lake melts away or is otherwise affected, for instance being moved, as may be the case when an ice-flow changes its direction.

The interest in Lake Vostok and other subglacial lakes has not yet answered all questions, partly because a drilling project was stopped in order to avoid contamination of a possible 'fossil' fauna or flora with present-day organisms (Nadis, 1999). Little information is therefore available about such aspects as (1) the amount of ice melting annually; (2) the mass balance of melting ice and freezing water; (3) the conditions limiting the size of the lake (ice dams? hard-rock topography?); (4) the characteristics (direction, transfer velocity, etc.) of the geothermal heat flux; (5) the movement of the ice immediately surrounding the lake, possibly temporarily blocking the outflow and eventually losing its function as a barrier again. It has been predicted already in 2000 that, in due time, Lake Vostok (or any other huge lake under the ice mass of Antarctica) might become the source of a giant subglacial meltwater outburst (Van Loon, 2000), and this prediction has since been proven to be correct, because a water mass of some 2 km³ travelled between 1996 and 1998 through the Adventure Subglacial Trench on Antarctica from one subglacial lake to another (Wingham et al., 2006; Carter et al., 2009).

The finding that huge water masses flow from one subglacial lake to another proves once more that the hydrology under large ice caps is still poorly understood (Wingham et al., 2006). It also proves that some ideas that are commonly taken for granted, commonly on the basis of physics that apparently do not prevail, are incorrect. It is often assumed, for instance, that – in the central parts of a large ice mass – no significant air-filled space can be present between the substratum and the ice, because of the downward pressure exerted by the weight of the ice. It is also commonly assumed that no

such space can be present between the water surface of a subglacial lake and the ice mass, because the "potentiometric surface is high up in the ice, just as in any confined groundwater aquifer". If no such air-filled spaces between the water surface and the ice would exist, it can, however, not be explained why huge water masses can embouch in a subglacial lake. Moreover, one might question how 'lake ice' can be formed if the water is under high pressure, and if there is no space for a volume increase. It is therefore, in contrast to common statements, not realistic to exclude by definition that large subglacial spaces occur through which megafloods can find their way.

Sequence of events characterising a giant subglacial outburst

Excellent progress has been made toward a refined theory of floodwater flow through glaciers (see, among others, Clarke, 2003), but the insight into the occurrence of outbursts and related processes under subglacial conditions is still small in comparison to the insight into proglacial flows: only two cases of floods from one subglacial site to another are well-known. The first is the 1996 Gjálp eruption in Iceland, where the meltwater from the eruption site accumulated in Grímsvötn Lake for several weeks before being released as a jökulhlaup that reached the area in front of the ice (Guðmundsson et al., 1997); it is known that the water volume of this lake can fluctuate considerably: in the days between 11 and 16 August 2004, it increased by 18 million cubic metres (Berthier et al., 2006). The second example regards the above-mentioned example in Antarctica, where a large subglacial water flow made the water level in one subglacial lake drop, and in another rise (Wingham et al., 2006). Differences in topographic height of the places between which the water flows can explain such transport, but differences in hydrostatic pressure may also play an important part.

There is clear sedimentary and geomorphological evidence from retreating ice margins

and formerly glaciated areas for subglacial floods (see, among others, Russell, 1994): the glaciological effect of floodwater movement through glaciers is reflected in ice fracturing and ice-surface sagging in the inlet region for the subglacial flood tract (Sturm and Benson, 1985). Hydrodynamical theories on the propagation and growth of subglacial floods have been obtained from observations in the inlet and outlet zones of flood-affected glaciers (Clarke, 2003). It seems therefore likely that huge lake outbursts, resulting in much larger floods than ever witnessed for subaerial jökulhlaups in historical times or reconstructed for prehistorical times, can occur underneath large ice masses. Only the following sequence of events need take place for the purpose:

1. Due to volcanic activity (Fig. 7), a temporarily increasing geothermal heat flux, or the supply through crevasses and tunnels of initially supraglacial meltwater, large water bodies are formed underneath a thick ice mass. Such water reservoirs may also develop in zones of equipotential convergence, at the point where hydraulic gradients are minimal. Ever growing lakes may thus originate, the final extent of which is determined by either a 'high' in the hard-rock substratum or an ice dam.
2. The water level rises as long as sufficient heat – from a volcanic eruption or from the geothermal heat flux – is supplied for additional melting of ice, or as long as additional water is supplied through upstream channels or through crevasses. This process goes on as long as the bordering ice mass or hard-rock 'high' is higher than the water level (this may last thousands of years).
3. The water level reaches the lowest upper limit of a hardrock barrier and starts overflowing, or the water pressure in the lake becomes high enough to force a water flow out of the lake by forming a tunnel underneath an ice dam or by flotation of such a dam (Tweed, 2000).
4. The result is, particularly in the case of an ice barrier, a rapid downwards erosion of the dam, resulting in an equally rapidly increasing release of a water mass with similarly increasing erosional power, so that a



Fig. 7. Eruption of the Grímsvötn volcano under the Vatnajökull glacier. Photo Oddur Sigurdsson.

- megaflood follows, which has all characteristics of a jökulhlaup (cf. Walder & Costa, 1996).
5. The process stops after a situation has been reached that either the lake has been emptied, or no further downwards erosion of the barrier takes place.

After this sequence of events has occurred, a new ice dam may eventually form and a new, similar sequence of events can start.

If sufficient air-filled space is available underneath the ice mass (and spaces must be available underneath temperate ice masses due to ice melting as a result of the geothermal heat flux in combination with the insulation capacity of ice), and if the distance to the ice front is sufficiently large, the torrential flood will gradually lose its power (by friction and by spreading over a large area) so that it need not necessarily preserve its torrential character when the water reaches the area in front of the ice. In such a case, the outburst will remain unnoticed: only an insignificant temporary increase of the sediment-loaded meltwater flow will possibly be recorded, contributing to the buildup of 'normal' glaciofluvial deposits and glaciolacustrine sediments (Gruszka, 2005).

Possible consequences

It is known from the Antarctic and Greenland ice sheets and many small glaciers that subglacial water can act as a lubricant, forming a 'film' over which ice masses can slide easily

(Björnsson, 1998). It should be emphasised in this context that such sliding is a process that differs fundamentally from the transport of, for instance, icebergs by ocean currents. In the latter case, buoyancy allows the water to carry the ice masses, whereas it is the almost negligible friction that allows an ice mass to slide downwards over the lubricant (sliding) layer. As a consequence, ice masses with considerable height can slide down over the lubricant water layer, also when the thickness of this layer is by far insufficient to allow such large ice masses to float within the water layer.

In addition, it is assumed on the basis of 'fossil' deposits that fairly strong subglacial meltwater streams can transport a large amount of clastic particles, thus forming low-viscosity slurries (Knight, 2003); such slurries may increase the effectiveness of the lubricant (sliding) layer underneath an ice mass. In the vicinity of the ice front, a water layer may therefore facilitate the start of surges, resulting in ice masses that move relatively fast over the sliding layer in a downslope direction (a slope of only a fraction of 1° is sufficient). Large icebergs can thus be transported to the ocean. The ice masses that slide down over the water layer can originate because more or less vertical fractures (crevasses) can be formed in the ice (Malthe-Sørensen et al., 1998; Waller et al., 2001), due to tensile stresses induced by the ice flow. Particularly lifting of ice by flowing water masses creates large tensile stresses and rupturing of the ice, which explains why large ice blocks can be transported by high-magnitude jökulhlaups. Shock waves may also play a (probably minor) role in the formation of ice blocks that can be transported over a lubricant layer. Such shock waves can arise from the sudden pressure exerted on ice walls if torrential floods find these walls on their ways.

It is obvious that the thickness (which influences the effectiveness) of a lubricant sheet of water depends on the amount of water supplied. Considering that increasingly larger floods can break up ever thicker ice masses, there must be a positive relationship between the force of a subglacial megaflood (and thus its mass) and the distance from the ice front where cracks may be formed that isolate ice

masses that may start sliding down over the lubricant water sheet. Large floods themselves may, moreover, favour the development of fractures in the ice (Russell et al., 1999b; Roberts et al., 2000b).

The above reasoning (Van Loon, 2000) comes – though following an entirely different approach – to a similar conclusion as reached by Johnson & Lauritzen (1995). They explain the global cooling of 116 ka ago as the result of a Heinrich event. The Heinrich event of 116 ka ago (reflected by a layer of dropstones in the Atlantic; see Scourse et al., 2000) occurred according to Johnson & Lauritzen (1995) as a consequence of a giant jökulhlaup. This flood must have originated after an ice dam that was formed due to surging and that was responsible for the presence of Lake Zissaga had failed; Lake Zissaga, with its 2,900,000 km² surface area and its 600,000 km³ water content, was 150 times larger than Lake Agassiz, which can be considered as its successor lake, covering only part of the Lake Zissaga area. The fundamental difference between the idea of Johnson & Lauritzen (1995) and the hypothesis presented here is that they reconstruct, like Teller et al. (2002) and Clarke et al. (2004), a jökulhlaup derived from a subaerial lake to explain climatic cooling (by triggering a Heinrich event); in contrast, the present contribution deals with the possibility of large subglacial lakes as the source of megafloods, which may also trigger such events. Another difference is that huge subaerial proglacial and supraglacial lakes require uncommon situations – such as the blocking of a huge depression by the Laurentide Ice Sheet – and are consequently relatively rare phenomena. In contrast, an irregular bedrock topography – as present under the Antarctic ice sheet – favours the continuous presence of huge water reservoirs. These form potential sources of sudden giant meltwater outbursts that may result in megafloods that might carry such large ice masses to the ocean that Heinrich-like events (with the potential consequence of global cooling) could take place.

Discussion

The larger a subglacial sheet-like flood is, the larger the ice mass that can be transported 'on its back' to the ocean. No reliable and statistically significant data about the relationship between jökulhlaup frequency and jökulhlaup size are available. It is therefore impossible to estimate with any degree of certainty the frequency of megafloods that originate and develop underneath huge icecaps and that might transport such huge icebergs to the sea that a Heinrich-like event would result. Johnson & Lauritzen (1995) argue that many Heinrich events may be due to processes identical - or at least comparable - to those that induced the 116 ka cooling).

Considering the frequency of (mostly small to moderate) jökulhlaups in Iceland during the 19th and 20th century, and considering the fact that in nature the frequency of specific phenomena decreases with increasing size/power, it must be considered very well possible that megafloods can occur underneath the land-ice masses of Antarctica every 10,000–100,000 years. Because the frequency of Heinrich events falls within this range, it would be scientifically challenging to evaluate the possibility that some phases of global cooling were - directly or indirectly - a result of surging that was triggered by extremely large subglacial floods. It may be true that 'real' Heinrich events are known only from the Atlantic (due to giant icebergs broken off the Laurentide ice sheet), but Heinrich-like events must also have occurred as a result of huge ice masses broken off the Antarctica ice sheet. This is indicated by significant layers of ice-rafted debris in the surrounding oceans (Hou et al., 1998) that coincide with low-temperature intervals on Earth, for instance 14.3–14.6 and 1.7–10.3 ka ago (Kanfoush et al., 2000). It should be kept in mind in this context that the existence of huge lakes on Antarctica, such as Lake Vostok, provides an ideal source for the water masses needed for such floods. Considering the occurrence of Heinrich events in the not so remote past, one could well imagine that the present-day conditions are still suitable for the occurrence of a new event: the ice or hardrock barrier dam-

ming off now the outlet of a sufficiently large subglacial lake may become overflow (or in the case of ice: undermined), triggering a giant subglacial outburst that could initiate a Heinrich event. The formation of a sufficiently thick lubricant layer with enough bearing capacity might have the same effect.

One might question why Heinrich events seem to have been restricted - or at least largely restricted - to the northern hemisphere (Scourse et al., 2000). This can be explained - if Heinrich events were triggered by catastrophic subglacial floods indeed - only by the fact that the conditions underneath the Laurentide Ice Sheet differed from those under the Antarctic ice cover. Analysis of such conditions (and their differences) is of great importance if the risk of megafloods under the Antarctic and the Greenland ice sheets is to be assessed. Relatively little is known about the Greenland conditions, as direct observations are impossible (Carter et al., 2009, call the transport of subglacial water "... an enigmatic and difficult to observe process"), and remote techniques (including geophysics) yield results that can, as a rule, be interpreted in various ways. One of the differences between Antarctica and Greenland is that the former is many times larger than the latter, whereas the ice-surface is more inclined in Greenland. In addition, it seems that the bedrock in the interior of Greenland does not have much relief. These two factors make the existence of numerous large subglacial lakes in Greenland unlikely; in fact no such lakes are known as yet from Greenland (M.J. Siegert, pers. comm., 2005). This is in strong contrast to the situation in Antarctica (Siegert et al., 2005) where subglacial lakes abound: apart from the 14,000 km² large Lake Vostok (Fig. 6), other subglacial lakes under Dome C amount to at least 15,000 km², and some additional 15,000 km² of subglacial lakes are known from under the remainder of the ice sheet. The total water volume underneath the ice is estimated to be 4,000–12,000 km³, possibly some 7140 km³ (Siegert, 2000), thus being much more than on Greenland. Moreover, the bedrock of Greenland has the shape of a bowl, which makes it much more likely that any hypothetical megaflood would be directed towards the

centre than towards the margin of the icecap. It is interesting in this context that few lakes have thus far been interpreted underneath ancient icecaps; one example is subglacial Lake Mc-Gregor, supposedly present under the Laurentide ice sheet in south-central Alberta (Munro-Stasiuk, 2003).

Processes like the interplay between ice movement, water/ice interaction, and geo-thermal-heat transfer might play a part in the occurrence of giant water outbursts under the Antarctic ice sheet. Such outbursts may, however, be so far away from the ice front that – however large the outburst may originally be – the effects in the more marginal parts become small. The diminishing effect may be due to the distribution of the water over an ever increasing surface area, to re-freezing of the water, and/or to loss of current velocity due to friction. The result of the decrease of the flood's power with increasing distance from the place of origin may be that no huge ice masses may be triggered to move in the more marginal parts of the ice. In that case, even a giant water outburst in the central part of the ice sheet may remain unnoticed. Obviously, the fact that no such outbursts have ever been noticed along the margin of the Antarctic ice sheet might also be ascribed to their non-existence. It is, for instance, principally possible that, apart from the 68 lakes known to be present under the East Antarctic ice sheet (Vincent, 1999), there is insufficient air-filled space in the direct vicinity of lakes under the Antarctic ice sheet to allow these lakes to overflow into such spaces. Another aspect to be considered is the topography of the hard-rock substratum. One could imagine – but there are insufficient detailed data available as yet – that the topography under the Antarctic ice sheet is so irregular, for instance due to glacial erosion during periods with no or much smaller ice caps (cf. Engström et al., 2000), that lake levels cannot rise sufficiently to start overflow. It is also possible that the physical characteristics of the hard-rock barriers in Antarctica do not allow them to be eroded strongly enough to start catastrophic floods. More detailed seismic mapping might provide the necessary data, changing the above idea into a testable hypothesis (cf. Van Loon, 2004).

Final remarks

Little is known about the occurrence of meltwater outbursts in the more central parts of huge icecaps. It is not known, for instance, what precise processes might be involved. In addition, it is not understood why Heinrich events are known well only from the northern hemisphere, and only from the Laurentide ice sheet. It is not well understood either what is the reason behind the rough cyclicity of these events.

No attention has been paid thus far to a possible triggering of Heinrich events by the sudden, giant outbursts of subglacial lakes. Several characteristics of both subglacial megafloods and Heinrich events could, however, be explained well if such a relationship exists. Admittedly, several processes (such as acceleration of ice sheets by basal decoupling or enhanced bed deformation) have been proposed already, and the proposed processes seem feasible. This does not imply, however, that alternative explanations should not be considered (the number of possible causes of the mass extinction at the P/T boundary is ever growing!). It is therefore suggested here that the possibility of a possible relationship between subglacial megafloods and Heinrich-like events be further investigated, if only because of the probable consequences of future Heinrich-like events for the global climate. It should be noticed in this context that a subglacial volcano that was active only some 2000 years ago and that should therefore be considered as dormant rather than as extinct, has recently been discovered beneath the West-Antarctic ice sheet (Corr & Vaughn, 2008) and that its activity has already been related to increased flowage of Antarctic ice masses.

As it seems that meltwater discharges during deglaciation show a complex interrelationship with climate, landscape evolution and the mechanisms determining glacier movement, not only subglacial volcanism but also present-day global warming might favour the conditions under which the sudden release of giant masses of subglacial meltwater (cf. Carter et al., 2009) might trigger a Heinrich-like event, with global consequences for society. It is note-

worthy to remark in this context that such an effect would induce a fall in temperature in large parts of the world, so that global warming might, in this respect, have a negative feedback.

Acknowledgements

I want to thank the colleagues who, after my presentation of these views at the 2000 GSA annual meeting in Reno, contacted me to discuss the various aspect of my hypothesis. Their remarks and suggestions were most helpful. I should mention in this context that several of the colleagues who initially criticized the hypothesis put forward here (mainly because of lack of data about subglacial floods) became less sceptical after, a few years later, the probably interrelated water-level fluctuations of two Antarctic lakes proved the subglacial flow, in whatever form, of huge water volumes.

I want to thank particularly my colleagues in Reykjavik, Iceland, who had some highly useful suggestions, and who provided me with some excellent literature about the topic. Magnús Guðmundsson (University of Iceland) kindly provided photographs taken by himself and Finnur Pálsson (Figs. 1 through 5, apart from Fig. 3-A)

Tomasz Zielinski and Beata Gruszka (both A. Mickiewicz University, Poznan), Andrew Russell (Newcastle University), Magnus Guðmundsson (University of Iceland) and Cathy Delaney (Manchester Metropolitan University) kindly commented upon earlier drafts of the present contribution, and suggested numerous improvements.

References

- Alho, P., Russell, A.J., Carrivick, J.L. & Käyhkö, J., 2005. Reconstruction of the largest Holocene jökulhaup within Jökulsá á Fjöllum, NE Iceland. *Quaternary Science Reviews* 24, 2319–2334.
- Berthier, E., Björnsson, H., Pálsson, F., Feigl, K.L., Llubes, M. & Rémy, F., 2006. The level of the Grímsvötn subglacial lake, Vatnajökull, Iceland, monitored with SPOT5 images. *Earth and Planetary Science Letters* 243, 293–302.
- Björnsson, H., 1974. Explanation of jökulhlaups from Grímsvötn, Vatnajökull, Iceland. *Jökull* 24, 1–26.
- Björnsson, H., 1992. Jökulhlaups in Iceland: prediction, characteristics and simulation. *Annals of Glaciology* 16, 95–106.
- Björnsson, H., 1998. Hydrological characteristics of the drainage system beneath a surging glacier. *Nature* 395, 771–774.
- Björnsson, H., 2002. Subglacial lakes and jökulhlaups in Iceland. *Global and Planetary Change* 35, 255–271.
- Björnsson, H. & Einarsson, P., 1991. Volcanoes beneath Vatnajökull, Iceland: evidence from radio-echo sounding, earthquakes and jökulhlaups. *Jökull* 40, 147–168.
- Björnsson, H. & Guðmundsson, M.T., 1993. Variations in the thermal output of the subglacial Grímsvötn caldera, Iceland. *Geophysical Research Letters* 20, 2127–2130.
- Björnsson, H., Pálsson, F. & Guðmundsson, M.T., 2000. Surface and bedrock topography of the Mýrdalsjökull ice cap, Iceland: the Katla caldera, eruption sites and routes of jökulhlaups. *Jökull* 49, 29–46.
- Björnsson, H., Pálsson, F., Flowers, G.E. & Guðmundsson, M.T., 2001. The extraordinary 1996 jökulhlaup from Grímsvötn, Vatnajökull, Iceland. *EOS* 82, 47.
- Brodzikowski, K. & Van Loon, A.J., 1987. A systematic classification of glacial and periglacial environments, facies and deposits. *Earth-Science Reviews* 24, 297–381.
- Brodzikowski, K. & Van Loon, A.J., 1991. *Glacigenic sediments*. Developments in Sedimentology 49. Elsevier, Amsterdam, 674 pp.
- Carter, S.P., Blankenship, D.D., Young, D.A., Peters, M.E., Holt, J.W. & Siegert M.J., 2009. Dynamic distributed drainage implied by the flow evolution of the 1996–1998 Adventure Trench subglacial lake discharge. *Earth and Planetary Science Letters* 283 24–37.
- Cassidy, N.J., Russell, A.J., Marren, P.M., Fay, H., Rushmer, E.L., Van Dijk, T.A.G.P. & Knudsen, Ó., 2003. GPR-derived architecture of November 1996 jökulhlaup deposits, Skeiðarársandur, Iceland. [In]: C.S. Bristow & H.M. Jol (eds): *Ground penetrating radar in seimentation*. Geological Society London Special Publication 211, 153–166.
- Calkin, P.E., 2002. Global glacial chronologies and causes of glaciation. [In]: J. Menzies (ed.): *Modern and past glacial environments*. Butterworth-Heinemann, Oxford, p. 15–52.
- Clarke, G.K.C., 1982. Glacier outburst floods from “Hazard Lake”, Yukon Territory, and the problem of flood magnitude prediction. *Journal of Glaciology* 28, 3–21.
- Clarke, G.K.C., 2003. Hydraulics of subglacial outburst floods: new insights from the Spring-Hutter formulation. *Journal of Glaciology* 49, 299–313.
- Clarke, G.K.C., Leverington, D.W., Teller, J.T. & Dyke, A.S., 2004. Paleohydraulics of the last outburst flood from glacial Lake Agassiz and the 8200 BP cold event. *Quaternary Science Reviews* 23, 389–407.
- Clayton, J.A. & Knox, J.C., 2008. Catastrophic flooding from Glacial Lake Wisconsin. *Geomorphology* 93, 384–397.

- Corr, H.F.J. & Vaughn, D.G., 2008. A recent volcanic eruption beneath the West Antarctic ice sheet. *Nature Geoscience* 1, 122–125.
- Engström, D.R., Fritz, S.C., Almendinger, J.E. & Juggins, S., 2000. Chemical and biological trends during lake evolution in recently deglaciated terrain. *Nature* 408, 161–166.
- Fay, H., 2002a. The formation of ice-block obstacle marks during the November 1996 glacier outburst flood (jökulhlaup), Skeiðarársandur, southern Iceland. [In:] I.P. Martini, V.R. Baker & G. Garzon (eds): *Flood and megaflood deposits: recent and ancient*. International Association of Sedimentologists Special Publication 32, 85–97.
- Fay, H., 2002b. Formation of kettle holes following a glacial outburst flood (jökulhlaup), Skeiðarársandur, southern Iceland. [In:] A. Snorasson, H.P. Finnsdóttir & M. Moss, M. (eds): *The extremes of the extremes: extraordinary floods – Proceedings of a symposium held at Reykjavik, Iceland, July 2000*. International Association of Hydrological Sciences Special Publication 271, 205–210.
- Fleisher, P.J., Bailey, P.K. & Cadwell, D.H., 2003. A decade of sedimentation in ice-contact, proglacial lakes, Bering Glacier, AK. [In:] J. Knight (ed.), *Subglacial lake sediments*. *Sedimentary Geology* 160, 309–324.
- Fowler, A.C. & Ng, F.S.L., 1996. The role of sediment transport in the mechanics of jökulhlaups – Proceedings of the International Symposium on Glacial Erosion and Sedimentation held at Reykjavik, Iceland, 20–25 August 1995. *Annals of Glaciology* 22, 255–259.
- Geirsdóttir, Á., Hardardóttir, J. & Sveinbjörndóttir, Á.E., 2000. Glacial extent and catastrophic meltwater events during the deglaciation of southern Iceland. *Quaternary Science Reviews* 19, 1749–1761.
- Gomez, B., Smith, L.C., Magilligan, F.J., Mertes, F.A.K. & Smith, N.D., 2000. Glacier outburst floods and outwash plain development: Skeiðarársandur, Iceland. *Terra Nova* 12, 126–131.
- Gomez, B., Russell, A.J., Smith, L.C. & Knudsen, Ó., 2002. Erosion and deposition in the proglacial zone: the 1996 jökulhlaup on Skeiðarársandur, southeast Iceland. [In:] Á. Snorasson, H.P. Finnsdóttir & M. Moss (eds): *The extremes of the extremes: extraordinary floods*. International Association of Hydrological Sciences Special Publication 271, 217–221.
- Gruszka, C., 2005. The Pleistocene lacustrine sediments in the Bełchatów mine (central Poland) – endogenic and exogenic controls. *Sedimentary Geology* 193, 149–166.
- Guðmundsson, M.T., Björnsson, H. & Pálsson, F., 1995. Changes in jökulhlaup sizes in Grímsvötn, Vatnajökull, Iceland, 1934–91, deduced from in-situ measurements of subglacial lake volume. *Journal of Geology* 41, 263–272.
- Guðmundsson, M.T., Sigmundsson, F. & Björnsson, H., 1997. Ice-volcano interaction of the 1996 Gjálp subglacial eruption, Vatnajökull, Iceland. *Nature* 389, 954–957.
- Hooke, R.L. & Jennings, C.E., 2006. On the formation of the tunnel valleys of the southern Laurentide ice sheet. *Quaternary Science Reviews* 25, 364–1372.
- Hou, H., Luo, Y., Zheng, H., Wang, B. & Tang, X., 1998. Heinrich layer in Antarctic marine sediments and its significance to global changes. *Chinese Science Bulletin* 43, 1830–1834.
- Johnson, R.G. & Lauritzen, S.-E., 1995. Hudson Bay-Hudson Strait jökulhlaups and Heinrich events: a hypothesis. *Palaeogeography, Palaeoclimatology, Palaeoecology* 117, 123–137.
- Jónsson, P., Sigurðsson, O., Snorasson, Á., Víkingsson, S., Kaldal, I. & Árnason, S., 1998. Course of events of the jökulhlaup on Skeiðarársandur, Iceland, in November 1996. [In:] J.C. Cañaveras, M., Ángeles García del Gura & J. Soria, J. (eds): *15th International Sedimentological Conference – Proceedings of the 15th International Sedimentological Conference, April 12–17, Alicante*. International Association of Sedimentologists, 456–457.
- Jouzel, J., Petit, J.R., Souchez, R., Barkov, N.I., Lipenkov, V.Ya., Raynaud, D., Stievenard, M., Vassiliev, N.I., Verbeke, V. & Vimeux, F., 1999. More than 200 meters of lake ice above subglacial Lake Vostok, Antarctica. *Science* 286, 2138–2141.
- Kanfoush, S.L., Hodell, D.A., Charles, C.D., Guilderson, T.P., Mortyn, P.G. & Ninnemann, U.S., 2000. Millennial-scale instability of the Antarctic ice sheet during the last glaciation. *Science* 288, 1815–1818.
- Kjartansson, G., 1951. *The eruption of Hekla, 1947–48, II. 4. Water flood and mud flows*. Societas Scientiarum Islandica, 60 pp.
- Knight, J., 2003. Temporal changes in subglacial meltwater activity: field evidence from the late Devensian in the north of Ireland. [In:] J. Knight (ed.): *Subglacial lake sediments*. *Sedimentary Geology* 160, 291–307.
- Kochel, R.C., Nickelsen, R.P. & Eaton, L.S., 2009. Catastrophic middle Pleistocene jökulhlaups in the upper Susquehanna River: Distinctive landforms from breakout floods in the central Appalachians. *Geomorphology* 110, 80–95.
- Kohler, J., 2007. Lubricating lakes. *Nature* 445, 830–831.
- Komatsu, G., Arzhannikov, S.G., Arzhannikova, A.V. & Ershov, K., 2007. Geomorphology of subglacial volcanoes in the Azas Plateau, the Tuva Republic, Russia. *Geomorphology* 88, 312–328.
- Krüger, J., 1994. Glacial processes, sediments, landforms, and stratigraphy in the terminus region of Myrdalsjökull, Iceland. *Folia Geographica Dania* 21, 1–233.
- Larsen, G., 2000. Holocene eruptions within the Katla volcanic system, south Iceland: characteristics and environmental impact. *Jökull* 49, 1–28.
- Larsen, G., Guðmundsson, M.T. & Björnsson, H., 1998. Eight centuries of periodic volcanism at the center of the Iceland hotspot revealed by glacier tephrostratigraphy. *Geology* 26, 943–946.
- Liestøl, O., 1956. Glacier-dammed lakes in Norway. *Norsk Geografisk Tidsskrift* 15, 122–149.
- Liu-Jinshi, 1992. Jökulhlaups in the Kunmalike River, southern Tien Shan Mountains, China. *Annals of Glaciology* 16, 85–88.
- Maizels, J.K., 1983. Channel changes, paleohydrology and deglaciation: evidence from some Lateglacial sandur

- deposits of NE Scotland. *Quaternary Studies in Poland* 4, 171–187.
- Maizels, J.K., 1989. Sedimentology, paleoflow dynamics and flood history of jökulhlaup deposits: paleohydrology of Holocene sediment sequences in S Iceland sandur deposits. *Journal of Sedimentary Petrology* 59, 204–223.
- Maizels, J.K., 1993. Lithofacies variations within sandur deposits: the role of runoff regime, flow dynamics and sediment supply characteristics. *Sedimentary Geology* 85, 299–325.
- Maizels, J.K., 1997. Jökulhlaup deposits in proglacial areas. *Quaternary Science Reviews* 16, 793–819.
- Maizels, J.K., 1991. Origin and evolution of Holocene sandurs in areas of jökulhlaup drainage, south Iceland. [In:] J.K. Maizels & C. Caseldine (eds): *Environmental change in Iceland: past and present*. Kluwer, Deventer, p. 267–300.
- Maizels, J.K. & Russell, A.J., 1992. Quaternary perspectives on jökulhlaup prediction. [In:] J.M. Gray (ed.): *Applications of Quaternary Research*. Quaternary Processes 2, 133–153.
- Malthe-Sørensen, A., Walmann, T., Jamtveit, B., Feder, J. & Jøssang, T., 1998. Modeling and characterization of fracture patterns in the Vatnajökull glacier. *Geology* 26, 931–934.
- Maria, A., Carey, S., Sigurdsson, H. & Kincaid, C., 2000. Source and dispersal of jökulhlaup sediments discharged to the sea following the 1996 Vatnajökull eruption. *Geological Society of America Bulletin* 112, 1507–1521.
- Marren, P.M., Russell, A.J. & Rushmer, E.L., 2009. Sedimentology of a sandur formed by multiple jökulhlaups, Kverkfjöll, Iceland. *Sedimentary Geology* 213, 77–88.
- Mathews, W.H. & Clague, J.J., 1993. The record of jökulhlaups from Summit Lake, northwestern British Columbia. *Canadian Journal of Earth Sciences* 30, 499–508.
- Mokhtari Fard, A., Ringberg, B., 2001. Sedimentological evidence of a meandering Younger Dryas subglacial conduit: Horn, SE central Sweden. [In:] A. Mokhtari Fard (ed.): *Recognition of abrupt climate change in clastic sedimentary environments*. Global and Planetary Change 28, 255–265.
- Munro-Stasiuk, M.J., 2003. Subglacial Lake McGregor, south-central Alberta, Canada. [In:] J. Knight (ed.): *Subglacial lake sediments*. *Sedimentary Geology* 160, 325–350.
- Nadis, S., 1999. Moves are afoot to probe the lake trapped beneath Antarctic ice. *Nature* 401, 203.
- Nye, J.F., 1976. Water flow in glaciers: jökulhlaups, tunnels and veins. *Journal of Glaciology* 17, 181–207.
- O'Connor, J.E., Baker, V.R., 1992. Magnitudes and implications of peak discharges from glacial Lake Missoula. *Geological Society of America Bulletin* 104, 267–279.
- Priscu, J.C., Adams, E.E., Lyons, W.B., Voytek, M.A., Mogk, D.W., Brown, R.L., McKay, Chr.P., Takacs, C.D., Welch, K.A., Wolf, C.F., Kirshtein, J.D. & Avci, R., 1999. Geomicrobiology of subglacial ice above Lake Vostok, Antarctica. *Science* 286, 2141–2144.
- Rist, S., 1983. Floods and flood danger in Iceland. *Jökull* 33, 119–132.
- Roberts, M.J., Russell, A.J., Tweed, F.S. & Knudsen, Ó., 2000a. Rapid sediment entrainment and englacial deposition during jökulhlaups. *Journal of Glaciology* 153, 349–351.
- Roberts, M.J., Russell, A.J., Tweed, F.S. & Knudsen, Ó., 2000b. Ice fracturing during jökulhlaups: implications for englacial floodwater routing and outlet development. *Earth Surface Processes and Landforms* 25, 1429–1446.
- Rudoy, A.N., 2002. Glacier-dammed lakes and geologic work of glacial superfloods in the Late Pleistocene, S Siberia, Altai Mts. *Quaternary International* 87, 119–140.
- Rudoy, A.N. & Baker, V.R., 1993. Sedimentary effects of cataclysmic late Pleistocene glacial outburst flooding, Altai Mts, Siberia. *Sedimentary Geology* 85, 53–62.
- Russell, A.J., 1993. Obstacle marks produced by flow around stranded ice blocks during a glacier outburst flood (jökulhlaup) in west Greenland. *Sedimentology* 40, 1091–1111.
- Russell, A.J., 1994. Subglacial jökulhlaup deposition, Jotunheimen, Norway. [In:] G.F. Dardis & A.M. McCabe (eds.): *Subglacial processes, sediments and landforms*. *Sedimentary Geology* 91, 131–144.
- Russell, A.J. & Knudsen, Ó., 1999a. Controls on the sedimentology of November 1996 jökulhlaup deposits, Skeiðarársandur, Iceland. [In:] N.D. Smith & J. Rogers (eds): *Fluvial Sedimentology VI*. International Association of Sedimentologists Special Publication 28, 315–329.
- Russell, A.J. & Knudsen, Ó., 1999b. An ice-contact rhythmite (turbidite) succession deposited during the November 1996 catastrophic outburst flood (jökulhlaup), Skeiðarárjökull, Iceland. *Sedimentary Geology* 127, 1–10.
- Russell, A.J. & Knudsen, Ó., 2002a. The effects of glacier-outburst flood flow dynamics in ice-contact deposits: November 1996 jökulhlaup, Skeiðarársandur, Iceland. [In:] I.P. Martini, V.R. Baker & G. Garzón (eds): *Flood and megaflood processes and deposits: recent and ancient examples*. International Association of Sedimentologists Special Publication 32, 67–83.
- Russell, A.J. & Knudsen, Ó., 2000b. The influence of channel flood history on the impact of the November 1996 jökulhlaup, Skeiðarársandur, Iceland. [In:] Á. Snorasson, H.P. Finnsdóttir & M. Moss (eds): *The extremes of the extremes: extraordinary floods*. International Association of Hydrological Sciences Publication 271, 243–247.
- Russell, A.J., Knudsen, Ó., Maizels, J.K. & Marren, Ph.M., 1999a. Channel cross-sectional area changes and peak discharge calculations in the Gígjukvísl river during the November 1996 jökulhlaup, Skeiðarársandur, Iceland. *Jökull* 47, 45–58.
- Russell, A.J., Tweed, F.S., Knudsen, Ó., Roberts, M.J. & Waller, R.I., 1999b. Ice fracturing and glacier sediment entrainment during two recent Icelandic jökulhlaups. *EOS* 80, 400.
- Russell, A.J., Knight, P.G. & Van Dijk, T.A.G.P., 2001a. Glacier surging as a control on the development

- of proglacial, fluvial landforms and deposits, Skeidarársandur, Iceland. [In:] A. Mokhtari Fard (ed.): Recognition of abrupt climate change in clastic sedimentary environments. *Global and Planetary Change* 28, 163–174.
- Russell, A.J., Knudsen, Ó., Fay, H., Marren, P.M., Heinz, J. & Tronicke, J., 2001b. Morphology and sedimentology of a giant supraglacial, ice-walled, jökulhlaup channel, Skeidarársandur, Iceland: implications for esker genesis. [In:] A. Mokhtari Fard (ed.): Recognition of abrupt climate change in clastic sedimentary environments. *Global and Planetary Change* 28, 193–216.
- Russell, A.J., Fay, H., Harris, T.D., Tweed, F. & Roberts, M.J., 2003. Preservation of jökulhlaups within subglacial sediments. *XVI INQUA Congress (Reno, July 2003) Programs with Abstracts* 42–1, 148.
- Scourse, D.E., Hall, I.R., McCave, I.N., Young, J.R. & Sugdon, C., 2000. The origin of Heinrich layers: evidence from H2 for European precursor events. *Earth and Planetary Science Letters* 182, 187–195.
- Siegert, M.J., 2000. Antarctic subglacial lakes. *Earth-Science Reviews* 50, 29–50.
- Siegert, M.J., Kwok, R., Mayer, Chr. & Hubbard, B., 2000. Water exchange between the subglacial Lake Vostok and the overlying ice sheet. *Nature* 403, 643–646.
- Siegert, M.J., Carter, S., Tabacco, I., Popov, S. & Blankenship, D., 2005. A revised inventory of Antarctic subglacial lakes. *Antarctic Science* 17, 453–460.
- Smith, D.G., Fisher, T.G., 1993. Glacial Lake Agassiz: The northwestern outlet and paleoflood. *Geology*, 21, 9–12.
- Smith, L.C., Alsdorf, D.E., Magilligan, F.J., Gomez, B., Mertes, L.A.K., Smith, N.D. & Garvin, J.B., 2000. Estimation of erosion, deposition, and net volumetric change caused by the 1996 Skeidarársandur jökulhlaup, Iceland, from synthetic aperture radar interferometry. *Water Resources Research* 36, 1583–1594.
- Souchez, R., Petit, J.-R., Jouzel, J., De Angelis, M. & Tison, J.-L., 2003. Reassessing Lake Vostok's behaviour from existing and new ice core data. *Earth and Planetary Science Letters* 217, 163–170.
- Stevenson, J.A., Smellie, J.L., McGarvie, D.W., Gilbert, J.S. & Cameron, B.I., 2009. Subglacial intermediate volcanism at Kerlingarfjöll, Iceland: Magma-water interactions beneath thick ice. *Journal of Volcanology and Geothermal Research* 185, 337–351.
- Sturm, M. & Benson, C.S., 1985. A history of jökulhlaups from Strandline Lake, Alaska, U.S.A. *Journal of Glaciology* 31, 272–280.
- Teller, J.T., Leverington, D.W. & Mann, J.D., 2002. Freshwater outbursts to the oceans from glacial Lake Agassiz and their role in climate change during the last deglaciation. *Quaternary Science Reviews* 21, 879–887.
- Thoma, M., Mayer, C. & Grosfeld, K., 2008. Sensitivity of subglacial Lake Vostok's flow regime on environmental parameters. *Earth and Planetary Science Letters* 269, 242–247.
- Thórarinnsson, S., 1953. Some new aspects of the Grímsvötn problem. *Journal of Glaciology* 2, 267–275.
- Thórarinnsson, S., 1957. The jökulhlaup from the Katla area in 1955 compared with other jökulhlaups in Iceland. *Jökull* 7, 21–25.
- Tómasson, H., 1974. Grímsvötnhlaup 1972, mechanism and sediment discharge. *Jökull* 24, 27–38.
- Tómasson, H., 1996. The jökulhlaup from Katla in 1918. *Annals of Glaciology* 22, 249–254.
- Tómasson, H., Ingólfsson, P., and Pálsson, S., 1980. Comparison of sediment load transport in the Skeidará jökulhlaups in 1972 and 1976. *Jökull* 30, 21–33.
- Tryggvason, E., 1960. Earthquakes, jökulhlaups and subglacial eruptions. *Jökull* 10, 18–22.
- Tweed, F.S., 2000. Jökulhlaup initiation by ice-dam flotation: the significance of glacier debris-content. *Earth Surface Processes and Landforms* 25, 105–108.
- Tweed, F.S., Russell, A.J., 1999. Controls on the formation and sudden drainage of glacier-impounded lakes: implications for jökulhlaup characteristics. *Progress in Physical Geography* 23, 79–110.
- Van Loon, A.J., 2000. What conditions are required for jökulhlaups? [In:] *Summit 2000 – Abstracts with programs Geological Society of America 112th Annual Meeting and Exposition (Reno, 2000): A-115–116*.
- Van Loon, A.J., 2004. The challenge of hypothesizing in the earth sciences. *Earth-Science Reviews* 65, 305–313.
- Vincent, W.F., 1999. Icy life on a hidden lake. *Science* 286, 2094–2095.
- Walder, J.S. & Costa, J.E., 1996. Outburst floods from glacier-dammed lakes: the effect of mode of drainage on flood magnitude. *Earth Surface Processes and Landforms* 21, 701–723.
- Waller, R.I., Russell, A.J., Van Dijk, T.A.G.P. & Knudsen, Ó., 2001. Jökulhlaup related ice fracture and the supraglacial routing of water and sediment, Skeidarársandur, Iceland. *Geografiska Annaler* 83A, 29–38.
- Wingham, D.J., Siegert, M.J., Shepherd, A. & Muir, A.S., 2006. Rapid discharge connects Antarctic subglacial lakes. *Nature* 440 1033–1036.
- Zhang-Xiangsong, 1992. Investigation of glacier bursts of the Yarkant River in Xinjiang, China. *Annals of Glaciology* 16, 135–139.