

Iridoids from *Cornus mas* L. and their potential as innovative ingredients in cosmetics

Zofia Nizioł-Łukaszewska^{1*}, Tomasz Wasilewski², Tomasz Bujak¹, Paweł Osika¹

¹The University of Information Technology and Management in Rzeszow, Department of Cosmetology, Kielnarowa 386a, 36-020 Tyczyn, Poland

²University of Technology and Humanities in Radom, Department of Chemistry, Chrobrego 27, Radom 26-600, Poland

*Corresponding author: e-mail: znizioł@wsiz.rzeszow.pl

Dogwood berries represent a valuable source of a variety of active ingredients. A group that deserves special attention comprises iridoids – compounds with potent antioxidant, antiinflammatory and antibacterial properties. The present study is an attempt to obtain an innovative plant material from dogwood berries. To this end, water and water/ethanol-based extracts (1:1) were prepared and, as the next step, an iridoids-rich fraction was isolated. The total content of iridoids was determined spectrophotometrically, and antioxidant properties of the isolates were concurrently assessed. Additionally, skin whitening activity of isolated fractions was assessed on the basis of tyrosinase inhibition measurement. The testing schedule also involved the formulation of model washing systems based on anionic surfactants. The effect of adding the fractions obtained by the above method on the irritant potential was assessed by determining the zein number

Keywords: dogwood, antioxidant activity, zein test, tyrosinase inhibition, iridoids.

INTRODUCTION

Cosmetic industry is one of the fastest growing industrial sectors both in Poland and across the world. As the market is very competitive, cosmetic manufacturers are required to offer innovative products. One method of obtaining products of this type is enhancing their formulas by adding a novel ingredient which is not commonly used in cosmetics. Recent years have seen a particular increase in consumer interest in cosmetics containing substances of plant origin. A number of scientific studies have shown that plant-based materials exhibit antibacterial, antiviral, antifungal and antiinflammatory properties which counteract adverse changes in the body, and have a positive effect on the condition and functioning of the skin^{1,2}. The group of beneficial plant-based materials also comprises dogwood berries. Biologically active substances contained in dogwood berries are known to have powerful antioxidant, antibacterial, toning and astringent effects. In addition, they effectively prevent inflammations in living organisms³⁻⁶. Dogwood has been found to contain numerous phenolic compounds, flavonoids, anthocyanins and vitamin C⁷⁻⁹.

Particular attention should be paid to iridoids, i.e. secondary metabolites in the form of cyclopentane monoterpenes. They may have a simple iridoid structure or occur as glycoside or ester compounds. Based on their chemical structure, iridoids can be divided into four main groups: iridoid glycosides, nonglycosidic (aglycone) iridoids, secoiridoids, and bisiridoids¹⁰. Dogwood berries have been shown to contain primarily oleanolic acid, ursolic acid, morroniside¹¹, loganin¹², sweroside, and cornuside. Iridoids have powerful antiinflammatory and antibacterial properties both in internal and external applications¹³. They are potent antioxidants, and may thus display anticancer activity in skin cancer^{14,15}, cancer of the colon, lungs, or nose and throat^{16,17}.

Since dogwood berries are associated with beneficial effects, extracts or substances isolated from them can be successfully used in cosmetic products. Active sub-

stances derived from dogwood and added to cosmetics successfully replace synthetic astringent agents which have a beneficial effect on human skin¹⁸. Iridoids reacting with amines have also been shown to yield a coloured product which is used as a colourant in the food and cosmetic industries. In addition, iridoids have found applications as hair dyes¹⁹.

The study reported here was an attempt to obtain an innovative cosmetic material. To achieve this goal, water and water/ethanol (1:1) extracts were prepared. Next, an iridoid fraction was isolated from them. The content of iridoids in each obtained fraction was determined, and an analysis of antioxidant properties was conducted. Moreover, each fraction was assessed to determine its effect on the irritant potential of model washing systems by determining the zein number. The test simulates the interaction that takes place between the cosmetics under study and the skin proteins.

MATERIAL AND METHODS

Chemicals

Biological activity tests were conducted using: DPPH (2,2-diphenyl-1-picrylhydrazyl, Sigma Aldrich), ABTS (2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid), Sigma Aldrich), 4-(dimetyloamino)benzaldehyde (Sigma Aldrich), aucubin (Sigma Aldrich), hydrochloric acid (36%, Chempur), ethyl alcohol (Heneywell), aluminium oxide (Sigma Aldrich), tyrosinase from mushroom (Sigma Aldrich), L-DOPA (Sigma Aldrich), hydroquinone (POCH Gliwice), di-potassium hydrogen phosphate pure p.a. (Chempur), potassium dihydrogen phosphate pure p.a. (POCH Gliwice). Measurements of the irritant potential were carried out with: zein from corn (Sigma Aldrich), sulfuric acid (98%, Chempur), copper sulphate pentahydrate (Chempur), potassium sulphate (Chempur), sodium hydroxide (Chempur), Tashiro indicator (Chempur). Model washing

systems were prepared from: sodium dodecyl sulphate (SLS, Sigma Aldrich) and sodium dodecyl ether sulphate (SLES, Brensurf, Brenntag). For a colorimetric analysis Aquamate Helios spectrophotometer were used.

Plant material

Dogwood (*Cornus mas L.* cultivar Bolestraszycki) fruits, collected in August 2016 in Bolestraszyce, were kindly supplied by Arboretum Bolesztraszyce.

Methods

Iridoid fraction derivation method

Isolation of the iridoid fraction, and determination of the total content of iridoids in the final product were performed using the method proposed by Baj et al.²⁰ with certain modifications.

To obtain appropriate product, freshly squeezed juice from pitted dogwood berries were used. The study was conducted using 25 g of dogwood berry juice which was extracted with 75 g of water (water extract, WE+IR). For the water/ethanol extract (WE+E+IR), extraction was performed in a water : ethanol mixture (1:1 w/w), with 75 g of extraction solvent in total. Both mixtures also contained 15 g of calcium carbonate each. The mixtures were boiled in a heating mantle under a reflux condenser for 30 minutes. The extracts thus prepared were decanted and filtered under reduced pressure through filter paper (Whatman no. 1). The resulting extracts were separated on a chromatography column filled with activated aluminium oxide (6 g). The first 5 mL portion of the filtrate was discarded.

Iridoid analysis

Analysis of total iridoid concentration

A 0.5 mL portion of the product was transferred to a 10 mL graduated tube, combined with 1 mL of methanol and 2 mL of 4-dimethylamino benzoic aldehyde solution with an addition of hydrochloric acid and 1.5 mL of distilled water. The mixture was heated for 3 minutes in boiling water bath, after which the test tube contents were immediately cooled down and filled with distilled water to 10 mL. After incubating the mixture at room temperature for 15 minutes absorbance was measured at the wavelength $\lambda = 540$ nm.

To calculate an amount of total iridoid concentration in final product, calibration curve of aucubin [0 – 100 mg/L] have been prepared with analogous procedure.

DPPH[•] radical scavenging activity

DPPH[•] radical scavenging by iridoid isolate was performed according to²¹. 1 mL of iridoid isolate or appropriate solvent was mixed with 1 mL 25 mM DPPH[•] solution in 96% ethanol. Following 40 min incubation at room temperature the absorbance of the sample was measured at $\lambda = 515$ nm using AquaMate spectrophotometer (Thermo Scientific). 96% ethanol was used as a blank sample. All samples were analyzed in triplicates. The percentage of DPPH[•] scavenging was calculated for each sample based on the equation:

$$\% \text{ of DPPH}^{\bullet} \text{ scavenging} = [1 - (As/Ac)] \times 100\%$$

where: As – absorbance of the sample; Ac – absorbance of the control sample (DPPH[•] solution).

ABTS^{•+} radical scavenging activity

Scavenging of ABTS^{•+} free radical was evaluated according to²². The scavenging reaction is based on decolourisation of the green ABTS radical cation (ABTS^{•+}). To prepare the ABTS^{•+} solution 19.5 mg ABTS and 3.3 mg potassium persulphate was mixed with 7 mL of phosphate buffer pH = 7.4 and dissolved for 16 hours in darkness. The solution was diluted to reach the absorbance at $\lambda = 414$ nm around 1.0. 20 μ L of iridoids fraction or appropriate solvent was mixed with 980 μ L diluted ABTS^{•+} solution and incubated for 10 min. The decrease in ABTS^{•+} absorbance was measured at $\lambda = 414$ nm using AquaMate spectrophotometer (Thermo Scientific), using distilled water as a blank. All samples were analyzed in triplicates. The percentage of ABTS^{•+} scavenging was calculated based on the equation:

$$\% \text{ of ABTS}^{\bullet+} \text{ scavenging} = [(1 - (As/Ac))] \times 100$$

where: As – absorbance of the sample; Ac – absorbance of the control sample (ABTS^{•+} solution).

Zein test

Irritant potential of the products was measured using zein test. In the surfactants solution zein protein is denatured and then is solubilized in the solution. This process simulates the behavior of surfactants in relation to the skin proteins. To 40 mL of the samples solution (10 wt%) was added 2 ± 0.05 g of zein from corn. The solutions with zein were shaken on a shaker with water bath (60 min at 35°C). The solutions were filtered on Whatman No. 1 filters and then centrifuged at 5000 rpm for 10 min. The nitrogen content in the solutions was determined by Kjeldahl method. 1 mL of the filtrate was mineralized in sulphuric acid (98%) containing copper sulphate pentahydrate and potassium sulphate. After mineralization the solution was transferred (with 50 mL of MiliQ water) into the flask of the Wagner–Parnas apparatus. 20 mL of sodium hydroxide (25 wt%) was added. The released ammonia was distilled with steam. Ammonia was bound by sulfuric acid (5 mL of 0.05 M H₂SO₄) in the receiver of the Wagner–Parnas apparatus. The unbound sulfuric acid was titrated with 0.1 M sodium hydroxide. Tashiro solution was used as an indicator. The zein number (ZN) was calculated from the equation:

$$ZN = (10 - V1) \cdot 100 \cdot 0.7 \text{ (mg N/100 mL)}$$

where V1 is the volume (mL) of sodium hydroxide used for titration of the sample.

The final result was the arithmetic mean of five independent measurements.

Patch test

The study was performed in a group of 20 healthy individuals (10 women and 10 men) aged between 20 and 40 years. None of the study participants had taken any medications for at least two weeks prior to the epidermal patch tests.

The epidermal patch tests were conducted using IQ Ultra test chambers from Chemotechnique Diagnostics. The aim of the patch test was to assess the irritant potential of water (WE+IR) and water/ethanol (WE+E+IR) isolates. An open test was also performed for an extract

containing no iridoids. 30 μ l portions of the test material were transferred into each chamber in accordance to the manufacturer's recommendations.

The epidermal patch tests were conducted in compliance with guidelines issued by the International Contact Dermatitis Research Group. Three identical patches containing the test emulsions were applied onto the skin of the test participants, in the region between the scapulas. The first patch was removed after 24 hours and the second (if no allergic reaction was observed) – after another 24 hours. Subsequent patches were removed 72, 96 and 168 hours after the application of the first patch test. The results were interpreted in accordance to the recognised international system for recording results of epidermal patch tests²³.

Tyrosinase inhibition activity

Tyrosinase inhibitory activity was determined by a spectrophotometric method described by Lim et. al²⁴ and modified by Studzińska-Sroka et. al²⁵. Briefly, 0.6 mL of isolate was mixed with 1.35 mL of 0.1 M of phosphate buffer (pH 6.8) and 0.75 mL of tyrosinase water solution (192 U/mL). Samples were incubated for 10 minutes in room temperature and after incubation 0.1 mL water solution of L-DOPA (4mM) was added. Absorbance of samples was measured after 20 minutes at $\lambda = 475$ nm. Each sample was accompanied by a reference that contains all components except L-DOPA, which was substituted with distilled water. The mixture of ingredients with distilled water in place of isolates was used as a control for reference. Hydroquinone was used as a standard as substance with well-known tyrosinase inhibitory properties. The percentage of tyrosinase inhibition was calculated as:

$$\frac{A_{control} - A_{sample}}{A_{control}} * 100\%$$

Presented values are the mean value from five independent measurements.

Statistical analysis

Each analysis was performed in triplicates. Obtained values were presented as mean \pm SD. Significant differences between obtained values were analyzed using GraphPad Prism 5.0 software using One-way ANOVA and Tukey's test. The differences were considered significant when $p < 0.05$.

RESULTS AND DISCUSSION

Multiple studies have shown that the content of iridoid compounds present in plant-based materials or extracts is directly correlated with their antioxidant potential²⁶⁻²⁸. Consequently, iridoids are a valuable material for a number of industries including the cosmetic industry. Dogwood berries, both fresh and processed, have been identified as a source of a broad spectrum of iridoid compounds at relatively high concentrations amounting to an average of 21.1 mg/g of fresh berry weight (assuming that berries contain 20% of fresh weight)²⁶⁻²⁹. The amount, however, is strictly related to a number of factors including plant variety, harvest time or degree of processing.

The process of isolation of iridoid compounds from the plant material is described in many procedures. In the majority of them, though, methanol is used as the elution solvent^{19,30}. However, the content of this alcohol markedly reduces the scope of potential applications of the extract as a material in the production of most cosmetics, and contributes to an increase in production costs due to the need to eliminate methanol. Consequently, using ethyl alcohol in the process of isolating the fraction rich in iridoid compounds seem to be a purposeful and justified strategy. Such an extract can then be added to cosmetic formulations as is, with ethanol used for isolation – ethanol is commonly present in cosmetics as a solvent, antibacterial agent or rheology modifier reducing the high viscosity of cosmetic products.

The results obtained in the study reveal that the use of water: ethanol (1:1) mixture to extract the iridoids allow to achieve a 4.5-fold increase in the concentration of iridoids in the final product relative to the product extracted with water only. The concentrations (expressed in miligram equivalents of aucubin per liter (mg(Aq)/L) are 69.9 mg/L for the water/ethanol isolate and 19.3 mg/L for the water fraction (Fig. 1).




Figure 1. Total concentration of iridoids expressed as milligram equivalents of aucubin [mg(Ac) / L]

Both tests investigating the antioxidant potential corroborated the thesis that the antioxidant capacity of solutions is directly proportional to their iridoid content^{26-29,31}. Within the studied range, the DPPH free-radical scavenging ability determined for the WE+E+IR was on average over 8.6% higher than for the WE+IR and for the ABTS radical it was nearly 1.5% higher. Importantly, for DPPH radicals the potential of the (WE+E+IR) was higher across the entire range of test concentrations. For the ABTS radical, a similar tendency was noted for the extract at concentrations exceeding approx. 1.5 mg/mL. What is interesting, at lower concentrations (between approx. 0.2 and 1.5 mg/mL) the ABTS radical scavenging ability was higher for the water isolate. The scavenging ability of both free radicals is shown in the charts below (Figure 2, 3).

The data given above demonstrate that the isolated iridoid fraction retains its high antioxidant potential despite performing the process of extraction at high temperature, which can lead to the decomposition of some antioxidants contained in fresh juice, e.g. vitamin C³²⁻³⁷. Whereas, as some authors report, the DPPH




Figure 2. Antioxidant properties of *Cornus mas* L. iridoid fractions against DPPH free radical




Figure 3. Antioxidant properties of *Cornus mas* L. iridoid fractions against ABTS free radical

free-radical scavenging ability for mashed dogwood berry is $73.30 \pm 4.07\%$, and for juice derived from *Cornus officinalis* it is $82.32 \pm 3.13\%$ ²⁶, the extracts obtained in the present study removed, respectively, $49.92\% \pm 3.21$ (for WE+IR) and $59.25\% \pm 3.82$ (for WE+E+IR) of the radicals at the concentration of 5 mg/mL. The relatively high antioxidant capacity of the obtained fraction, and its improved stability in relation to fresh juice, justify the use of the isolate as a material for the production of cosmetics.

The study involved an assessment of the capacity of iridoids isolated from dogwood berries to reduce the irritant potential of washing cosmetics. To this end, the irritant potential was determined for model systems based on washing compounds that are commonly used in the formulation of cosmetics, including sodium lauryl sulphate (SLS) and sodium laureth sulphate (SLES).

The literature data indicate that the irritant effect triggered by surfactants stems primarily from their interactions with the protein on the skin surface. The most potent irritant activity is ascribed to anionic surfactants which interact with proteins via ionic bonds. Nonionic and amphoteric washing agents exhibit no irritant effects because of weak hydrogen interactions with the surface protein^{38–42}.

The results indicate that iridoids have no significant impact on reducing the irritant activity of SLS and SLES. For pure solutions of these compounds, the results were

approx. 250 mg N/100 mL (SLS) and approx. 80 mg N/100 mL (SLES). The addition of the iridoid fraction isolated using water and water/ethanol causes a slight, falling within the margin of error, decrease in irritant activity both in SLS and SLES solutions. The results of tests determining the irritant potential support the classification of iridoids as active ingredients of body care cosmetics, in which they contribute to enhancing the antioxidant and nourishing effects (Fig. 4).




Figure 4. Zein value expressed in mgN/100 ml for 10% (w/w) surfactant solutions (SLS- sodium lauryl sulphate, SLES- sodium laureth sulphate), solutions with an addition of 5% (w/w) of ethanol EtOH instead of water and solutions with an 10% (w/w) addition of isolates

Patch tests are relatively cheap and uncomplicated to perform, which makes them one of the most widely used methods for the diagnosis of contact allergies. They are widely used for the evaluation of safety of cosmetic products. An assessment of the irritant reaction (IR) is usually performed using a 4-grade scale developed by Draeos and Dover⁴³.

The study reported above proves that the addition of WE+IR and WE+E+IR triggers no allergic responses in adults. Based on the study, it can thus be concluded that the extracts under investigation can be successfully used as an active ingredient in cosmetics.

Tyrosinase is a copper-containing enzyme which is a key enzyme in melanin biosynthesis and plays an important role in melanin pigmentation distribution⁴⁴. An accumulation of an abnormal amount of melanin in different specific parts of skin might result in more pigmented patches, which might be considered as an aesthetic problem. Bioactive compounds with strong tyrosinase inhibitory properties are valuable material for cosmetic industry – especially in field of anti-aging cosmetics. Our research proved that both isolates (WE+IR and WE+E+IR) are very effective tyrosinase inhibitors – the percentage of inhibition was up to two fold higher than for hydroquinone (44.2%) and were equal 83.4% for WE+IR and 88.9% for WE+E+IR (Fig. 5). Obtained values of inhibition are also significantly higher from values obtained by the other authors for water extracts from *Sambucus nigra* L. flower (62.5%), *Calendula* L. flower (55.3%) and *Euphrasia* L. herb (48.2%)²⁵. Results




Figure 5. Tyrosinase inhibition [%] for *Cornus mas* L. iridoid fractions in comparison with hydrochinone as well described tyrosinase inhibitor

shows that tyrosinase inhibitory properties of obtained isolates are relatively high, which make them effective cosmetic ingredients against skin hyperpigmentation.

CONCLUSION

The iridoid-containing extracts investigated in the study had a high antioxidant activity. The study demonstrated the water/ethanol isolate have a considerably higher antioxidant potential. Another finding was that the antioxidant capacity of solutions is directly proportional to the content of iridoids. The results of tests determining the safety of cosmetic use support the classification of iridoids as active ingredients of body care cosmetics, in which they contribute to enhancing the antioxidant and nourishing effects.

LITERATURE CITED

- Bulter, H. (2000). Poucher's perfumes, cosmetics and soaps (10th ed.). London, United Kingdom: Kluwer Academic Publishers.
- Elser, P. & Maibach, H. (2000). Cosmeceuticals and active cosmetics. New York, USA: Taylor & Francis Group.
- Wang, W., Sun, F., An, Y., Ai, H., Zhang L., Huang, W. & Li, L. (2009). Morroniside protects human neuroblastoma SH-SY5Y cells against hydrogen peroxide-induced cytotoxicity. *Eur. J. Pharmacol.* 613(1–3), 19–23. DOI :10.1016/j.ejphar.2009.04.013.
- Seeram, N.P. & Nair, M.G. (2002). Inhibition of lipid peroxidation and structure-activity-related studies of the dietary constituents anthocyanins, anthocyanidins, and catechins. *J. Agric. Food Chem.* 50(9), 5308–5312. DOI: 10.1021/jf025671.
- Vareed, S.K., Reddy, M.K., Schutzki, R.E. & Nair, M.G. (2006). Anthocyanins in *Cornus alternifolia*, *Cornus controversa*, *Cornus kousa* and *Cornus florida* fruits with health benefits. *Life Sci.* 11,78(7), 777–784. DOI: 10.1016/j.lfs.2005.05.094.
- Kyriakopoulos, A.M. & Dinda, B. (2015). *Cornus mas* (Linnaeus) novel devised medicinal preparations: bactericidal effect against *Staphylococcus aureus* and *Pseudomonas aeruginosa*. *Molecules* 20, 11202–11218. DOI:10.3390/molecules200611202.
- Tural, S. & Koca, I. (2008). Physicochemical and antioxidant properties of Cornelian Cherry fruits (*Cornus mas* L.) grown in Turkey. *Sci. Hort.* 116, 362–366. DOI: 10.1016/J.Scienta.2008.02.003.
- Klimenko, S.W. (2004). The cornelian cherry (*Cornus mas* L.) collection, preservation and utilization of genetic resources. *J. Fruit Orn. Plant Res.* 12, 93–98. INRMM:13555513.

- Pawlowska, A.M., Camangi, F. & Braca, A. (2010). Quali-quantitative analysis of flavonoids of *Cornus mas* L. (*Cornaceae*) fruits. *Food Chem.* 119, 1257–1261. DOI: 10.1016/j.foodchem.2009.07.063.

- Niemmen, M., Suomi, J., Nouhuys, S., Sauri, P. & Riekkola, M.L. (2003). Effect of iridoid glycoside content on oviposition host plant choice and parasitism in a specialist herbivore. *J. Chem. Ecol.* 29(4). ID: 12775146.

- Vodovotz, Y., Constantine, G., Rubin, J., Csete, M., Voit E.O. & An G. (2009). Mechanistic simulations of inflammation: current state and future prospects. *Math. Biosci.* 217, 1–10. DOI: 10.1016/j.mbs.2008.07.013.

- Calixto, J.B., Campos, M.M., Otuki, M.F. & Santos, A.R.S. (2004). Anti-inflammatory compounds of plant origin. Part II. Modulation of pro-inflammatory cytokines, chemokines and adhesion molecules. *Planta Med.* 70, 93–103. DOI: 10.1055/s-2004-815483.

- Háznagy-Radnai, E., Balogh, A., Czige, S., Máthé, I., Hohmann, J. & Blazsó, G. (2011). Antiinflammatory activities of hungarian stachys species and their iridoids. *Phytother Res.* 26(4), 505–509. DOI: 10.1002/ptr.3582.

- Masuda, M., Itoh K., Murata K., Naruto S., Uwaya A., Isami, F. & Matsuda, H. (2012). Inhibitory effects of *Morinda citrifolia* extract and its constituents on melanogenesis in murine B16 melanoma cells. *Biol. Pharm. Bull.* 35(1), 7–83. DOI: 10.2145/biol.2012.3254.

- Akihisa, T., Seino, K., Kaneko, E., Watanabe, K., Tochizawa, S., Fukatsu, M., Banno, N., Metori, K. & Kimura, Y. (2010). Melanogenesis inhibitory activities of iridoid, hemiterpene, and fatty acid-glycosides from the fruits of *Morinda citrifolia* (Noni). *J. Oleo Sci.* 59(1), 49–57. DOI: 10.5650/jos.59.49.

- Saracoglu, I. & Harput, U.S. (2012). In vitro cytotoxic activity and structure activity relationships of iridoid glucosides derived from *Veronica species*. *Phytother. Res.* 26(1), 148–152. DOI: 10.1002/ptr.3546.

- Saracoglu, I., Oztunca, F.H., Nagatsu, A. & Harput, U.S. (2011). Iridoid content and biological activities of *Veronica cuneifolia subsp. cuneifolia* and *Veronica cymbalaria*. *Pharm. Biol.* 49(11), 1150–1157. DOI: 10.3109/13880209.2011.575790.

- Es Haghi, M., Dehghan, G., Banihabib, N., Zare, S., Mikaili, P. & Panahi, F. (2012). Protective effects of *Cornus mas* fruit extract on carbon tetrachloride induced nephrotoxicity in rats. *Indian J. Nephrol.* 24(5), 291–296. DOI: 10.4103/0971-4065.133000.

- Koichiro, K. & Shoichi, H. (1992). U.S. Patent no. 5,078,750. Washington, D.C.: U.S. Patent and Trademark Office.

- Baj, T., Kołtunowska, D., Głowniak, K. & Wolski, T. (2015). Determination of aucubin in aerial parts of *Buddleja davidii* Franch. using different TLC-detection methods. XX. 46 symposium on essential oils. Lublin, Poland, Annales Universitatis Mariae Curie-Skłodowska.

- Brand-Williamis, W., Cuvelier, M. & Berset, C. (1995). Use of a free radical method to evaluate antioxidant activity. *LWT Food Sci. Technol.* 28, 25–30. DOI: 0023-6438/95/010025.

- Re, R., Pellegrini, N., Protegente, A., Pannala, A., Yang, M. & Rice-Evans, C. (1999). Antioxidant activity applying and improved ABTS radical cation decolorization assay. *Free Radic. Biol. Med.* 26, 1231–1237. DOI: 0891-5849(98)00315-3.

- Płocica, J., Tal-Figiel, B. & Figiel, W. (2014). Badania reologiczne i sensoryczne stosowane do oceny preparatów kosmetycznych. *Ś.P.K.* 17(1), 68–73.

- Lim, T.Y., Lim, Y.Y. & Yule, C.M. (2009). Evaluation of antioxidant, antibacterial and anti-tyrosinase activities of four *Macaranga* species. *Food Chem.* 114, 594–599. DOI: 10.1016/j.foodchem.2008.09.093.

- Studzińska-Sroka, E., Frątczak, A. & Bylka, W. (2016). Evaluation of properties inhibiting tyrosinase by selected plant extracts. *Pol. J. Cosmetol.* 19(1), 51–55.

- West, B.J., Deng, S., Jensen, J., Palu, A.K. & Berrio, F.L. (2012). Antioxidant, toxicity, and iridoid tests of pro-

cessed cornelian cherry fruits. IJFST. 7, 1392–1397. DOI: 10.1111/j.1365-2621.2012.02985.

27. Kwak, H., Kim, H.J., Lee, K.H., Kang, S.C. & Zee, O.P. (2009). Antioxidative iridoid glycosides and phenolic compounds from *Veronica peregrina*. *Jong. Arch. Pharm. Res.* 32(2), 207–213. DOI: 10.1007/s12272-009-1137-x.

28. Wael, M., Abdel-Mageed, A., Enaam, Y., Backheet, A. Azza, A., Khalifa, A. Zedan, Z., Ibraheim, A. & Samir, A. (2007). Antiparasitic antioxidant phenylpropanoids and iridoid glycosides from *Tecoma mollis*. *Fitoterapia* 83(3), 500–507. DOI: 10.1016/j.fitote.2011.12.025.

29. Kucharska, M. (2012). *Związki aktywne owoców derenia*. Wrocław, Poland, Uniwersytet Przyrodniczy we Wrocławiu.

30. Xiao, N.M., Xue, X.Y., Feng, J.T., Zhang, X.L. & Liang, X.M. (2011). Isolation and purification of unstable iridoid glucosides from traditional chinese medicine by preparative high performance liquid chromatography coupled with solid-phase extraction. *Chem. Res. Chinese Univ.* 27(3), 392–396. DOI: 1005-9040(2011)-03-392-05.

31. Norlia, M, Siti, A.M, Mashitah, M.Y. & Jolius, G. (2014). Influence of solvent polarity and conditions on extraction of antioxidant, flavonoids and phenolic content from *Averrhoa bilimbi*. *J. Food Sci. Eng.* 4, 255–260. DOI: 10.17265/2159-5828/2014.05.006.

32. Dai, J. & Mumper, R.J. (2010). Plant phenolics: extraction, analysis and their antioxidant and anticancer properties. *Molecules* 21:15(10), 7313–7352. DOI: 10.3390/molecules15107313.

33. Pacifico, S., D'Abrosca, B., Pascarella, M.T., Letizia, M., Uzzo, P., Piscopo, V. & Fiorentino, A. (2009). Antioxidant efficacy of iridoid and phenylethanoid glycosides from the medicinal plant *Teucrium chamaedris* in cell-free systems. *Bioorg. Med. Chem.* (17), 6173–6179. DOI: 10.1016/j.bmc.2009.07.065.

34. Polinicencu, C., Popescu, H. & Nistor, C. (1980). Vegetal extracts for cosmetic use: extracts from fruits of *Cornus mas*. Preparation and characterization. *Cluj. Med.* 53, 160–163.

35. Pantelidis, G.E., Vasilakakis, M., Manganaris, G.A. & Diamantidis, G.R. (2007). Antioxidant capacity, phenol, anthocyanin and ascorbic acid contents in raspberries, blackberries, red currants, gooseberries and cornelian cherries. *Food Chem.* 102, 777–783. DOI: 10.1016/j.foodchem.2006.06.021.

36. Demir, F. & Kalyoncu, I.H. (2003). Some nutritional, pomological and physical properties of cornelian cherry (*Cornus mas* L.). *J. Food Eng.* 60, 335–341. DOI: 10.1016/S0260-8774(03)00056-6.

37. Celep, E., Aydın, A. & Yesilada, E. (2012). A comparative study on the in vitro antioxidant potentials of three edible fruits: cornelian cherry, Japanese persimmon and cherry laurel. *Food. Chem. Toxicol.* 50, 3329–3335. DOI: 10.1016/j.fct.2012.06.010.

38. Jackson, C.T., Paye, M. & Maibach, H. (2014). Mechanism of skin irritation by surfactants and anti-irritants for surfactants base products (pp. 353–358). In Barel, A., Paye, M. & Maibach, H. *Handbook of cosmetic science and technology*. Fourth edition. Bosta Rocon. USA: CRC Press Taylor & Francis Group.

39. Lips, A., Ananthapadmanabhan, K.P., Vethamuthu, M., Hua, X.Y., Yang, L., Vincent, C., Deo N. & Somasundaran, P. (2007). Role of surfactant micelle charge in protein denaturation and surfactant-induced skin irritation, (pp. 177–189). In Rhein, L., Schlossman, M., O'Lenick, A. & Somasundaran, P. *Surfactants in personal care products and decorative cosmetics third edition*. Bosta rocon, USA: CRC Press Taylor & Francis Group.

40. Agner, T. & Serup, J. (1990). Sodium lauryl sulphate for irritant patch testing—A dose-response study using bioengineering methods for determination of skin irritation. *J. Invest. Dermatol.* 95, 543–547. DOI: 022-202X/90/S03.50.

41. Polefka, T. (1999). Surfactants interaction with skin. In Broze, G. *Handbook of detergents. Part A: Properties*. Surf. Sci. Ser. Vol. 82. Marcel Dekker Publ.

42. Bujak, T., Wasilewski, T. & Nizioł-Łukaszewska, Z. (2015). Role of macromolecules in the safety of use of body wash cosmetics, *Coll. Surf. B.* 135, 497–503. DOI: 10.1016/j.colsurfb.2015.07.051.

43. Draelos, Z.D. & Dover, J.S. (2011). *Kosmeceutyki*. Wrocław, Poland: Elsevier, Urban&Partner.

44. Kima, Y.J. & Uyamab, H. (2005). Tyrosinase inhibitors from natural and synthetic sources: structure, inhibition mechanism and perspective for the future. *Cell. Mol. Life Sci.* 62, 1707–1712. DOI: 10.1007/s00018-005-5054-y.