


This article was downloaded by: [185.55.64.226]

On: 15 March 2015, At: 01:32

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



International Journal of Occupational Safety and Ergonomics

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/tose20>

Estimation of Aerobic Capacity and Determination of Its Associated Factors Among Male Workers of Industrial Sector of Iran

Hadi Daneshmandi^a, Abdolreza Rajae Fard^a & Alireza Choobineh^b

^a School of Health, Shiraz University of Medical Sciences, Shiraz, Iran

^b Research Center for Health Science, Shiraz University of Medical Sciences, Shiraz, Iran

Published online: 08 Jan 2015.

To cite this article: Hadi Daneshmandi, Abdolreza Rajae Fard & Alireza Choobineh (2013) Estimation of Aerobic Capacity and Determination of Its Associated Factors Among Male Workers of Industrial Sector of Iran, *International Journal of Occupational Safety and Ergonomics*, 19:4, 667-673

To link to this article: <http://dx.doi.org/10.1080/10803548.2013.11077014>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

Estimation of Aerobic Capacity and Determination of Its Associated Factors Among Male Workers of Industrial Sector of Iran

Hadi Daneshmandi
Abdolreza Rajaei Fard

School of Health, Shiraz University of Medical Sciences, Shiraz, Iran

Alireza Choobineh

Research Center for Health Science, Shiraz University of Medical Sciences, Shiraz, Iran

Introduction. The aim of this study was to estimate maximal aerobic capacity ($V_{O_{2max}}$), to determine its associated factors among workers of industrial sector of Iran and to develop a regression equation for subjects' $V_{O_{2max}}$. **Methods.** In this study, 500 healthy male workers employed in Shiraz industries participated voluntarily. The subjects' $V_{O_{2max}}$ was assessed with the ergocycle test according to the Åstrand protocol. Required data was collected with a questionnaire covering demographic details (i.e., age, job tenure, marital status, education, nature of work, shift work, smoking and weekly exercises). **Results.** The subject's mean $V_{O_{2max}}$ was 2.69 ± 0.263 L/min. The results showed that there was an association between $V_{O_{2max}}$ and age, BMI, hours of exercise and smoking, but there was no association between $V_{O_{2max}}$ and height, weight, nature of work and working schedule. On the basis of the results, regression equations were developed to estimate $V_{O_{2max}}$. **Conclusion.** Final regression equation developed in this study may be used to estimate $V_{O_{2max}}$ reliably without the need to use other laboratory instruments for aerobic measurement.

ergocycle test Åstrand protocol aerobic capacity $V_{O_{2max}}$

1. INTRODUCTION

Heavy and dynamic physical work is still common in many industrial activities such as mining, building and agriculture [1, 2]. In the developed countries, 10–20% of workers are engaged in muscular demanding jobs, while in the developing countries all kinds of intensive work are common [3]. Unbalanced relationship between physical demands and workers' capacities caused adverse health outcomes [1, 4, 5]. In physically demanding jobs, the aerobic capacity (V_{O_2}) of workers determines man's power productivity [6]. Assessing V_{O_2} is essential to fit the job to the

worker's physiological capacity and to prevent over strain and negative consequences of demanding muscular exertion of daily tasks. Knowledge about the level of physical work capacity is inevitable to ensure physiologically safe working conditions.

The capacity of performing prolonged dynamic work is determined with V_{O_2} , which is assessed by $V_{O_{2max}}$ [7, 8]. According to some studies, general health status and physical exercise have positive influence on the physical capacity [7, 8, 9], whereas age has negative effects [1, 8, 10, 11, 12, 13].

This article was extracted from the thesis written by Mr. Hadi Daneshmandi, MSc., a student of Ergonomics and was financially supported by Shiraz University of Medical Sciences grant 89-5301. The authors would like to thank Dr. H.R. Mostafavie and Mrs. A Hosseini from Nader Kazemi clinic for their valuable cooperation in the stage of data gathering and $V_{O_{2max}}$ measurement.

Correspondence should be sent to Alireza Choobineh, Department of Ergonomics, School of Health, Shiraz University of Medical Sciences, PO Box 71645-11, Shiraz, Iran. E-mail: alrchoobin@sums.ac.ir.

The best method to measure $V_{O_{2max}}$ is a direct measurement of V_{O_2} in the maximal exercise test, which is frequently used by athletes [14]. For industrial workers, much more suitable is indirect method estimating $V_{O_{2max}}$, which puts less physiological pressure on the subject [15].

There have been few studies in Iran on estimating physical capacity of different population groups, especially workers of industrial sector, but there is no national data on physiological characteristics. Therefore, this study was conducted to a) establish the level of V_{O_2} , b) determine its associated factors among industrial workers of Shiraz and c) to develop regression equation to estimate the subjects' V_{O_2} . The findings of this study can help to fit the job to the workers' characteristics and to select personnel during the process of employment.

2. MATERIAL AND METHODS

2.1. Study Subjects

Subjects of this cross-sectional study were 500 healthy, occupationally active, blue-collar male workers of industrial sectors of Shiraz. The study was conducted from October 2010 to February 2011. The subjects were randomly selected from those referred to a governmental clinic for periodic medical examinations. Each day, researchers randomly selected 10 workers out of 40 from 50 different factories, who came to the clinic. All subjects participated in the study voluntarily and received information on the aims and protocol of the study. The study was conducted in accordance with the Helsinki Declaration of 1964 as revised in 1984. All subjects signed a consent form before the study. The study was reviewed and approved by Shiraz University of Medical Sciences ethics committee.

2.2. Measurements

2.2.1. Demographic characteristics

The subjects filled a self-administered questionnaire with questions on demographics and job related variables (i.e., age, job title, job tenure,

marital status, education, static or dynamic nature of work, shift work, smoking and weekly exercise hours).

2.2.2. $V_{O_{2max}}$

The subjects performed a 6-min sub-maximal exercise test on a cycle-ergometer (Monark, Sweden) based on the Åstrand protocol to calculate V_{O_2} [16]. During the test, heart rate was monitored and a wireless transmitter transmitted the heart beat data to the base unit. The subjects' $V_{O_{2max}}$ (expressed both in L/min and ml/min/kg) was calculated with obtained data and a special software. The tests were performed in the laboratory, between 8 am and 2 pm. The average temperature was 20.5 °C, relative humidity 41% and barometric pressure 857 hPa.

2.2.3. Height and weight

The subjects' height was measured with a tape in the standard position (Pakhsh Abzar, Iran) [16]. The subjects' weight was measured with a digital scales (Beurer, Germany). During the measurements, the subjects wore light clothing only. Body mass index (BMI) was calculated for each subject.

2.3. Data Analysis and Statistical Procedures

SPSS version 16 was used for statistical analyses. Nonparametric analyses were applied because the data was not normally distributed. Kruskal–Wallis and Mann–Whitney U tests were used to compare the mean values of $V_{O_{2max}}$ in age, weight, height and BMI groups. A linear regression analysis examined relationship between $V_{O_{2max}}$ and different variables separately, and a multiple regression analysis developed the overall regression equation for $V_{O_{2max}}$ estimation in the study population; $p < .05$ was significant.

3. RESULTS

Table 1 presents demographic characteristics of the subjects. Table 2 presents mean (M), standard deviation (SD), minimum and maximum V_{O_2} .

TABLE 1. Characteristics of Subjects (N = 500)

Variable	M (SD)	Range
Age (years)	32.01 (7.66)	20–59
Weight (kg)	77.10 (12.41)	50–110.2
Height (cm)	176.10 (5.93)	159–190
BMI	24.82 (3.58)	15.90–33.57
Job tenure (years)	8.50 (6.27)	0.17–30
Exercise per week (h)	2.95 (3.49)	0–12

Variable	n (%)
Marital status	
single	104 (20.8)
married	396 (79.2)
Education	
primary	134 (26.8)
secondary	258 (51.6)
tertiary	108 (21.6)
Nature of work	
static	255 (51)
dynamic	245 (49)
Type of industry	
chemical	177 (35.4)
food	167 (33.4)
other ¹	156 (31.2)
Work shift	
rotating	234 (46.8)
fixed day	266 (53.2)
Cigarette smoking	
smoker	60 (12)
nonsmoker	440 (88)

Notes. 1 = including metal structure manufacturing, construction, textile and wood industries. BMI = body mass index.

TABLE 2. Aerobic capacity (V_{O2}) of Subjects (N = 500)

V _{O2}	M	SD	Range
L/min	2.69	0.263	1.71–3.50
ml/min/kg	35.95	7.390	19.50–58.04

TABLE 3. Aerobic Capacity (V_{O2}) by Age (N = 500)

Age (years)	V _{O2max} (ml/min/kg)			V _{O2max} (L/min)		
	M	SD	Range	M	SD	Range
20–29 (n = 227)	38.82	7.30	23.91–58.04	2.84	0.165	2.45–3.50
30–39 (n = 198)	34.78	6.59	22.22–57.40	2.66	0.183	2.00–2.97
40–49 (n = 55)	31.14	5.21	22.26–45.50	2.38	0.241	1.78–2.73
50–59 (n = 20)	27.97	5.10	19.50–35.74	2.13	0.27	1.71–2.56

Notes. V_{O2max} = maximal aerobic capacity.

Table 3 presents V_{O2} of the subjects by age groups. The results show that V_{O2} decreases with age. Kruskal–Wallis analysis reveals that mean values of V_{O2max} in the age groups are significantly different (*p* < .001). Moreover, Mann–Whitney *U* test shows that mean values of V_{O2max} of each pair of the age groups are significantly different (*p* < .001).

Linear regression analysis reveals a reverse relationship between V_{O2max} and age (*r* = −.796). It is defined with Equation 1 (Figure 1):

$$V_{O2max} \text{ (L/min)} = (-0.027 \times \text{age}) + 3.572. \quad (1)$$

Table 4 presents V_{O2} by BMI [17]. The subject with normal BMI (18.5–24.9) has the highest value of V_{O2max}. Statistical analysis shows that mean V_{O2max} values in the BMI groups are significantly different (*p* < .001). Mann–Whitney *U* test also shows significant differences in V_{O2} between normal and overweight groups (*p* < .001), and normal and obese groups (*p* = .005).

The linear regression analysis reveals relationship between V_{O2max} and BMI (*r* = −.158). It is defined with Equation 2 (Figure 2):

$$V_{O2max} \text{ (L/min)} = (-0.012 \times \text{BMI}) + 2.982. \quad (2)$$

According to Equation 2, V_{O2} declines when BMI increases. The linear regression analysis also shows a direct relationship between V_{O2max} and weekly exercise hours (*r* = .37) (Figure 3).

The multiple regression analysis reveals that there is a strong relationship between V_{O2max}, age, BMI and weekly exercises (*r* = .818). It is defined with Equation 3:

$$V_{O2max} \text{ (L/min)} = (-0.026 \times \text{age}) - (0.002 \times \text{BMI}) + (0.014 \times \text{exercise}) + 3.516. \quad (3)$$

Downloaded by [185.55.64.226] at 01:32 15 March 2015




Figure 1. Relationship between aerobic capacity (V_{O_2}) and age of subjects ($N = 500$).




Figure 2. Relationship between aerobic capacity (V_{O_2}) and body mass index (BMI) of subjects ($N = 500$).




Figure 3. Relationship between aerobic capacity (V_{O_2}) and weekly exercise of subjects ($N = 500$).

Downloaded by [185.55.64.226] at 01:32 15 March 2015

TABLE 4. Aerobic Capacity (V_{O_2}) by BMI ($N = 500$)

BMI [17]	V_{O_2max} (L/min)			V_{O_2max} (ml/min/kg)		
	M	SD	Range	M	SD	Range
Underweight ($n = 23$)	2.62	0.30	1.91–2.96	48.73	6.57	34.11–58.04
Normal ($n = 245$)	2.74	0.26	1.78–3.50	49.17	6.18	33.12–58.09
Overweight ($n = 196$)	2.65	0.24	1.71–3.24	31.64	3.57	22.22–42.16
Obese ($n = 36$)	2.61	0.27	1.97–2.97	26.21	3.29	19.50–30.90

Notes. BMI = body mass index; underweight = BMI <18.5, normal = BMI 18.5–24.9, overweight = BMI 25–29.9, obese = BMI ≥30; V_{O_2max} = maximal aerobic capacity.

TABLE 5. Aerobic capacity (V_{O_2}) by Work Properties and Smoking Habit ($N = 500$)

Variable	V_{O_2max} (L/min)		V_{O_2max} (ml/min/kg)		p^*
	M	SD	M	SD	
Nature of work					
static ($n = 255$)	2.68	0.29	35.67	7.75	.668
dynamic ($n = 245$)	2.70	0.23	36.23	7.00	
Work shift					
rotating ($n = 234$)	2.69	0.23	35.69	6.87	.552
fixed day ($n = 266$)	2.69	0.28	36.17	7.82	
Cigarette smoking					
smoker ($n = 60$)	2.57	0.28	36.94	7.28	< .001
nonsmoker ($n = 440$)	2.70	0.25	37.81	7.40	

Notes. * = Independent Mann–Whitney U test between two groups.

Table 5 presents the result of V_{O_2} on the basis of the nature of work, working schedule and smoking habit. Independent Mann–Whitney U test shows that only smoking has influence on V_{O_2} . The mean value of V_{O_2max} is significantly lower among smokers than nonsmokers ($p < .001$).

Further experiments were conducted to check to what extent the final regression equation could estimate V_{O_2} . The value of V_{O_2} of 10 workers (selected from the subjects of this study) was estimated with developed regression equation. The value of V_{O_2max} of 10 workers was measured with the Åstrand protocol [16]. Compared results of statistical analysis indicates that the mean values

of V_{O_2max} obtained with the two methods are not significantly different ($p > .05$) (Table 6).

4. DISCUSSION

The subjects of the study were young (32.01 ± 7.66 years) and tall (176.10 ± 5.93 cm). The subjects were male workers with BMI within the normal range (24.82 ± 3.58). Mean value of V_{O_2max} was 2.69 ± 0.263 L/min (35.95 ± 7.39 ml/min/kg).

The value of V_{O_2} measured in this study was close to the findings of Tuxworth and Shahnavaaz. In their study, V_{O_2max} of 45 Iranian male workers of a steel corporation was measured with the step test and it was 2.65 L/min [18]. In contrast, V_{O_2} of the present study subjects was lower than V_{O_2} of the trained men who's V_{O_2max} was 4.451 ± 0.629 L/min [19], and of the Netherlands male nonsmokers who's V_{O_2max} was 3.89 ± 0.92 L/min [20].

TABLE 6. V_{O_2} Measured with Final Regression Equation and Åstrand Protocol ($n = 10$)

Method	V_{O_2max} (L/min)		p^*
	M	SD	
Åstrand protocol	2.49	0.41	.348
Regression equation	2.58	0.14	

Notes. * = paired-sample t test.

Additionally, the linear regression analysis showed significant relationship between $V_{O_{2max}}$ and age, BMI and weekly exercises. These findings are similar to the findings of other studies [10, 11, 12, 13, 21, 22, 23, 24]. The developed final regression equation showed these relationships and made it possible to estimate V_{O_2} of any individual with the smaller number of measurements.

Similarly to the results of Betik and Hepple's study, smoking reduced V_{O_2} among workers as compared with their nonsmoker colleagues [24]. Moreover, $V_{O_{2max}}$ among workers with exercise habit was significantly higher than among those who did not exercise. This finding is in line with the previous study [25]. The findings of the present study revealed that work schedule had no association with V_{O_2} and confirmed the results of the previous study [26].

Estimation of V_{O_2} with the final regression equation and the Åstrand protocol yielded similar results. This indicated that the regression equation developed in this study might be used to measure $V_{O_{2max}}$.

5. CONCLUSION

There is a relationship between $V_{O_{2max}}$ and age, BMI, hours of exercise and smoking habit, but weight, height, nature of work and working schedule have no influence on V_{O_2} . The final regression equation developed in this study shows relationship between $V_{O_{2max}}$ and age, BMI and exercise, and may be used to estimate V_{O_2} without using additional laboratory instruments for cardiorespiratory capacity measurement.

REFERENCES

1. Yoopat P, Toicharoen P, Boontong S, Glinsukon T, Vanwongerghem K, Louhevaara V. Cardiorespiratory capacity of Thai workers in different age and job categories. *J Physiol Anthropol Appl Human Sci*. 2002; 21(2):121–8.
2. Åstrand PO, Rodahl K, Dahl HA, Strømme SB. *Textbook of work physiology: physiological bases of exercise*. 4th ed. Champaign, IL, USA: Human Kinetics; 2003.
3. Louhevaara V, Kilbom Å. Dynamic work assessment. In: Wilson JR, Corlett N, editors. *Evaluation of human work*. 3rd ed. Boca Raton, FL, USA: CRC Press; 2005. p. 429–52.
4. Kumashiro M. Practical measurement of psychophysiological functions for determining workloads. In: Wilson JR, Corlett N, editors. *Evaluation of human work*. 3rd ed. Boca Raton, FL, USA: CRC Press; 2005. p. 605–28.
5. Holtermann A, Jørgensen MB, Gram B, Christensen JR, Faber A, Overgaard K, et al. Worksite interventions for preventing physical deterioration among employees in job-groups with high physical work demands: background, design and conceptual model of FINALE. *BMC Public Health*. 2010; 10:120. Retrieved September 24, 2013, from: <http://www.biomedcentral.com/1471-2458/10/120>.
6. Ramsey JD, Burford CL, Beshir MY, Jensen RC. Effect of workplace thermal conditions on safe work behavior. *J Safety Res*. 1983;14(3):105–14.
7. Bugajska J, Makowiec-Dąbrowska T, Bortkiewicz A, Gadzicka E, Marszałek A, Lewandowski Z, Konarska M. Physical capacity of occupationally active population and capability to perform physical work. *International Journal Occupational Safety and Ergonomics (JOSE)*. 2011;17(2): 127–8. Retrieved September 24, 2013, from: <http://www.ciop.pl/43470>.
8. Huggett DL, Connelly DM, Overend TJ. Maximal aerobic capacity testing of older adults: a critical review. *J Gerontol A Biol Sci Med Sci*. 2005;60(1):57–66.
9. Guyton AC, Hall JE. *Textbook of medical physiology*. 11th ed. Philadelphia, PA, USA: Saunders Elsevier; 2006.
10. Tanaka H, Desouza CA, Jones PP, Stevenson ET, Davy KP, Seals DR. Greater rate of decline in maximal aerobic capacity with age in physically active vs. sedentary healthy women. *J Appl Physiol*. 1997;83 (6): 1947–53. Retrieved September 24, 2013, from: <http://jap.physiology.org/content/83/6/1947.long>.
11. Beere PA, Russell SD, Morey MC, Kitzman DW, Higginbotham MB. Aerobic exercise training can reverse age-related

- peripheral circulatory changes in healthy older men. *Circulation*. 1999;100(10): 1085–94. Retrieved September 24, 2013, from: <http://circ.ahajournals.org/content/100/10/1085.long>.
12. Schiller BC, Casas YG, Desouza CA, Seals DR. Maximal aerobic capacity across age in healthy Hispanic and Caucasian women. *J Appl Physiol*. 2001;91(3): 1048–54. Retrieved September 24, 2013, from: <http://jap.physiology.org/content/91/3/1048.long>.
 13. Ładyga M, Faff J, Burkhard-Jagodzińska K. Age-related decrease of the indices of aerobic capacity in the former elite rowers and kayakers. *Biol Sport*. 2008;25(3): 245–60. Retrieved September 24, 2013, from: <http://biolsport.com/abstracted.php?level=5&ICID=890322>.
 14. Ładyga M, Faff J. Assessment of the accuracy of prediction of the maximal oxygen uptake based on submaximal exercises in the former elite rowers and paddlers. *Biol Sport*. 2005; 22(2):125–34. Retrieved September 24, 2013, from: <http://biolsport.com/abstracted.php?level=5&ICID=891539>.
 15. Tayyari F, Smith JL. Occupational ergonomics: principles and applications. London, UK: Chapman & Hall; 1997.
 16. Dwyer GB, Davis SE. ACSM's health-related physical fitness assessment manual. 2nd ed. Hagerstown, MD, USA: Lippincott Williams & Wilkins; 2008.
 17. Lee RD, Nieman DC. Nutritional assessment. 3rd ed. New York, NY, USA: McGraw-Hill; 2003.
 18. Tuxworth W, Shahnawaz H. The design and evaluation of a step test for the rapid prediction of physical work capacity in an unsophisticated industrial work force. *Ergonomics*. 1977;20(2):181–91.
 19. Malek MH, Housh TJ, Berger DE, Coburn JW, Beck TW. A new non-exercise-based $\dot{V}_{O_{2max}}$ prediction equation for aerobically trained men. *J Strength Cond Res*. 2005;19(3):559–65.
 20. Zoladz JA, Duda K, Majerczak J. \dot{V}_{O_2} /power output relationship and the slow component of oxygen uptake kinetics during cycling at different pedaling rates: relationship to venous lactate accumulation and blood acid-base balance. *Physiol Res*. 1998;47(6):427–38. Retrieved September 24, 2013, from: http://www.biomed.cas.cz/physiolres/pdf/47/47_427.pdf.
 21. Hepple RT, Hagen JL, Krause D. Oxidative capacity interacts with oxygen delivery to determine maximal O_2 uptake in rat skeletal muscles *in situ*. *J Physiol*. 2002;541(Pt 3): 1003–12. Retrieved September 24, 2013, from: <http://jp.physoc.org/content/541/3/1003.long>.
 22. Uth N, Sørensen H, Overgaard K, Pedersen PK. Estimation of $\dot{V}_{O_{2max}}$ from the ratio between HR_{max} and HR_{rest} —the heart rate ratio method. *Eur J Appl Physiol*. 2004;91(1): 111–5.
 23. Grassi GP, Turci M, Sforza C. Aerobic fitness and somatic growth in adolescents: a cross sectional investigation in a high school context. *J Sports Med Phys Fitness*. 2006;46(3):412–8.
 24. Betik AC, Hepple RT. Determinants of $\dot{V}_{O_{2max}}$ decline with aging: an integrated perspective. *Appl Physiol Nutr Metab*. 2008; 33(1):130–40.
 25. Byars A, Greenwood M, Greenwood L, Simpson WK. The effectiveness of a pre-exercise performance drink (PRX) on indices of maximal cardiorespiratory fitness. *J Int Soc Sports Nutr*. 2006;3:56–9.
 26. Virtanen M, Vahtera J, Pentti J, Honkonen T, Elovainio M, Kivimäki M. Job strain and psychologic distress influence on sickness absence among Finnish employees. *Am J Prev Med*. 2007;33(3):182–7.