

## EXISTENCE OF POSITIVE RADIAL SOLUTIONS TO A $p$ -LAPLACIAN KIRCHHOFF TYPE PROBLEM ON THE EXTERIOR OF A BALL

John R. Graef, Doudja Hebboul, and Toufik Moussaoui

*Communicated by Vicentiu D. Rădulescu*

**Abstract.** In this paper the authors study the existence of positive radial solutions to the Kirchhoff type problem involving the  $p$ -Laplacian

$$-\left(a + b \int_{\Omega_e} |\nabla u|^p dx\right) \Delta_p u = \lambda f(|x|, u), \quad x \in \Omega_e, \quad u = 0 \text{ on } \partial\Omega_e,$$

where  $\lambda > 0$  is a parameter,  $\Omega_e = \{x \in \mathbb{R}^N : |x| > r_0\}$ ,  $r_0 > 0$ ,  $N > p > 1$ ,  $\Delta_p$  is the  $p$ -Laplacian operator, and  $f \in C([r_0, +\infty) \times [0, +\infty), \mathbb{R})$  is a non-decreasing function with respect to its second variable. By using the Mountain Pass Theorem, they prove the existence of positive radial solutions for small values of  $\lambda$ .

**Keywords:** Kirchhoff problem,  $p$ -Laplacian, positive radial solution, variational methods.

**Mathematics Subject Classification:** 35A01, 35A15, 35B38, 35D30, 35J92.

### 1. INTRODUCTION

The aim of this work is to prove the existence of positive radial solutions on the exterior of a ball to the Kirchhoff type problem

$$\begin{cases} -\left(a + b \int_{\Omega_e} |\nabla u|^p dx\right) \Delta_p u = \lambda f(|x|, u), & x \in \Omega_e, \\ u(x) = 0, & |x| = r_0, \\ u(x) \rightarrow 0, & |x| \rightarrow \infty, \end{cases} \quad (1.1)$$

where  $a$  and  $b$  are positive constants,  $\lambda > 0$  is a parameter,  $\Omega_e = \{x \in \mathbb{R}^N : |x| > r_0\}$ ,  $r_0 > 0$ ,  $N > p > 1$ ,  $\Delta_p$  is the  $p$ -Laplacian operator ( $\Delta_p u = \operatorname{div}(|\nabla u|^{p-2} \nabla u)$ ), and  $f : [r_0, +\infty) \times [0, +\infty) \rightarrow \mathbb{R}$  is continuous and is non-decreasing in its second variable.

Note that Kirchhoff type problems are nonlinear, and as such present several interesting challenges; see, for instance, the recent work in [1, 2, 12–15, 18, 21, 24] for various issues and applications. Additional work on  $p$ -Laplacian problems can be found in [6–10, 17, 20, 23] and other related results in [16, 22].

In [7, 8, 17, 23], the equations considered are of the form

$$\Delta_p u = \lambda f(u) \quad \text{in } \Omega$$

with Dirichlet boundary conditions, where  $\Omega$  is a bounded domain in  $\mathbb{R}^N$ . Concerning the existence of positive radial solutions to a class of  $p$ -Laplacian problems on the exterior of a ball, we mention the papers [9] for  $p = 2$  and [20] for any  $p > 1$ . In these articles the equation is of the form  $-\Delta_p u = \lambda K(|x|)f(u)$  and the authors appeal to the Mountain Pass Theorem (MPT).

Notice that our problem (1.1) can be written as

$$\begin{cases} -M\left(\int_{\Omega_e} |\nabla u|^p dx\right) \Delta_p u = \lambda f(|x|, u), & x \in \Omega_e, \\ u(x) = 0, & |x| = r_0, \\ u(x) \rightarrow 0, & |x| \rightarrow \infty, \end{cases}$$

where  $M(\zeta) = a + b\zeta$ .

In the case where  $p = 2$  and  $M$  is any positive function defined on  $\mathbb{R}^+$  (with some additional conditions), problems of the type

$$\begin{cases} -M\left(\int_{\Omega} |\nabla u|^2 dx\right) \Delta u = f(x, u), & \text{in } \Omega, \\ u = 0, & \text{on } \partial\Omega, \end{cases}$$

where  $\Omega$  a bounded domain in  $\mathbb{R}^N$ , have physical motivations. For example, the Kirchhoff operator  $M(\int_{\Omega} |\nabla u|^2 dx) \Delta u$  appears in nonlinear vibration equations such as

$$\begin{cases} u_{tt} - M\left(\int_{\Omega} |\nabla u|^2 dx\right) \Delta u = f(x, u), & \text{in } \Omega \times (0, T), \\ u = 0, & \text{on } \partial\Omega \times (0, T), \\ u(x, 0) = u_0(x), \quad u_t(x, 0) = u_1(x). \end{cases}$$

Such equations generalize to higher dimensions the equation studied by Kirchhoff [19],

$$\rho \frac{\partial^2 u}{\partial t^2} - \left( \frac{P_0}{h} + \frac{E}{2L} \int_0^L \left| \frac{\partial u}{\partial x} \right|^2 dx \right) \frac{\partial^2 u}{\partial x^2} = 0$$

as an extension of the classical D'Alembert wave equation for free vibrations of elastic strings.

Kirchhoff type problems have been treated in many papers. For example, in [2], by using truncations and the MPT, the authors proved the existence of solutions to the problem

$$\begin{cases} -M\left(\int_{\Omega} |\nabla u|^2 dx\right) \Delta u = f(x, u), & \text{in } \Omega, \\ u = 0, & \text{on } \partial\Omega, \end{cases}$$

where  $\Omega$  is a bounded smooth domain in  $\mathbb{R}^N$ . In [18], He *et al.* considered a similar problem where  $\Omega$  is a bounded domain in  $\mathbb{R}^3$  or is all of  $\mathbb{R}^3$ , and instead of  $f(t, u)$ , they had  $f(u) + h$  with  $h \geq 0$  and  $h \in L^2(\Omega)$ . In [24], Wang *et al.* also took  $\Omega$  to also be a bounded and smooth domain in  $\mathbb{R}^N$  and used the MPT to prove the existence of solutions to the problem

$$\begin{cases} -M\left(\int_{\Omega} |\nabla u|^p dx\right) \Delta_p u = \lambda f(x, u) + |u|^{p^*-2}u, & \text{in } \Omega, \\ u = 0, & \text{on } \partial\Omega, \end{cases}$$

for all  $\lambda$  greater than some  $\lambda^* > 0$ , where  $p^* = \frac{Np}{N-p}$ . An important feature of that study is that  $M$  could be zero at zero. Additional recent results on Kirchhoff type problems can be found in [1, 12–15, 21].

Extending the ideas in [9, 20], instead of  $\Delta_p u$ , we consider a Kirchhoff type operator and generalize the term  $K(|x|)f(u)$  to  $f(|x|, u)$ , where  $f : [r_0, +\infty) \times [0, +\infty) \rightarrow \mathbb{R}$  is continuous, non-decreasing in its second variable, and satisfies:

(F1) there exist continuous functions  $A, B : [r_0, +\infty) \rightarrow (0, +\infty)$  with  $q > 2p - 1$  and  $\mu \in \left(0, \frac{N-p}{p-1}\right)$  such that

$$A(\xi)(t^q - 1) \leq f(\xi, t) \leq B(\xi)(t^q + 1) \quad \text{for all } (\xi, t) \in [r_0, +\infty) \times [0, +\infty)$$

where  $A(\xi), B(\xi) \leq \frac{1}{\xi^{N+\mu}}$  for  $\xi \gg 1$ ,

(F2) for all  $\xi \in [r_0, +\infty)$ ,  $f(\xi, 0) < 0$ ,

(F3) (Ambrosetti-Rabinowitz condition) There exists  $\theta > 2p$  such that, for all sufficiently large  $t$ ,

$$tf(\xi, t) > \theta F(\xi, t) \quad \text{for all } \xi \geq r_0,$$

where  $F(\xi, t) = \int_0^t f(\xi, \sigma) d\sigma$ .

Applying the change of variables  $r = |x|$  and  $s = \left(\frac{r}{r_0}\right)^{\frac{p-N}{p-1}}$  transforms (1.1) into the boundary-value problem (see, for example, [6])

$$\begin{cases} -\left(a + \alpha \int_0^1 |u'|^p d\sigma\right) (\phi_p(u'))' = \lambda h(s) f(r_0 s^{\frac{p-1}{p-N}}, u(s)), & s \in (0, 1), \\ u(0) = u(1) = 0, \end{cases} \quad (1.2)$$

where

$$\alpha = bN\omega_N r_0^{N-p} \left( \frac{N-p}{p-1} \right)^{p-1}, \quad \phi_p(\zeta) = |\zeta|^{p-2}\zeta, \quad h(s) = \left( r_0 \frac{(p-1)}{(N-p)} \right)^p s^{-\frac{p(N-1)}{N-p}},$$

and  $\omega_N$  is the volume of the unit ball in  $\mathbb{R}^N$ .

**Remark 1.1.** If in (F1) we assume that  $\mu \geq \frac{N-p}{p-1}$ , this would imply that the functions defined by  $h(s)B(r_0 s^{\frac{p-1}{p-N}})$  or  $h(s)A(r_0 s^{\frac{p-1}{p-N}})$  are dominated in neighborhoods of zero by a continuous function on  $[0, 1]$ , and in fact, we would have a simpler situation. But  $\mu \in \left(0, \frac{N-p}{p-1}\right)$  implies the singularity of these functions at  $s = 0$ , but they would still belong to  $L^1(0, 1)$ .

**Remark 1.2.** Consider the function

$$f(\xi, t) = \frac{1}{2\xi^{\frac{7}{2}}}(2t^4 - 1).$$

Then  $f$  satisfies all of the above conditions for  $N = 3$ ,  $p = 2$ ,  $q = 4$ ,  $\mu = \frac{1}{2}$ , and  $\theta = \frac{9}{2}$ .

As a second example, we have the following.

**Remark 1.3.** Take  $N = 4$ ,  $p = 2$ , and  $q = 4 > 2p - 1 = 3$ . We need  $\mu \in (0, 2)$  so choose  $\mu = 1$ . Let

$$A(\xi) = \frac{2 + \sin \xi}{\xi^5} \quad \text{and} \quad B(\xi) = \frac{(4 - \cos \xi)(\xi^2 + 1)}{\xi^7}.$$

Then

$$f(\xi, t) = \frac{3(\xi^2 + 1)(e^{t^2} - 2)(t^4 + 1)}{\xi^7 e^{t^2}}$$

satisfies all of the above conditions for some  $\theta > 4$ .

Next, we define what is meant by a solution of our problem.

**Definition 1.4.** We say that  $u \in W_0^{1,p}(0, 1)$  is a weak solution of problem (1.2) if

$$(a + \alpha \|u\|_{1,p}^p) \int_0^1 |u'(s)|^{p-2} u'(s) v'(s) ds = \lambda \int_0^1 h(s) f(r_0 s^{\frac{p-1}{p-N}}, u(s)) v(s) ds$$

for all  $v \in W_0^{1,p}(0, 1)$ .

We will establish the following theorem, which is our main result in this paper.

**Theorem 1.5.** *Assume that (F1)–(F3) hold. Then (1.2) admits a positive weak solution for  $\lambda \approx 0$ .*

## 2. PRELIMINARIES

In order to apply variational techniques such as the MPT, we extend the function  $f$  to  $[r_0, +\infty) \times \mathbb{R}$  by setting  $f(\xi, t) = f(\xi, 0)$  for  $(\xi, t) \in [r_0, +\infty) \times (-\infty, 0)$ . We also need the Banach spaces  $W_0^{1,p}(0, 1)$ ,  $C[0, 1]$ , and  $L^r(0, 1)$  equipped their respective norms  $\|\cdot\|_{1,p}$ ,  $\|\cdot\|_\infty$ , and  $\|\cdot\|_r$ . We recall that  $W_0^{1,p}(0, 1)$  is compactly embedded in  $C[0, 1]$ , and this implies that  $\|u\|_\infty \leq k\|u\|_{1,p}$  for every  $u$  in  $W_0^{1,p}(0, 1)$ , where  $k$  is a fixed positive constant (see [5]).

**Remark 2.1.** Let

$$D = \{(\xi, t) \in [r_0, +\infty) \times \mathbb{R} : f(\xi, t) \geq 0\}$$

and

$$D^c = \{(\xi, t) \in [r_0, +\infty) \times \mathbb{R} : f(\xi, t) < 0\}.$$

On  $D$ , we have

$$|f(\xi, t)| = f(\xi, t) \leq B(\xi)(t^q + 1),$$

and on  $D^c$ ,

$$|f(\xi, t)| \leq A(\xi).$$

Hence, for all  $(\xi, t) \in [r_0, +\infty) \times \mathbb{R}$ ,

$$|f(\xi, t)| \leq \max(A(\xi), B(\xi))(|t|^q + 1),$$

and for every compact interval  $I \subset \mathbb{R}$ , there exists a constant  $M_I$  such that

$$|f(\xi, t)| \leq M_I \max(A(\xi), B(\xi)) \quad \text{for all } \xi \geq r_0 \text{ and all } t \in I.$$

**Remark 2.2.** If  $f$  satisfies (F1) and (F3), then:

(F4) There exists a continuous function  $\theta_1 : [r_0, +\infty) \rightarrow (0, +\infty)$  and a constant  $C > 0$  such that

$$\theta_1(\xi) \leq \frac{C}{\xi^{N+\mu}}$$

and

$$tf(\xi, t) > \theta F(\xi, t) - \theta_1(\xi) \quad \text{for all } (\xi, t) \in [r_0, +\infty) \times [0, +\infty).$$

**Remark 2.3.** We note that (F1) implies that there exist continuous functions  $A_1, B_1 : [r_0, +\infty) \rightarrow (0, +\infty)$  and a positive constant  $C_1$  such that

$$F(\xi, t) \leq B_1(\xi)(|t|^{q+1} + 1) \quad \text{for all } (\xi, t) \in [r_0, +\infty) \times \mathbb{R},$$

and

$$F(\xi, t) \geq A_1(\xi)(t^{q+1} - C_1) \quad \text{for all } (\xi, t) \in [r_0, +\infty) \times [0, +\infty).$$

Furthermore,  $A_1(\xi), B_1(\xi) \leq \frac{1}{\xi^{N+\mu}}$  for  $\xi \gg 1$ , where  $\mu$  is given in (F1). Notice that the second inequality above follows from the fact that  $\frac{t^{q+1}}{q+1} - t \geq \frac{t^{q+1}}{2(q+1)}$  for all  $t \geq (2(q+1))^{\frac{1}{q}}$ .

**Lemma 2.4.** Let  $J : W_0^{1,p}(0,1) \rightarrow \mathbb{R}$  be defined by

$$J(u) = \frac{1}{p} \hat{M}(\|u\|_{1,p}^p) - \lambda K(u)$$

where

$$K(u) = \int_0^1 h(s) F(r_0 s^{\frac{p-1}{p-N}}, u(s)) ds$$

and

$$\hat{M}(t) = \int_0^t M(\sigma) d\sigma \quad \text{with} \quad M(t) = a + \alpha t.$$

Then  $J$  is well defined, continuously differentiable, and for all  $v \in W_0^{1,p}(0,1)$ , its Gâteaux derivative is given by

$$J'(u)(v) = M(\|u\|_{1,p}^p) \int_0^1 |u'(s)|^{p-2} u'(s) v'(s) ds - \lambda \int_0^1 h(s) f(r_0 s^{\frac{p-1}{p-N}}, u(s)) v(s) ds.$$

*Proof.* It is clear that  $\hat{M}(\|u\|_{1,p}^p)$  is finite, and since  $W_0^{1,p}(0,1) \hookrightarrow C[0,1]$ , by Remark 2.1, for  $I = [-\|u\|_\infty, \|u\|_\infty]$ , there exists  $M_I > 0$  such that

$$|f(r_0 s^{\frac{p-1}{p-N}}, \sigma)| \leq M_I \max(A(r_0 s^{\frac{p-1}{p-N}}), B(r_0 s^{\frac{p-1}{p-N}})) \quad \text{for all } \sigma \in I.$$

Therefore,

$$\int_0^1 |h(s)| |F(r_0 s^{\frac{p-1}{p-N}}, u(s))| ds \leq M_I \|u\|_\infty \int_0^1 \max(A(r_0 s^{\frac{p-1}{p-N}}), B(r_0 s^{\frac{p-1}{p-N}})) |h(s)| ds < \infty.$$

The functional  $J$  is continuous. Moreover, if we set  $L(u) = \|u\|_{1,p}^p$ , then  $\hat{M} \circ L$  is differentiable, and for all  $v \in W_0^{1,p}(0,1)$ ,

$$\frac{1}{p} (\hat{M} \circ L)'(u)(v) = M(\|u\|_{1,p}^p) \int_0^1 |u'(s)|^{p-2} u'(s) v'(s) ds.$$

On the other hand, from the continuity of  $f$ , we see that  $K$  is Gâteaux differentiable and its Gâteaux derivative is continuous. Hence,  $K$  is continuously differentiable and

$$K'(u)(v) = \int_0^1 h(s) f(r_0 s^{\frac{p-1}{p-N}}, u(s)) v(s) ds \quad \text{for all } v \in W_0^{1,p}(0,1).$$

Therefore,  $J$  is continuously differentiable, and for all  $v \in W_0^{1,p}(0, 1)$ ,

$$J'(u)(v) = M(\|u\|_{1,p}^p) \int_0^1 |u'(s)|^{p-2} u'(s) v'(s) - \lambda \int_0^1 h(s) f(r_0 s^{\frac{p-1}{p-N}}, u(s)) v(s) ds$$

as we wished to show.  $\square$

In order to prove our main result, Theorem 1.5 above, we will apply the Mountain Pass Theorem stated below.

**Theorem 2.5** (Mountain Pass Theorem [3]). *Let  $X$  be a Banach space and let  $J \in C^1(X; \mathbb{R})$  satisfy:*

- (i) (Palais–Smale condition) *any sequence  $(u_n) \subset X$  such that  $(J(u_n))$  is bounded and  $J'(u_n) \rightarrow 0$  as  $n \rightarrow \infty$  possesses a convergent subsequence,*
- (ii)  $J(0) = 0$ ,
- (iii) *there exist  $\nu, R > 0$  such that  $J(u) \geq \nu$  for all  $u$  with  $\|u\|_X = R$ ,*
- (iv) *there exists  $e \in X$  such that  $\|e\|_X > R$  and  $J(e) < 0$ .*

In addition, let

$$\Gamma := \{\gamma \in C([0, 1], X) : \gamma(0) = 0, \gamma(1) = e\}$$

and

$$\hat{c} := \inf_{\gamma \in \Gamma} \max_{t \in [0, 1]} J(\gamma(t)).$$

Then  $\hat{c}$  is a critical value of the functional  $J$ .

### 3. PROOF OF THEOREM 1.5

In this section, we construct the proof of our main result. We begin by recalling that proving Theorem 1.5 is equivalent to proving that the functional  $J$  defined above admits a positive critical point for  $\lambda \approx 0$  (see[4]). As was seen in Lemma 2.4, the functional  $J$  is in  $C^1(W_0^{1,p}(0, 1), \mathbb{R})$ , so we need to prove that  $J$  satisfies the conditions of the MPT.

In the following, for all  $s \in (0, 1]$ , we denote by  $\tilde{A}(s)$ ,  $\tilde{B}(s)$ ,  $\tilde{A}_1(s)$ ,  $\tilde{B}_1(s)$ , and  $\tilde{\theta}_1(s)$  the quantities  $A(r_0 s^{\frac{p-1}{p-N}})$ ,  $B(r_0 s^{\frac{p-1}{p-N}})$ ,  $A_1(r_0 s^{\frac{p-1}{p-N}})$ ,  $B_1(r_0 s^{\frac{p-1}{p-N}})$ , and  $\theta_1(r_0 s^{\frac{p-1}{p-N}})$ , respectively.

#### 3.1. THE PALAIS–SMALE CONDITION

In order to show that our functional  $J$  satisfies the Palais–Smale condition, we first recall the following proposition.

**Proposition 3.1** ([11]). *Let  $\psi : W^{1,p}(0, 1) \rightarrow [0, +\infty)$  be defined by*

$$\psi(u) = \frac{1}{p} \int_0^1 |u'(s)|^p ds.$$

Then  $\psi'$  exists and

$$\langle \psi'(u), v \rangle = \int_0^1 |u'(s)|^{p-2} u'(s) v'(s) ds.$$

In addition, if  $u_n \rightharpoonup u$  and  $\limsup_{n \rightarrow +\infty} \langle \psi'(u_n), u_n - u \rangle \leq 0$ , then  $u_n \rightarrow u$  strongly in  $W^{1,p}(0,1)$ .

**Lemma 3.2.** *The functional  $J$  satisfies the Palais-Smale condition.*

*Proof.* Let  $(u_n)_n \subset W_0^{1,p}(0,1)$  such that  $(J(u_n))_n$  is bounded and  $J'(u_n) \rightarrow 0$  as  $n \rightarrow \infty$ . First, we will prove that  $(u_n)_n$  is bounded in  $W_0^{1,p}(0,1)$ . Assume to contrary that  $(u_n)_n$  is such that  $J'(u_n) \rightarrow 0$  as  $n \rightarrow \infty$ , there exists  $M > 0$  such that  $|J(u_n)| \leq M$  for all  $n \geq 1$ , but  $\|u_n\|_{1,p} \rightarrow \infty$  as  $n \rightarrow \infty$ .

We consider the quantity

$$\frac{\theta J(u_n) - \langle J'(u_n), u_n \rangle}{\|u_n\|_{1,p}},$$

where  $\theta > 2p$  is chosen as in (F3). Since  $J(u_n)$  is bounded and  $J'(u_n) \rightarrow 0$  as  $n \rightarrow \infty$ ,

$$\lim_{n \rightarrow \infty} \frac{\theta J(u_n) - \langle J'(u_n), u_n \rangle}{\|u_n\|_{1,p}} = 0.$$

However, we have

$$\begin{aligned} \theta J(u_n) - \langle J'(u_n), u_n \rangle &= a \left( \frac{\theta}{p} - 1 \right) \|u_n\|_{1,p}^p + \alpha \left( \frac{\theta}{2p} - 1 \right) \|u_n\|_{1,p}^{2p} \\ &\quad - \lambda \int_0^1 h(s) \left( \theta F(r_0 s^{\frac{p-1}{p-N}}, u_n(s)) - f(r_0 s^{\frac{p-1}{p-N}}, u_n(s)) u_n(s) \right) ds \\ &= a \left( \frac{\theta}{p} - 1 \right) \|u_n\|_{1,p}^p + \alpha \left( \frac{\theta}{2p} - 1 \right) \|u_n\|_{1,p}^{2p} - \lambda (I_1 + I_2), \end{aligned}$$

where

$$I_1 = \int_{\{u_n \geq 0\}} h(s) \left( \theta F(r_0 s^{\frac{p-1}{p-N}}, u_n(s)) - f(r_0 s^{\frac{p-1}{p-N}}, u_n(s)) u_n(s) \right) ds$$

and

$$I_2 = \int_{\{u_n < 0\}} h(s) \left( \theta F(r_0 s^{\frac{p-1}{p-N}}, u_n(s)) - f(r_0 s^{\frac{p-1}{p-N}}, u_n(s)) u_n(s) \right) ds,$$

with

$$\{u_n \geq 0\} = \{s \in [0,1] : u_n(s) \geq 0\}$$

and

$$\{u_n < 0\} = \{s \in [0,1] : u_n(s) < 0\}.$$



Using (F4), we can write

$$I_1 \leq \int_{\{u_n \geq 0\}} h(s)\theta_1(r_0 s^{\frac{p-1}{p-N}}) ds \leq \int_{[0,1]} h(s)\theta_1(r_0 s^{\frac{p-1}{p-N}}) ds \leq \|h\tilde{\theta}_1\|_1,$$

and on  $\{u_n < 0\}$ ,

$$\begin{aligned} \theta F(r_0 s^{\frac{p-1}{p-N}}, u_n(s)) - f(r_0 s^{\frac{p-1}{p-N}}, u_n(s))u_n(s) &= (\theta - 1)f(r_0 s^{\frac{p-1}{p-N}}, 0)u_n(s) \\ &\leq (\theta - 1)A(r_0 s^{\frac{p-1}{p-N}})\|u_n\|_\infty \\ &\leq k(\theta - 1)A(r_0 s^{\frac{p-1}{p-N}})\|u_n\|_{1,p}, \end{aligned}$$

so that

$$\begin{aligned} \frac{\theta J(u_n) - \langle J'(u_n), u_n \rangle}{\|u_n\|_{1,p}} &\geq a\left(\frac{\theta}{p} - 1\right)\|u_n\|_{1,p}^{p-1} + \alpha\left(\frac{\theta}{2p} - 1\right)\|u_n\|_{1,p}^{2p-1} \\ &\quad - \lambda \frac{\|h\tilde{\theta}_1\|_{L^1}}{\|u_n\|_{1,p}} - k\lambda(\theta - 1)\|h\tilde{A}\|_1. \end{aligned}$$

Taking the limit as  $n \rightarrow +\infty$ , we obtain a contradiction. Thus,  $(u_n)$  is bounded in  $W_0^{1,p}(0,1)$  and this implies that there exists a subsequence, again calling it  $(u_n)$ , that converges weakly in  $W_0^{1,p}(0,1)$  and strongly in  $C[0,1]$ .

We want to show that  $u_n \rightarrow u$  strongly in  $W_0^{1,p}(0,1)$ . Since for all  $v \in W_0^{1,p}(0,1)$ ,

$$J'(u_n)(v) = M(\|u_n\|_{1,p}^p) \int_0^1 |u'_n(s)|^{p-2} u'_n(s) v'(s) ds - \lambda \int_0^1 h(s) f(r_0 s^{\frac{p-1}{p-N}}, u_n(s)) v(s) ds,$$

we have

$$\left| \int_0^1 |u'_n|^{p-2} u'_n (u'_n - u') \right| \leq \frac{|J'(u_n)(u_n - u)| + \lambda \left| \int_0^1 h(s) f(r_0 s^{\frac{p-1}{p-N}}, u_n(s)) (u_n - u) ds \right|}{a}.$$

Since  $J'(u_n) \rightarrow 0$  and  $(u_n)$  is bounded in  $W_0^{1,p}(0,1)$ , we have  $J'(u_n)(u_n - u) \rightarrow 0$  as  $n \rightarrow +\infty$ . On the other hand, since  $u_n \rightarrow u$  strongly in  $C[0,1]$  and since  $(u_n)$  is bounded in the same space, we have (see Remark 2.1)

$$\left| \int_0^1 h(s) f(r_0 s^{\frac{p-1}{p-N}}, u_n(s)) (u_n(s) - u(s)) ds \right| \leq M_I \|u_n - u\|_\infty \|h \max(\tilde{A}, \tilde{B})\|_1 \rightarrow 0, \quad (3.1)$$

where  $I = [-M, M]$  is such that  $u, u_n \in [-M, M]$  for  $s \in [0,1]$ . This implies that

$$\left| \int_0^1 |u'_n(s)|^{p-2} u'_n(s) (u'_n(s) - u'(s)) ds \right| \rightarrow 0$$

as  $n \rightarrow \infty$ . By Proposition 3.1,  $u_n \rightarrow u$  strongly in  $W_0^{1,p}(0,1)$ , which completes the proof of the lemma.  $\square$

3.2. THE GEOMETRY OF  $J$ 

We begin by pointing out that  $J(0) = 0$ . Next, we prove two lemmas that will be needed to complete the proof that  $J$  satisfies the Mountain Pass Theorem.

**Lemma 3.3.** *For any positive function  $w$  in  $W_0^{1,p}(0,1)$  satisfying  $\|w\|_{1,p} = 1$ , we have  $\lim_{\sigma \rightarrow +\infty} J(\sigma w) = -\infty$  for any  $\sigma > 0$ .*

*Proof.* Let  $w \in W_0^{1,p}(0,1)$  be such that  $w$  is positive with  $\|w\|_{1,p} = 1$ , and let  $\sigma > 0$  be a parameter. We have

$$J(\sigma w) = \frac{1}{p} \hat{M}(\|\sigma w\|_{1,p}^p) - \lambda \int_0^1 h(s) F(r_0 s^{\frac{p-1}{p-N}}, \sigma w(s)) ds$$

with

$$\hat{M}(\|\sigma w\|_{1,p}^p) = a\sigma^p + \frac{\alpha}{2}\sigma^{2p}.$$

By (F1) and an integration,

$$\int_0^1 h(s) F(r_0 s^{\frac{p-1}{p-N}}, \sigma w(s)) ds \geq \int_0^1 h(s) A(r_0 s^{\frac{p-1}{p-N}}) \left( \frac{(\sigma w(s))^{q+1}}{q+1} - \sigma w(s) \right) ds. \quad (3.2)$$

Then,

$$\begin{aligned} J(\sigma w) &\leq \frac{a}{p}\sigma^p + \frac{\alpha}{2p}\sigma^{2p} - \lambda \frac{\sigma^{q+1}}{q+1} \int_0^1 h(s) A(r_0 s^{\frac{p-1}{p-N}}) (w(s))^{q+1} ds \\ &\quad + \lambda \sigma \int_0^1 h(s) A(r_0 s^{\frac{p-1}{p-N}}) w(s) ds. \end{aligned}$$

But

$$0 < \int_0^1 h(s) A(r_0 s^{\frac{p-1}{p-N}}) (w(s))^{q+1} ds \leq k^{q+1} \|h\tilde{A}\|_1 \|w\|_{1,p}^{q+1} < \infty$$

and

$$0 < \int_0^1 h(s) A(r_0 s^{\frac{p-1}{p-N}}) w(s) ds \leq k \|h\tilde{A}\|_1 \|w\|_{1,p} < \infty,$$

so  $\lim_{\sigma \rightarrow +\infty} J(\sigma w) = -\infty$  since  $q > 2p - 1$ . This proves the lemma.  $\square$

**Lemma 3.4.** *There exists  $\lambda_0 > 0$  such that for all  $\lambda \in (0, \lambda_0)$  and  $u \in W_0^{1,p}(0, 1)$  be such that  $\|u\|_{1,p} = \lambda^{\frac{-1}{2(q+1-2p)}}$ , we have  $J(u) \geq \frac{\alpha}{4p} \lambda^{\frac{-p}{q+1-2p}}$ .*

*Proof.* Let  $\lambda > 0$  and  $u \in W_0^{1,p}(0, 1)$  be such that

$$\|u\|_{1,p} = \lambda^{\frac{-1}{2(q+1-2p)}}.$$

We have

$$\begin{aligned} J(u) &\geq \frac{a}{p} \lambda^{\frac{-p}{2(q+1-2p)}} + \frac{\alpha}{2p} \lambda^{\frac{-p}{q+1-2p}} - \lambda \int_0^1 h(s) B_1(r_0 s^{\frac{p-1}{p-N}}) (|u(s)|^{q+1} + 1) ds \\ &\geq \frac{a}{p} \lambda^{\frac{-p}{2(q+1-2p)}} + \frac{\alpha}{2p} \lambda^{\frac{-p}{q+1-2p}} - \lambda \|u\|_{\infty}^{q+1} \|h\tilde{B}_1\|_1 - \lambda \|h\tilde{B}_1\|_1 \\ &\geq \frac{a}{p} \lambda^{\frac{-p}{2(q+1-2p)}} + \frac{\alpha}{2p} \lambda^{\frac{-p}{q+1-2p}} - k \lambda \lambda^{\frac{-(q+1)}{2(q+1-2p)}} - \lambda \|h\tilde{B}_1\|_1 \\ &\geq \lambda^{\frac{-p}{q+1-2p}} \left( \frac{\alpha}{2p} + \frac{a}{p} \lambda^{\frac{p}{2(q+1-2p)}} - k \|h\tilde{B}_1\|_1 \lambda^{\frac{1}{2}} - \|h\tilde{B}_1\|_1 \lambda^{\frac{q+1-p}{q+1-2p}} \right). \end{aligned}$$

Since

$$\lim_{\lambda \rightarrow 0} \frac{a}{p} \lambda^{\frac{p}{2(q+1-2p)}} - k \|h\tilde{B}_1\|_1 \lambda^{\frac{1}{2}} - \|h\tilde{B}_1\|_1 \lambda^{\frac{q+1-p}{q+1-2p}} = 0,$$

there exists  $\lambda_0 > 0$  such that for all  $\lambda \in (0, \lambda_0)$ ,

$$J(u) \geq \frac{\alpha}{4p} \lambda^{\frac{-p}{q+1-2p}},$$

which proves the lemma.  $\square$

From Lemma 3.3 and Lemma 3.4, we can deduce that conditions (iii) and (iv) of Theorem 2.5 are satisfied. We have then proved that the functional  $J$  admits a critical value for  $\lambda \approx 0$ . We need to show that this critical point is positive.

### 3.3. POSITIVITY OF THE MPT SOLUTION

Let  $r = \frac{1}{q+1-2p}$ . We start with two lemmas.

**Lemma 3.5.** *Let  $u_\lambda$  be a mountain pass solution to (1.1). Then there exists  $M_0 > 0$  and  $\lambda_1 > 0$  such that*

$$\|u_\lambda\|_{\infty} \geq M_0 \lambda^{-r \frac{p}{q+1}}$$

for all  $\lambda \in (0, \lambda_1)$ .

*Proof.* Since  $u_\lambda$  is a mountain pass solution, we have

$$\begin{aligned}
\lambda \int_0^1 h(s) f(r_0 s^{\frac{p-1}{p-N}}, u_\lambda(s)) u_\lambda(s) ds &= a \|u_\lambda\|_{1,p}^p + \alpha \|u_\lambda\|_{1,p}^{2p} \\
&= pJ(u_\lambda) + \frac{\alpha}{2} \|u_\lambda\|_{1,p}^{2p} + p\lambda \int_0^1 h(s) F(r_0 s^{\frac{p-1}{p-N}}, u_\lambda) ds \\
&\geq pJ(u_\lambda) + p\lambda \int_{\{u_\lambda < 0\}} h(s) F(r_0 s^{\frac{p-1}{p-N}}, u_\lambda(s)) ds \\
&\quad + p\lambda \int_{\{u_\lambda \geq 0\}} h(s) F(r_0 s^{\frac{p-1}{p-N}}, u_\lambda(s)) ds \\
&\geq pJ(u_\lambda) + p\lambda \int_{\{u_\lambda \geq 0\}} h(s) F(r_0 s^{\frac{p-1}{p-N}}, u_\lambda(s)) ds.
\end{aligned}$$

In view of Remark 2.3 and the fact that  $u_\lambda$  satisfies

$$J(u_\lambda) \geq \frac{\alpha}{4p} \lambda^{-rp} \quad \text{for all } \lambda \in (0, \lambda_0),$$

we see that

$$\begin{aligned}
\lambda \int_0^1 h(s) f(r_0 s^{\frac{p-1}{p-N}}, u_\lambda) u_\lambda ds &\geq \frac{\alpha}{4} \lambda^{-rp} + p\lambda \int_{\{u_\lambda \geq 0\}} h(s) A_1(r_0 s^{\frac{p-1}{p-N}}) u_\lambda^{q+1} ds \\
&\quad - C_1 p\lambda \int_{\{u_\lambda \geq 0\}} h(s) A_1(r_0 s^{\frac{p-1}{p-N}}) ds \\
&\geq \frac{\alpha}{4} \lambda^{-rp} - C_1 p\lambda \|h\tilde{A}_1\|_1 \geq \frac{\alpha}{8} \lambda^{-rp},
\end{aligned}$$

for all  $\lambda \in \left(0, \min\left(\lambda_0, \left(\frac{\alpha}{8p\|h\tilde{A}_1\|_1}\right)^{\frac{1}{rp+1}}\right)\right)$ . By Remark 2.1, for all

$$\lambda \in \left(0, \min\left(\lambda_0, \left(\frac{\alpha}{8p\|h\tilde{A}_1\|_1}\right)^{\frac{1}{rp+1}}, 1\right)\right),$$

we have

$$\|u_\lambda\|_\infty^{q+1} + \|u_\lambda\|_\infty \geq \frac{\alpha}{8\|h \max(\tilde{A}, \tilde{B})\|_1} \lambda^{-rp}.$$

Then, for all

$$\lambda \in \left(0, \min \left( \lambda_0, \left( \frac{\alpha}{8p\|h\tilde{A}_1\|_1} \right)^{\frac{1}{rp+1}}, 1, \left( \frac{\alpha}{16\|h \max(\tilde{A}, \tilde{B})\|_1} \right)^{\frac{1}{rp}} \right) \right),$$

we have  $\|u_\lambda\|_\infty \geq 1$ , so

$$\|u_\lambda\|_\infty^{q+1} \geq \frac{\alpha}{16\|h \max(\tilde{A}, \tilde{B})\|_1} \lambda^{-rp},$$

or

$$\|u_\lambda\|_\infty \geq \left( \frac{\alpha}{16\|h \max(\tilde{A}, \tilde{B})\|_1} \right)^{\frac{1}{q+1}} \lambda^{-r \frac{p}{q+1}}.$$

Taking

$$M_0 = \left( \frac{\alpha}{16\|h \max(\tilde{A}, \tilde{B})\|_1} \right)^{\frac{1}{q+1}}$$

and

$$\lambda_1 = \min \left( \lambda_0, \left( \frac{\alpha}{8p\|h\tilde{A}_1\|_1} \right)^{\frac{1}{rp+1}}, 1, \left( \frac{\alpha}{16\|h \max(\tilde{A}, \tilde{B})\|_1} \right)^{\frac{1}{rp}} \right)$$

completes the proof of the lemma.  $\square$

**Lemma 3.6.** *Let  $u_\lambda$  be a mountain pass solution of (1.2). Then there exists  $C_0 > 0$  and  $\lambda_2 > 0$  such that*

$$\|u_\lambda\|_{1,p} \leq C_0 \lambda^{-r},$$

for all  $\lambda \in (0, \lambda_2)$ .

*Proof.* Since  $u_\lambda$  is a solution of (1.2), by Remark 2.2, we have

$$\begin{aligned}
a\|u_\lambda\|_{1,p}^p + \frac{\alpha}{2}\|u_\lambda\|_{1,p}^{2p} &= pJ(u_\lambda) + p\lambda \int_0^1 h(s)F(r_0s^{\frac{p-1}{p-N}}, u_\lambda(s))ds \\
&= pJ(u_\lambda) + p\lambda \int_{\{u_\lambda < 0\}} h(s)u_\lambda(s)f(r_0s^{\frac{p-1}{p-N}}, 0)ds \\
&\quad + p\lambda \int_{\{u_\lambda \geq 0\}} h(s)F(r_0s^{\frac{p-1}{p-N}}, u_\lambda(s))ds \\
&\leq pJ(u_\lambda) + p\lambda \int_{\{u_\lambda < 0\}} h(s)u_\lambda(s)f(r_0s^{\frac{p-1}{p-N}}, 0)ds \\
&\quad + p\lambda \int_{\{u_\lambda \geq 0\}} \frac{h(s)}{\theta} \left( u_\lambda(s)f(r_0s^{\frac{p-1}{p-N}}, u_\lambda(s)) + \theta_1(r_0s^{\frac{p-1}{p-N}}) \right) ds \\
&\leq pJ(u_\lambda) + p\lambda \int_{\{u_\lambda < 0\}} h(s)u_\lambda(s)f(r_0s^{\frac{p-1}{p-N}}, 0)ds \\
&\quad + \frac{p}{\theta}\lambda \int_0^1 h(s) \left( u_\lambda(s)f(r_0s^{\frac{p-1}{p-N}}, u_\lambda(s)) + \theta_1(r_0s^{\frac{p-1}{p-N}}) \right) ds \\
&\quad - \frac{p}{\theta}\lambda \int_{\{u_\lambda < 0\}} h(s) \left( u_\lambda(s)f(r_0s^{\frac{p-1}{p-N}}, 0) + \theta_1(r_0s^{\frac{p-1}{p-N}}) \right) ds \\
&\leq pJ(u_\lambda) + \frac{p}{\theta}\lambda \int_0^1 h(s)u_\lambda(s)f(r_0s^{\frac{p-1}{p-N}}, u_\lambda(s))ds \\
&\quad + \frac{p}{\theta}\lambda \|h\tilde{\theta}_1\|_1 + p\lambda \left(1 - \frac{1}{\theta}\right) \int_{\{u_\lambda < 0\}} h(s)u_\lambda(s)f(r_0s^{\frac{p-1}{p-N}}, 0)ds \\
&\leq pJ(u_\lambda) + \frac{p}{\theta}(a\|u_\lambda\|_{1,p}^p + \alpha\|u_\lambda\|_{1,p}^{2p}) \\
&\quad + \frac{p}{\theta}\lambda \|h\tilde{\theta}_1\|_1 + p\lambda k \|u_\lambda\|_{1,p} \|h\max(\tilde{A}, \tilde{B})\|_1.
\end{aligned}$$

Then,

$$a\left(1 - \frac{p}{\theta}\right)\|u_\lambda\|_{1,p}^p + \left(\frac{\alpha}{2} - \frac{p\alpha}{\theta}\right)\|u_\lambda\|_{1,p}^{2p} \leq pJ(u_\lambda) + \frac{p}{\theta}\lambda \|h\tilde{\theta}_1\|_1 + p\lambda k' \|u_\lambda\|_{1,p}, \quad (3.3)$$

where  $k' = k\|h\max(\tilde{A}, \tilde{B})\|_1$ . On the other hand, since  $u_\lambda$  is a mountain pass solution, we have  $J(u_\lambda) \leq \max_{\sigma \geq 0} J(\sigma w)$  where  $w > 0$  is such that  $\|w\|_{1,p} = 1$ , and so

in view of Remark 2.3,

$$J(u_\lambda) \leq \max_{\sigma \geq 0} \frac{a}{p} \sigma^p + \frac{\alpha}{2p} \sigma^{2p} - \frac{\lambda}{q+1} D_1 \sigma^{q+1} + C_1 \lambda \|h\tilde{A}_1\|_1,$$

where

$$0 < D_1 := \int_0^1 h A_1(r_0 s^{\frac{p-1}{p-N}}) w(s)^{q+1} ds \leq k^{q+1} \|h\tilde{A}_1\|_1 \|w\|_{1,p}^{q+1} < \infty.$$

Let

$$\begin{aligned} P(\sigma) &= \frac{\alpha}{2p} \sigma^{2p} + \frac{a}{p} \sigma^p - \frac{C\lambda}{q+1} \sigma^{q+1} + C_1 \lambda \|h\tilde{A}_1\|_1, \\ P_1(\sigma) &= \left(\frac{a}{p} + \frac{\alpha}{2p}\right) \sigma^p - \frac{C\lambda}{q+1} \sigma^{q+1} + C_1 \lambda \|h\tilde{A}_1\|_1 \end{aligned}$$

and

$$P_2(\sigma) = \left(\frac{a}{p} + \frac{\alpha}{2p}\right) \sigma^{2p} - \frac{C\lambda}{q+1} \sigma^{q+1} + C_1 \lambda \|h\tilde{A}_1\|_1.$$

On  $[0, 1]$ ,  $P(\sigma) \leq P_1(\sigma)$  and on  $(1, +\infty)$ ,  $P(\sigma) \leq P_2(\sigma)$ . Also,  $P_1(\sigma)$  is maximized for  $\sigma_1 = \tilde{K}_1^{\frac{1}{q+1-p}} \lambda^{\frac{1}{q+1-p}}$  and  $P_2(\sigma)$  is maximized for  $\sigma_2 = \tilde{K}_2^r \lambda^{-r}$ , where  $\tilde{K}_1 = \frac{2a+\alpha}{2C}$  and  $\tilde{K}_2 = \frac{2a+\alpha}{C}$ . Note that if  $\lambda \leq 1$ , then  $\lambda \leq \lambda^{-2pr}$ ,  $\lambda \leq \lambda^{\frac{-p}{q+1-p}}$  and  $\lambda^{\frac{-p}{q+1-p}} \leq \lambda^{-2pr}$ . Therefore,

$$\begin{aligned} pP_1(\sigma) + \frac{p}{\theta} \lambda \|h\tilde{\theta}_1\|_1 &\leq \left(a + \frac{\alpha}{2}\right) \tilde{K}_1^{\frac{p}{q+1-p}} \lambda^{\frac{-p}{q+1-p}} + \lambda p \left(C_1 \|h\tilde{A}_1\|_1 + \frac{\|h\tilde{\theta}_1\|_1}{\theta}\right) \\ &\leq \lambda^{\frac{-p}{q+1-p}} \left(\left(a + \frac{\alpha}{2}\right) \tilde{K}_1^{\frac{p}{q+1-p}} + p \left(C_1 \|h\tilde{A}_1\|_1 + \frac{\|h\tilde{\theta}_1\|_1}{\theta}\right)\right) \\ &\leq \lambda^{-2pr} \left(\left(a + \frac{\alpha}{2}\right) \tilde{K}_1^{\frac{p}{q+1-p}} + p \left(C_1 \|h\tilde{A}_1\|_1 + \frac{\|h\tilde{\theta}_1\|_1}{\theta}\right)\right) \\ &= \tilde{C}_1 \lambda^{-2pr} \end{aligned}$$

and

$$\begin{aligned} pP_2(\sigma) + \frac{p}{\theta} \lambda \|h\tilde{\theta}_1\|_1 &\leq \left(a + \frac{\alpha}{2}\right) \tilde{K}_2^{2pr} \lambda^{-2pr} + \lambda p \left(C_1 \|h\tilde{A}_1\|_1 + \frac{\|h\tilde{\theta}_1\|_1}{\theta}\right) \\ &\leq \lambda^{-2pr} \left(\left(a + \frac{\alpha}{2}\right) \tilde{K}_2^{2pr} + p \left(C_1 \|h\tilde{A}_1\|_1 + \frac{\|h\tilde{\theta}_1\|_1}{\theta}\right)\right) \\ &= \tilde{C}_2 \lambda^{-2pr}. \end{aligned}$$

Setting  $\tilde{C}_3 = \max(\tilde{C}_1, \tilde{C}_2)$  gives

$$pJ(u_\lambda) + \frac{p}{\theta} \lambda \|h\tilde{\theta}_1\|_1 \leq \tilde{C}_3 \lambda^{-2pr},$$

and from (3.3), we have

$$\left(\frac{\alpha}{2} - \frac{p\alpha}{\theta}\right) \|u_\lambda\|_{1,p}^{2p} \leq \tilde{C}_3 \lambda^{-2pr} + p\lambda k' \|u_\lambda\|_{1,p}.$$

By Lemma 3.5, for all  $\lambda \in (0, \lambda_1)$ ,

$$\|u_\lambda\|_{1,p} \geq \frac{1}{k} \|u_\lambda\|_\infty \geq \frac{M_0}{k} \lambda^{-r \frac{p}{q+1}}.$$

Then for all  $\lambda \in \left(0, \min\left(\lambda_1, \left(\frac{M_0}{k}\right)^{\frac{q+1}{rp}}\right)\right)$ , we have  $\|u_\lambda\|_{1,p} \geq 1$ , so

$$\left(\frac{\alpha}{2} - \frac{p\alpha}{\theta}\right) \|u_\lambda\|_{1,p}^{2p} \leq \tilde{C}_3 \lambda^{-2pr} + p\lambda k' \|u_\lambda\|_{1,p}^{2p}.$$

This implies

$$\left(\frac{\alpha}{2} - \frac{p\alpha}{\theta} - p\lambda k'\right) \|u_\lambda\|_{1,p}^{2p} \leq \tilde{C}_3 \lambda^{-2pr}.$$

Hence, for all  $\lambda \in \left(0, \min\left(\lambda_1, \left(\frac{M_0}{k}\right)^{\frac{q+1}{rp}}, \frac{\alpha(\theta-2p)}{4\theta p k'}\right)\right)$ , we have

$$\frac{\alpha(\theta-2p)}{4\theta} \|u_\lambda\|_{1,p}^{2p} \leq \tilde{C}_3 \lambda^{-2pr}.$$

Taking  $C_0 = \frac{4\theta \tilde{C}_3}{\alpha(\theta-2p)}$  and  $\lambda_2 = \min\left(\lambda_1, \left(\frac{M_0}{k}\right)^{\frac{q+1}{rp}}, \frac{\alpha(\theta-2p)}{4\theta p k'}\right)$ , we see that the lemma is proved.  $\square$

To prove the positivity of the mountain pass solution, assume to the contrary, that there exists a sequence  $\{(\lambda_i, u_{\lambda_i})\}_{i=1}^\infty \subset (0, 1) \times C([0, 1])$  of mountain pass solutions to (1.2) such that  $\lambda_i \rightarrow 0$  as  $i \rightarrow \infty$  and  $m(\{x \in (0, 1) : u_{\lambda_i}(x) \leq 0\}) > 0$ . Let  $w_i = \frac{u_{\lambda_i}}{\|u_{\lambda_i}\|_\infty}$ . Since

$$-(\phi_p(u'_{\lambda_i}))' = \frac{\lambda_i h(s) f(r_0 s^{\frac{p-1}{p-N}}, u_{\lambda_i})}{a + \alpha \|u_{\lambda_i}\|_{1,p}^p},$$

we have

$$-(\phi_p(w'_i))' = \frac{\lambda_i h(s) f(r_0 s^{\frac{p-1}{p-N}}, u_{\lambda_i})}{a + \alpha \|u_{\lambda_i}\|_{1,p}^p} \|u_{\lambda_i}\|_\infty^{1-p}.$$

From Remark 2.1 and Lemmas 3.5 and 3.6, we obtain

$$\begin{aligned} \left| \frac{\lambda_i f(r_0 s^{\frac{p-1}{p-N}}, u_{\lambda_i})}{a + \alpha \|u_{\lambda_i}\|_{1,p}^p} \|u_{\lambda_i}\|_\infty^{1-p} \right| &\leq \left( \frac{\lambda_i \|u_{\lambda_i}\|_\infty^{q+1-p}}{\alpha \|u_{\lambda_i}\|_{1,p}^p} + \frac{\lambda_i M_0^{1-p} \lambda_i^{-\frac{rp(1-p)}{q+1}}}{a} \right) \max(\tilde{A}, \tilde{B}) \\ &\leq \left( \frac{\lambda_i k^{q+1-p} \|u_{\lambda_i}\|_{1,p}^{\frac{1}{r}} + \frac{M_0^{1-p}}{a}}{\alpha} \right) \max(\tilde{A}, \tilde{B}) \\ &\leq \left( \frac{\lambda_i k^{q+1-p} C_0^{\frac{1}{r}} \lambda_i^{-1} + \frac{M_0^{1-p}}{a}}{\alpha} \right) \max(\tilde{A}, \tilde{B}) \\ &\leq D_2 \max(\tilde{A}(s), \tilde{B}(s)), \end{aligned}$$



where

$$D_2 = \frac{k^{q+1-p}C_0^{\frac{1}{r}}}{\alpha} + \frac{M_0^{1-p}}{a}.$$

So for all  $s \in (0, 1)$ , the sequence  $\left\{ \frac{\lambda_i f(r_0 s^{\frac{p-1}{p-N}}, u_{\lambda_i})}{a + \alpha \|u_{\lambda_i}\|_{1,p}^p} \|u_{\lambda_i}\|_{\infty}^{1-p} \right\}$  is bounded. Thus, there exists a subsequence (named the same) that converges to a limit  $z_1(s)$ . Moreover,  $z_1(s) \geq 0$  since

$$z_1(s) = \lim_{i \rightarrow \infty} \frac{\lambda_i f(r_0 s^{\frac{p-1}{p-N}}, u_{\lambda_i})}{a + \alpha \|u_{\lambda_i}\|_{1,p}^p} \|u_{\lambda_i}\|_{\infty}^{1-p} \geq \lim_{i \rightarrow \infty} \frac{\lambda_i f(r_0 s^{\frac{p-1}{p-N}}, 0)}{a + \alpha \|u_{\lambda_i}\|_{1,p}^p} \|u_{\lambda_i}\|_{\infty}^{1-p} = 0.$$

Hence, for all  $s \in (0, 1)$ , the sequence  $\left\{ \frac{\lambda_i h(s) f(r_0 s^{\frac{p-1}{p-N}}, u_{\lambda_i})}{a + \alpha \|u_{\lambda_i}\|_{1,p}^p} \|u_{\lambda_i}\|_{\infty}^{1-p} \right\}$  converges to  $z(s) = h(s)z_1(s) \geq 0$ .

Let  $s_i \in (0, 1)$  be a maximum of  $w_i$ . Then,

$$\phi_p(w'_i(s)) = \int_s^{s_i} (-\phi_p(w'_i(\sigma)))' d\sigma = \int_s^{s_i} \frac{\lambda_i h(\sigma) f(r_0 \sigma^{\frac{p-1}{p-N}}, u_{\lambda_i}(\sigma))}{a + \alpha \|u_{\lambda_i}\|_{1,p}^p} \|u_{\lambda_i}\|_{\infty}^{1-p} d\sigma.$$

From (3.4),

$$|w'_i(s)|^{p-1} = |\phi_p(w'_i(s))| \leq \int_s^{s_i} C \max(\tilde{A}(\sigma), \tilde{B}(\sigma)) h(\sigma) d\sigma \leq C \|\max(\tilde{A}, \tilde{B})h\|_1,$$

so  $|w'_i(s)| \leq \|\max(\tilde{A}, \tilde{B})h\|_1^{\frac{1}{p-1}}$  for all  $s \in [0, 1]$ . By the Arzelà-Ascoli theorem, there exists  $w \in C([0, 1])$  such that  $w_i \rightarrow w$  in  $C([0, 1])$ .

Since  $(s_i)$  is bounded, there exists a subsequence (again denote by  $(s_i)$ ) that converges to some  $s_0$ . Again by (3.4), we have

$$\left| \frac{\lambda_i f(r_0 s^{\frac{p-1}{p-N}}, u_{\lambda_i})}{a + \alpha \|u_{\lambda_i}\|_{1,p}^p} \right| \|u_{\lambda_i}\|_{\infty}^{1-p} \leq C \max(\tilde{A}(s), \tilde{B}(s)) h(s).$$

Since  $\max(\tilde{A}, \tilde{B})h \in L^1(0, 1)$ , by the Lebesgue dominated convergence theorem,

$$\int_s^{s_i} \frac{\lambda_i h(\sigma) f(r_0 \sigma^{\frac{p-1}{p-N}}, u_{\lambda_i}(\sigma))}{a + \alpha \|u_{\lambda_i}\|_{1,p}^p} \|u_{\lambda_i}\|_{\infty}^{1-p} d\sigma \rightarrow \int_s^{s_0} z(\sigma) d\sigma.$$

Therefore,

$$\phi_p^{-1} \left( \int_s^{s_i} \frac{\lambda_i h(\sigma) f(r_0 \sigma^{\frac{p-1}{p-N}}, u_{\lambda_i}(\sigma))}{a + \alpha \|u_{\lambda_i}\|_{1,p}^p} \|u_{\lambda_i}\|_{\infty}^{1-p} d\sigma \right) \rightarrow \phi_p^{-1} \left( \int_s^{s_0} z(\sigma) d\sigma \right),$$

so we get

$$\int_0^\tau \phi_p^{-1} \left( \int_s^{s_i} \frac{\lambda_i h(\sigma) f(r_0 \sigma^{\frac{p-1}{p-N}}, u_{\lambda_i}(\sigma))}{a + \alpha \|u_{\lambda_i}\|_{1,p}^p} \|u_{\lambda_i}\|_\infty^{1-p} d\sigma \right) ds \rightarrow \int_0^\tau \phi_p^{-1} \left( \int_s^{s_0} z(\sigma) d\sigma \right) ds.$$

We see that

$$w_i(\tau) \rightarrow \int_0^\tau \phi_p^{-1} \left( \int_s^{s_0} z(\sigma) d\sigma \right) ds = w(\tau),$$

and so

$$w_i'(\tau) = \phi_p^{-1} \left( \int_\tau^{s_i} \frac{\lambda_i h(\sigma) f(r_0 \sigma^{\frac{p-1}{p-N}}, u_{\lambda_i}(\sigma))}{a + \alpha \|u_{\lambda_i}\|_{1,p}^p} \|u_{\lambda_i}\|_\infty^{1-p} d\sigma \right)$$

converges to  $\phi_p^{-1} \left( \int_\tau^{s_0} z(\sigma) d\sigma \right) = w'(\tau)$  for all  $\tau \in [0, 1]$ . Hence,  $-(\phi_p(w'))' = z \geq 0$  with  $w(0) = 0 = w(1)$ . Since  $\|w\|_\infty = 1$ , clearly  $w \neq 0$ . Then, since  $w$  is concave,  $w > 0$  in  $(0, 1)$ ,  $w'(0) > 0$ , and  $w'(1) < 0$ . Because  $w_i \rightarrow w$  in  $C([0, 1])$ , we conclude that  $w_i(s) > 0$  for all  $s \in (0, 1)$  for  $i$  sufficiently large. Hence,  $u_{\lambda_i}(s) > 0$  for all  $s \in (0, 1)$  for  $i$  sufficiently large. This contradicts  $m(\{x \in (0, 1) : u_{\lambda_i}(x) \leq 0\}) > 0$  for all sufficiently large  $i$ .

Thus, the mountain pass solution is positive, and this completes the proof of Theorem 1.5.


## REFERENCES

- [1] N. Aissaoui, W. Long, *Positive solutions for a Kirchhoff equation with perturbed source terms*, Acta Math. Scientia **42** (2022), 1817–1830.
- [2] C.O. Alves, F.J.S.A. Corrêa, T.F. Ma, *Positive solutions for a quasilinear elliptic equation of Kirchhoff type*, Comput. Math. Appl. **49** (2005), 85–93.
- [3] A. Ambrosetti, P.H. Rabinowitz, *Dual variational methods in critical point theory and applications*, J. Funct. Anal. **14** (1973), 349–381.
- [4] M. Badiale, E. Serra, *Semilinear Elliptic Equations for Beginners. Existence Results via the Variational Approach*, Universitext, Springer, London, 2011.
- [5] H. Brézis, *Functional Analysis, Sobolev Spaces and Partial Differential Equations*, Springer, New York, 2010.
- [6] D. Butler, E. Ko, E.K. Lee, R. Shivaji, *Positive radial solutions for elliptic equations on exterior domains with nonlinear boundary conditions*, Comm. Pure Appl. Anal. **13** (2014), 2713–2731.
- [7] A. Castro, R. Shivaji, *Nonnegative solutions for a class of nonpositone problems*, Proc. Roy. Soc. Edinburgh Sect. A **108** (1988), 291–302.
- [8] A. Castro, D.G. de Figueiredo, E. Lopera, *Existence of positive solutions for a semipositone  $p$ -Laplacian problem*, Proc. Roy. Soc. Edinburgh Sect. A **146** (2016), 475–482.

- [9] R. Dhanya, Q. Morris, R. Shivaji, *Existence of positive radial solutions for superlinear semipositone problems on the exterior of a ball*, J. Math. Anal. Appl. **434** (2016), 1533–1548.
- [10] M. Ding, C. Zhang, S. Zhou, *Local boundedness and Hölder continuity for the parabolic fractional  $p$ -Laplace equations*, Calc. Var. Partial Differential Equations **60** (2021), Article no. 38.
- [11] L. Gasinski, N.S. Papageorgiou, *Nonsmooth Critical Point Theory and Nonlinear Boundary Value Problems*, Series in Mathematical Analysis and Applications, vol. 8, Chapman & Hall/CRC, Boca Raton, 2005.
- [12] J.R. Graef, S. Heidarkhani, L. Kong, *A variational approach to a Kirchhoff-type problem involving two parameters*, Results Math. **63** (2013), 877–889.
- [13] J.R. Graef, S. Heidarkhani, L. Kong, *Variational-hemivariational inequalities of Kirchhoff-type with small perturbations of nonhomogeneous Neumann boundary conditions*, Math. Eng. Sci. Aero. **8** (2017), 345–357.
- [14] J.R. Graef, S. Heidarkhani, L. Kong, S. Moradi, *On an anisotropic discrete boundary value problem of Kirchhoff type*, J. Difference Equ. Appl. **27** (2021), 1103–1119.
- [15] J.R. Graef, S. Heidarkhani, L. Kong, A. Ghobadi, *Existence of multiple solutions to a  $P$ -Kirchhoff problem*, Differ. Equ. Appl. **14** (2022), 227–237.
- [16] L. Guo, Y. Sun, G. Shi, *Ground states for fractional nonlocal equations with logarithmic nonlinearity*, Opuscula Math. **42** (2022), 157–178.
- [17] D.D. Hai, *Positive radial solutions for singular quasilinear elliptic equations in a ball*, Publ. Res. Inst. Math. Sci. **50** (2014), 341–362.
- [18] W. He, D. Qin, Q. Wu, *Existence, multiplicity and nonexistence results for Kirchhoff type equations*, Adv. Nonlinear Anal. **10** (2021), 616–635.
- [19] G. Kirchhoff, *Mechanik*, Teubner, Leipzig, 1883.
- [20] Q. Morris, R. Shivaji, I. Sim, *Existence of positive radial solutions for a superlinear semipositone  $p$ -Laplacian problem on the exterior of a ball*, Proc. Roy. Soc. Edinburgh Sect. A **148** (2018), 409–428.
- [21] H. Pi, Y. Zeng, *Existence results for the Kirchhoff type equation with a general nonlinear term*, Acta Math. Scientia **42** (2022), 2063–2077.
- [22] D. Qin, V.D. Radulescu, X. Tang, *Ground states and geometrically distinct solutions for periodic Choquard–Pekar equations*, J. Differential Equations **275** (2021), 652–683.
- [23] J. Smoller, A. Wasserman, *Existence of positive solutions for semilinear elliptic equations in general domains*, Arch. Ration. Mech. Anal. **98** (1987), 229–249.
- [24] L. Wang, K. Xie, B. Zhang, *Existence and multiplicity of solutions for critical Kirchhoff-type  $p$ -Laplacian problems*, J. Math. Anal. Appl. **458** (2018), 361–378.

John R. Graef (corresponding author)

john-graef@utc.edu

 <https://orcid.org/0000-0002-8149-4633>

University of Tennessee at Chattanooga

Department of Mathematics

Chattanooga, TN 37403, USA

Doudja Hebboul

doudja.hebboul@g.ens-kouba.dz

Ecole Normale Supérieure

Laboratory of Partial Differential Equations and History of Mathematics

Kouba, Algiers, Algeria

Toufik Moussaoui

toufik.moussaoui@g.ens-kouba.dz

Ecole Normale Supérieure

Laboratory of Fixed Point Theory and Applications

Kouba, Algiers, Algeria

*Received: August 17, 2022.*

*Revised: November 7, 2022.*

*Accepted: November 8, 2022.*