

Dark and singular optical solitons with competing nonlocal nonlinearities

QIN ZHOU^{1*}, LAN LIU¹, HUIJUAN ZHANG¹, MOHAMMAD MIRZAZADEH², ALI H. BHRAWY^{3,4},
ESSAID ZERRAD⁵, SEITHUTI MOSHOKOA⁶, ANJAN BISWAS^{7,3}

¹School of Electronics and Information Engineering, Wuhan Donghu University,
Wuhan 430212, P.R. China

²Department of Engineering Sciences, Faculty of Technology and Engineering,
East of Guilan, University of Guilan, 44891-63157 Rudsar-Vajargah, Iran

³Department of Mathematics, Faculty of Science, King Abdulaziz University,
Jeddah-21589, Saudi Arabia

⁴Department of Mathematics, Faculty of Science, Beni-Suef University, Beni-Suef, Egypt

⁵Department of Physics and Engineering, Delaware State University, Dover, DE 19901-2277, USA

⁶Department of Mathematics and Statistics, Tshwane University of Technology,
Pretoria-0008, South Africa

⁷Department of Mathematical Sciences, Delaware State University, Dover, DE 19901-2277, USA

*Corresponding author: qinzhou@whu.edu.cn, qzh@whu.edu.cn

In this work, we study the dynamics of optical solitons in a synthetic nonlocal nonlinear media. The nonlinear dynamical model which describes the propagation of optical solitons in the weakly nonlocal nonlinear media with parabolic law nonlinearity is investigated analytically. The tool of integration that is the Riccati equation mapping approach is introduced to extract exact traveling wave solutions. As a result, an explicit dark soliton, singular soliton and periodic solutions are derived.

Keywords: solitons, parabolic law nonlinearity, competing nonlinearities.

1. Introduction

The study of dynamical behaviours of optical solitons in nonlocal nonlinear media has greatly attracted researchers' attention in recent years [1–30]. Nonlocality of nonlinearity, which includes weak nonlocality, general nonlocality and strong nonlocality, exists in nematic liquid crystals, plasmas and many other nonlinear systems [4, 9, 13, 17]. In the paraxial approximation, the propagation of optical solitons in a nonlocal nonlinear

medium is ruled by the well-known nonlocal nonlinear Schrödinger equation (NNLSE) in the form [7, 10, 17, 18]

$$i \frac{\partial u}{\partial z} + \frac{1}{2} \frac{\partial^2 u}{\partial z^2} + \Delta n u = 0 \quad (1a)$$

where $u(x, z)$ represents the normalized complex wave envelope that is the function of transverse coordinate x and longitudinal coordinate z ;

$$\Delta n(x, z) = s \int_{-\infty}^{+\infty} R(x' - x) |u(x', z)|^2 dx' \quad (1b)$$

refers to the refractive index change that depends on optical pulse intensity $|u(x, z)|^2$, in which $R(x)$ is the nonlocal response function, and $s = \pm 1$ determines the types of nonlinearity, specifically, the plus (minus) sign corresponds to the self-focusing (self-defocusing) nonlinearity.

For strongly nonlocal medium, Δn is proportional to x^2 , then NNLSE (Eq. (1a)) reduces to a linear differential equation, also known as the Snyder–Mitchell model.

For weakly nonlocal medium $\Delta n = s[|u|^2 + \gamma \partial_x^2(|u|^2)]$ in which γ is a small constant. Then, in this case, NNLSE becomes [10]

$$i \frac{\partial u}{\partial z} + \frac{1}{2} \frac{\partial^2 u}{\partial x^2} + s|u|^2 u + \gamma s \frac{\partial^2(|u|^2)}{\partial x^2} u = 0 \quad (2)$$

Recently, there has been a growing interest in the study of optical solutions in the nonlocal nonlinear media with competing nonlinearities where the nonlinear response is the result of contributions from nonlocal nonlinearity and other nonlinear effects. CHEN *et al.* [4] reported the dynamical behaviours of dark solitons in a medium with competing nonlocal nonlinearity and quintic nonlinearity. Bang's research team [7, 9, 13, 16] studied the properties of optical solitons in a medium with competing nonlocal self-focusing and self-defocusing nonlinearities. In our previous studies [5, 10, 12, 15, 17], exact solitons in a medium with competing weakly nonlocal nonlinearity and parabolic law nonlinearity, as well as a medium with competing nonlocal self-focusing nonlinearity, self-defocusing nonlinearity and quintic nonlinearity, are derived.

The aim of the present work is to study the dynamics of optical solitons in a medium with competing weakly nonlocal nonlinearity and parabolic law nonlinearity. Without loss of generality, the nonlinear dynamical model is given by [5, 10]

$$i \frac{\partial u}{\partial z} + a \frac{\partial^2 u}{\partial x^2} + b \frac{\partial^2(|u|^2)}{\partial x^2} u + c|u|^2 u + \mu|u|^4 u = 0 \quad (3)$$

where a represents the coefficients of diffraction, while b gives the coefficients of weakly nonlocal nonlinearity, and finally the terms with c and μ account for the parabolic law (cubic-quintic) nonlinearity [5, 10, 31, 32].

It should be noted that Eq. (3) has been investigated analytically by traveling wave hypothesis, Jacobi's elliptic equations expansion approach and ansatz scheme [5, 10]. This work is an extension of earlier studies. We will use a different algorithm to integrate Eq. (3). It is the generalized Riccati equation expansion technique [33–41]. Finally, explicit traveling wave solutions, along with the integrability conditions, are reported.

2. Exact solitons

Assume that Eq. (3) admits the following stationary solutions:

$$u(x, z) = \psi(x) \exp(i\lambda z) \quad (4)$$

where λ is the propagation constant.

Substituting the above hypothesis (4) into Eq. (3) yields

$$(a + 2b\psi^2) \frac{d^2\psi}{dx^2} + 2b\psi \left(\frac{d\psi}{dx} \right)^2 - \lambda\psi + c\psi^3 + \mu\psi^5 = 0 \quad (5)$$

Below, we will perform the Riccati equation mapping scheme to construct explicit traveling wave solutions to Eq. (5). Based on the homogeneous balance principle, Eq. (5) has the solution in the form

$$\psi(x) = l_0 + l_1 \varphi(x) \quad (6)$$

where l_0 and l_1 are real constants to be determined later, and $\varphi(x)$ is the solution of the following second-order nonlinear ordinary differential equation:

$$\varphi'(x) = \alpha + \beta\varphi(x) + \delta\varphi^2(x) \quad (7)$$

with α , β and δ being real constants. Equation (7) is the famous generalized Riccati equation, exact solutions of which are given in [33–41].

Substituting Eqs. (6) and (7) into Eq. (5), and vanishing the coefficients of φ^i ($i = 0, 1, 2, 3, 4, 5$) yield a system of nonlinear algebraic equations which solves to:

$$l_0 = 0 \quad (8)$$

$$l_1 = \pm \sqrt{-\frac{6b\delta^2}{\mu}} \quad (9)$$

$$\alpha = \frac{a\mu - 3bc}{24b^2\delta} \quad (10)$$

$$\beta = 0 \quad (11)$$

$$\lambda = \frac{a^2\mu - 3abc}{12b^2} - \frac{(a\mu - 3bc)^2}{48\mu b^2} \quad (12)$$

where a, b, c, μ and δ are arbitrary constants.

Equation (9) naturally introduces the constraint

$$b\mu < 0 \quad (13)$$

which means that the coefficients of weakly nonlocal nonlinearity and cubic-quintic nonlinearity have the opposite sign.

Equation (12) gives the expression of propagation constant.

Finally, we can obtain analytical traveling wave solutions to Eq. (3), which are listed as follows:

Case 1: periodic solutions and periodic singular solutions. Equation (3) admits 10 periodic traveling wave solutions, which are given by

$$u_1(x, z) = \pm \sqrt{-\frac{6b\alpha\delta}{\mu}} \tan(\sqrt{\alpha\delta}x) \exp(i\lambda z) \quad (14)$$

$$u_2(x, z) = \mp \sqrt{-\frac{6b\alpha\delta}{\mu}} \cot(\sqrt{\alpha\delta}x) \exp(i\lambda z) \quad (15)$$

$$u_3(x, z) = \pm \sqrt{-\frac{6b\alpha\delta}{\mu}} \left[\tan(\sqrt{4\alpha\delta}x) \pm \sec(\sqrt{4\alpha\delta}x) \right] \exp(i\lambda z) \quad (16)$$

$$u_4(x, z) = \mp \sqrt{-\frac{6b\alpha\delta}{\mu}} \left[\cot(\sqrt{4\alpha\delta}x) \pm \csc(\sqrt{4\alpha\delta}x) \right] \exp(i\lambda z) \quad (17)$$

$$u_5(x, z) = \pm \sqrt{-\frac{3b\alpha\delta}{2\mu}} \left[\tan\left(\frac{\sqrt{\alpha\delta}}{2}x\right) - \coth\left(\frac{\sqrt{\alpha\delta}}{2}x\right) \right] \exp(i\lambda z) \quad (18)$$

$$u_6(x, z) = \pm \sqrt{-\frac{6b\alpha\delta}{\mu}} \frac{\pm\sqrt{A^2 - B^2} - A \cos(\sqrt{4\alpha\delta}x)}{A \sinh(\sqrt{4\alpha\delta}) + B} \exp(i\lambda z) \quad (19)$$

$$u_7(x, z) = \pm \sqrt{-\frac{6b\alpha\delta}{\mu}} \frac{\mp\sqrt{A^2 - B^2} - A \cos(\sqrt{4\alpha\delta}x)}{A \sinh(\sqrt{4\alpha\delta}) + B} \exp(i\lambda z) \quad (20)$$

$$u_8(x, z) = \mp \sqrt{-\frac{6b\alpha\delta}{\mu}} \frac{\cos(\sqrt{\alpha\delta}x)}{\sin(\sqrt{4\alpha\delta}) \pm 1} \exp(i\lambda z) \quad (21)$$

$$u_9(x, z) = \pm \sqrt{-\frac{6b\alpha\delta}{\mu}} \frac{\sin(\sqrt{\alpha\delta}x)}{\cos(\sqrt{4\alpha\delta}) \pm 1} \exp(i\lambda z) \quad (22)$$

$$u_{10}(x, z) = \pm 2 \sqrt{-\frac{6b\alpha\delta}{\mu}} \frac{\sin\left(\frac{\sqrt{\alpha\delta}}{2}\right) \cos\left(\frac{\sqrt{\alpha\delta}}{2}\right)}{2\cos^2\left(\frac{\sqrt{\alpha\delta}}{2}\right) - 1} \exp(i\lambda z) \quad (23)$$

which naturally pose the constraint conditions

$$b\alpha\delta\mu < 0 \quad (24)$$

$$A^2 - B^2 > 0 \quad (25)$$

where both A and B are constants.

Here, we noted that the periodic solutions and periodic singular solutions are not necessary because these periodic solutions are of no interest in optical fibers at all.

Case 2: soliton solutions. Equation (3) admits a dark soliton solution

$$u_{11}(x, z) = \mp \sqrt{\frac{6b\alpha\delta}{\mu}} \tanh(\sqrt{-\alpha\delta}x) \exp(i\lambda z) \quad (26)$$

and singular soliton solutions

$$u_{12}(x, z) = \mp \sqrt{\frac{6b\alpha\delta}{\mu}} \coth(\sqrt{-\alpha\delta}x) \exp(i\lambda z) \quad (27)$$

which naturally introduce the restriction

$$b\alpha\delta\mu > 0 \quad (28)$$

3. Conclusion

The dynamics of optical solitons in a medium with competing weakly nonlocal nonlinearity and parabolic law nonlinearity has been studied analytically. The integration tool that is the Riccati equation mapping approach is adopted to retrieve exact solutions. Finally, ten periodic traveling wave solutions and two soliton solutions are reported. The presented results show that the first restriction for the existence of those explicit traveling wave solutions is $b\mu < 0$, *i.e.*, the coefficients of weakly nonlocal nonlinearity and cubic-quintic nonlinearity have the opposite sign.

In the future studies, we plan to present the analytic study on optical solitons in a medium with competing weakly nonlocal nonlinearity and other types of non-Kerr law nonlinearity. They are power law and dual-power law. The results will be published in other papers.

Acknowledgements – This work was supported by the Natural Science Foundation of Hubei Province of China under grant number 2015CFC891.

References

- [1] MIHALACHE D., MAZILU D., LEDERER F., MALOMED B.A., KARTASHOV Y.V., CRASOVAN L.-C., TORNER L., *Three-dimensional spatiotemporal optical solitons in nonlocal nonlinear media*, Physical Review E **73**(2), 2006, article 025601R.
- [2] MIHALACHE D., *Linear and nonlinear light bullets: Recent theoretical and experimental studies*, Romanian Journal of Physics **57**(1–2), 2012, pp. 352–371.
- [3] NIKOLOV N.I., NESHEV D., KRÓLIKOWSKI W., OLE BANG, RASMUSSEN J.J., CHRISTIANSEN P.L., *Attraction of nonlocal dark optical solitons*, Optics Letters **29**(3), 2004, pp. 286–288.
- [4] CHEN L., WANG Q., SHEN M., ZHAO H., LIN Y.-Y., JENG C.-C., LEE R.-K., KRÓLIKOWSKI W., *Nonlocal dark solitons under competing cubic-quintic nonlinearities*, Optics Letters **38**(1), 2013, pp. 13–15.
- [5] QIN ZHOU, QIUPING ZHU, SAVESCU M., BHRRAWY A., BISWAS A., *Optical solitons with nonlinear dispersion in parabolic law medium*, Proceedings of the Romanian Academy, Series A **16** (2), 2015, pp. 152–159.
- [6] MIHALACHE D., *Multidimensional localized structures in optics and Bose–Einstein condensates: a selection of recent studies*, Romanian Journal of Physics **59**(3–4), 2014, pp. 295–312.
- [7] KRÓLIKOWSKI W., OLE BANG, *Solitons in nonlocal nonlinear media: exact solutions*, Physical Review E **63**(1), 2001, article 016610.
- [8] WEI-PING ZHONG, BELIĆ M., *Three-dimensional optical vortex and necklace solitons in highly nonlocal nonlinear media*, Physical Review A **79**(2), 2009, article 023804.
- [9] KONG Q., WANG Q., BANG O., KRÓLIKOWSKI W., *Analytical theory for the dark-soliton interaction in nonlocal nonlinear materials with an arbitrary degree of nonlocality*, Physical Review A **82**(1), 2010, article 013826.
- [10] QIN ZHOU, DUANZHENG YAO, SIJING DING, YAFANG ZHANG, FENG CHEN, FANG CHEN, XIAONA LIU, *Spatial optical solitons in fifth order and seventh order weakly nonlocal nonlinear media*, Optik – International Journal for Light and Electron Optics **124**(22), 2013, pp. 5683–5686.
- [11] XINGHUI GAO, CHENGYUN ZHANG, DONG TANG, HUI ZHENG, DAQUAN LU, WEI HU, *High-order dark solitons in nonlocal nonlinear media*, Journal of Modern Optics **60**(15), 2013, pp. 1281–1286.
- [12] QIN ZHOU, DUANZHENG YAO, XIAONA LIU, FANG CHEN, SIJING DING, YAFANG ZHANG, FENG CHEN, *Exact solitons in three-dimensional weakly nonlocal nonlinear time-modulated parabolic law media*, Optics and Laser Technology **51**, 2013, pp. 32–35.
- [13] QIAN KONG, MING SHEN, ZHENYI CHEN, QI WANG, RAY-KUANG LEE, KRÓLIKOWSKI W., *Dark solitons in nonlocal media with competing nonlinearities*, Physical Review A **87**(6), 2013, article 063832.
- [14] MING SHEN, YUANYAO LIN, CHIEN-CHUNG JENG, RAY-KUANG LEE, *Vortex pairs in nonlocal nonlinear media*, Journal of Optics **14**(6), 2012, article 065204.
- [15] QIN ZHOU, QIUPING ZHU, YAXIAN LIU, PING YAO, BHRRAWY A.H., MORARU L., BISWAS A., *Bright-dark combo optical solitons with non-local nonlinearity in parabolic law medium*, Optoelectronics and Advanced Materials – Rapid Communications **8**(9–10), 2014, pp. 837–839.
- [16] ESBENSEN B.K., WLOTZKA A., BACHE M., BANG O., KRÓLIKOWSKI W., *Modulational instability and solitons in nonlocal media with competing nonlinearities*, Physical Review A **84**(5), 2011, article 053854.
- [17] QIN ZHOU, QIUPING ZHU, *Theoretical study of dark solitons in media with competing nonlocal nonlinearities and local quintic nonlinearity*, Journal of Modern Optics **61**(18), 2014, pp. 1465–1469.

- [18] DING NA, GUO QI, *The $(1+1)$ -dimensional spatial solitons in media with weak nonlinear nonlocality*, Chinese Physics B **18**(10), 2009, pp. 4298–4302.
- [19] TSOY E.N., *Solitons in weakly nonlocal media with cubic-quintic nonlinearity*, Physical Review A **82**(6), 2010, article 063829.
- [20] YANWEI DU, ZHONGXIANG ZHOU, HAO TIAN, DAJUN LIU, *Bright solitons and repulsive in-phase interaction in media with competing nonlocal Kerr nonlinearities*, Journal of Optics **13**(1), 2011, article 015201.
- [21] RUI GUO, YUE-FENG LIU, HUI-QIN HAO, FENG-HUA QI, *Coherently coupled solitons, breathers and rogue waves for polarized optical waves in an isotropic medium*, Nonlinear Dynamics **80**(3), 2015, pp. 1221–1230.
- [22] QIN ZHOU, QIUPING ZHU, BHRAWY A.H., BISWAS A., *Combined optical solitons with nonlinear dispersion and spatio-temporal dispersion*, Optoelectronics and Advanced Materials – Rapid Communications **9**(1–2), 2015, pp. 1–4.
- [23] GANG-WEI WANG, TIAN-ZHOU XU, EBADI G., JOHNSON S., STRONG A.J., BISWAS A., *Singular solitons, shock waves, and other solutions to potential KdV equation*, Nonlinear Dynamics **76**(2), 2014, pp. 1059–1068.
- [24] QIN ZHOU, DUANZHENG YAO, QIANQIAN XU, XIAONA LIU, *Optical soliton perturbation with time- and space-dependent dissipation (or gain) and nonlinear dispersion in Kerr and non-Kerr media*, Optik – International Journal for Light and Electron Optics **124**(16), 2013, pp. 2368–2372.
- [25] ZHOU Q., ZHU Q., BHRAWY A.H., MORARU L., BISWAS A., *Optical solitons with spatially-dependent coefficients by Lie symmetry*, Optoelectronics and Advanced Materials – Rapid Communications **8**(7–8), 2014, pp. 800–803.
- [26] QIN ZHOU, *Analytical solutions and modulation instability analysis to the perturbed nonlinear Schrödinger equation*, Journal of Modern Optics **61**(6), 2014, pp. 500–503.
- [27] QIN ZHOU, QIUPING ZHU, YAXIAN LIU, BISWAS A., BHRAWY A.H., KHAN K.R., MAHMOOD M.F., BELIC M., *Solitons in optical metamaterials with parabolic law nonlinearity and spatio-temporal dispersion*, Journal of Optoelectronics and Advanced Materials **16**(11–12), 2014, pp. 1221–1225.
- [28] MIHALACHE D., *Stable vortex solitons and robust soliton complexes in optical media with competing nonlinearities*, Romanian Journal of Physics **50**(3–4), 2005, pp. 357–364.
- [29] MIHALACHE D., MAZILU D., *Ginzburg–Landau spatiotemporal dissipative optical solitons*, Romanian Reports in Physics **60**(3), 2008, pp. 749–761.
- [30] SAVESCU M., ALSHAERY A.A., HILAL E.M., BHRAWY A.H., QIN ZHOU, BISWAS A., *Optical solitons in DWDM system with four-wave mixing*, Optoelectronics and Advanced Materials – Rapid Communications **9**(1–2), 2015, pp. 14–19.
- [31] VEGA-GUZMAN J., ZHOU Q., ALSHAERY A.A., HILAL E.M., BHRAWY A.H., BISWAS A., *Optical solitons in cascaded system with spatio-temporal dispersion by ansatz approach*, Journal of Optoelectronics and Advanced Materials **17**(1–2), 2015, pp. 165–171.
- [32] YANAN XU, QIN ZHOU, BHRAWY A.H., KHAN K.R., MAHMOOD M.F., BISWAS A., BELIC M., *Bright solitons in optical metamaterials by traveling wave hypothesis*, Optoelectronics and Advanced Materials – Rapid Communications **9**(3–4), 2015, pp. 384–387.
- [33] QIN ZHOU, QIUPING ZHU, BISWAS A., *Optical solitons in birefringent fibers with parabolic law nonlinearity*, Optica Applicata **44**(3), 2014, pp. 399–409.
- [34] SHUN-DONG ZHU, *The generalizing Riccati equation mapping method in non-linear evolution equation: application to $(2+1)$ -dimensional Boiti–Leon–Pempinelle equation*, Chaos, Solitons and Fractals **37**(5), 2008, pp. 1335–1342.
- [35] QIN ZHOU, LAN LIU, YAXIAN LIU, HUA YU, PING YAO, CHUN WEI, HUIJUAN ZHANG, *Exact optical solitons in metamaterials with cubic–quintic nonlinearity and third-order dispersion*, Nonlinear Dynamics **80**(3), 2015, pp. 1365–1371.
- [36] VITANOV N.K., *Application of simplest equations of Bernoulli and Riccati kind for obtaining exact traveling-wave solutions for a class of PDEs with polynomial nonlinearity*, Communications in Nonlinear Science and Numerical Simulation **15**(8), 2010, pp. 2050–2060.

- [37] QI WANG, YONG CHEN, HONGQING ZHANG, *A new Riccati equation rational expansion method and its application to (2+1)-dimensional Burgers equation*, *Chaos, Solitons and Fractals* **25**(5), 2005, pp. 1019–1028.
- [38] LIANG-QIAN KONG, CHAO-QING DAI, *Some discussions about variable separation of nonlinear models using Riccati equation expansion method*, *Nonlinear Dynamics* **81**(3), 2015, pp. 1553–1561.
- [39] QIN ZHOU, QIUPING ZHU, HUA YU, YAXIAN LIU, CHUN WEI, PING YAO, BHRAWY A.H., BISWAS A., *Bright, dark and singular optical solitons in a cascaded system*, *Laser Physics* **25**(2), 2015, article 025402.
- [40] QIN ZHOU, QIUPING ZHU, YAXIAN LIU, HUA YU, PING YAO, BISWAS A., *Thirring optical solitons in birefringent fibers with spatio-temporal dispersion and Kerr law nonlinearity*, *Laser Physics* **25**(1), 2015, article 015402.
- [41] BISWAS A., MIRZAZADEH M., SAVESCU M., MILOVIC D., KHAN K.R., MAHMOOD M.F., BELIC M., *Singular solitons in optical metamaterials by ansatz method and simplest equation approach*, *Journal of Modern Optics* **61**(19), 2014, pp. 1550–1555.

*Received August 16, 2015
in revised form September 16, 2015*