

Number of Zeros of a Polynomial (Lacunary-type) in a Disk

Idrees Qasim, Tawheeda Rasool and Abdul Liman

ABSTRACT: The problem of finding out the region which contains all or a prescribed number of zeros of a polynomial $P(z) := \sum_{j=0}^n a_j z^j$ has a long history and dates back to the earliest days when the geometrical representation of complex numbers was introduced. In this paper, we present certain results concerning the location of the zeros of Lacunary-type polynomials $P(z) := a_0 + \sum_{j=\mu}^n a_j z^j$ in a disc centered at the origin.

AMS Subject Classification: 30A01, 30C10, 30C15.

Keywords and Phrases: Zeros; Lacunary polynomial; Prescribed region.

1. Introduction and Statement of Results

The problem of locating some or all the zeros of a given polynomial as a function of its coefficients is of long standing interest in mathematics. This fact can be visualized by glancing at the references in the comprehensive books of Marden [9] and Milovanovic, Mitrinovic and Rassias [10], Rahman and Schmeisser [12] and by noting the abundance of recent publications on the subject [7, 8, 13].

Regarding the least number of zeros of polynomial $P(z) = \sum_{j=0}^n a_j z^j$ in a given circle Mohammad [11] proved the following:

Theorem A. Let $P(z) := \sum_{j=0}^n a_j z^j$ be a polynomial of degree n such that

$$a_n \geq a_{n-1} \geq a_{n-2} \geq \cdots \geq a_1 \geq a_0 > 0,$$

then the number of zeros of $P(z)$ in $|z| \leq \frac{1}{2}$ does not exceed

$$1 + \frac{1}{\log 2} \log \frac{a_n}{a_0}.$$

Dewan [3] generalized Theorem A to polynomials with complex coefficients and proved the following result:

Theorem B. Let $P(z) := \sum_{j=0}^n a_j z^j$ be a polynomial of degree n with complex coefficients. If $\operatorname{Re} a_j = \alpha_j$ and $\operatorname{Im} a_j = \beta_j$, $j = 0, 1, 2, \dots, n$ such that

$$\alpha_n \geq \alpha_{n-1} \geq \alpha_{n-2} \geq \dots \geq \alpha_1 \geq \alpha_0 > 0,$$

then the number of zeros of $P(z)$ in $|z| \leq \frac{1}{2}$ does not exceed

$$1 + \frac{1}{\log 2} \log \frac{\alpha_n + \sum_{j=0}^n |\beta_j|}{|a_0|}.$$

In this direction, recently Irshad et al [1] proved the following:

Theorem C. Let $P(z) := \sum_{j=0}^n a_j z^j$ be a polynomial of degree n with complex coefficients such that for some $\lambda \geq 1$, $0 \leq k \leq n$,

$$|a_n| \leq |a_{n-1}| \leq \dots \leq |a_{k+1}| \leq \lambda |a_k| \geq |a_{k-1}| \geq \dots \geq |a_1| \geq |a_0|,$$

and for some real β ,

$$|\arg a_j - \beta| \leq \alpha \leq \frac{\pi}{2}, \quad j = 0, 1, 2, \dots, n$$

then the number of zeros of $P(z)$ in $|z| \leq \frac{1}{2}$ does not exceed

$$\left. \begin{aligned} & \frac{1}{\log 2} \log \left\{ \frac{2\lambda |a_k| \cos \alpha + 2|\lambda - 1| |a_k| \sin \alpha}{|a_0|} \right. \\ & \left. + \frac{|a_n| (\sin \alpha - \cos \alpha + 1) + 2 \sin \alpha \sum_{j=0}^{n-1} |a_j| + 2|1 - \lambda| |a_k|}{|a_0|} \right\}. \end{aligned} \right\}$$

Chan and Malik [2] introduced the class of Lacunary polynomials of the form $P(z) := a_0 + \sum_{j=\mu}^n a_j z^j$, where $a_0 \neq 0$. Notice that when $\mu = 1$, we simply have the class of all polynomials of degree n . In [5] and [6] Landau proved that every trinomial

$$a_0 + a_1 z + a_n z^n, a_1 a_n \neq 0, n \geq 2$$

has at least one zero in the circle $|z| \leq 2 \frac{|a_0|}{|a_1|}$ and that of quadrinomial

$$a_0 + a_1 z + a_m z^m + a_n z^n, a_1 a_m a_n \neq 0, 2 \leq m \leq n$$

has at least one zero in the circle $|z| \leq \frac{17}{3} \left| \frac{a_0}{a_1} \right|$. These two polynomials are of the Lacunary-type $P(z) := a_0 + \sum_{j=\mu}^n a_j z^j$.

The aim of this paper is to study the number of zeros in a disc centered at the origin for such class of polynomials. We begin by proving the following result putting restrictions on the moduli of the coefficients. In fact we prove:

Theorem 1. *Let $P(z) := a_0 + \sum_{j=\mu}^n a_j z^j$, $1 \leq \mu \leq n - 1$, $a_0 \neq 0$, be a polynomial of degree n . If for some real α and β*

$$|\arg a_j - \beta| \leq \alpha \leq \frac{\pi}{2}, \quad \mu \leq j \leq n$$

and for some $t > 0$ and some k with $\mu \leq k \leq n$,

$$t^\mu |a_\mu| \leq \dots \leq t^{k-1} |a_{k-1}| \leq t^k |a_k| \geq t^{k+1} |a_{k+1}| \geq \dots \geq t^{n-1} |a_{n-1}| \geq t^n |a_n|$$

then the number of zeros of $P(z)$ in $|z| \leq \frac{t}{2}$ does not exceed

$$\frac{1}{\log 2} \log \frac{M}{|a_0|},$$

where

$$M = 2|a_0|t + |a_\mu|t^{\mu+1}(1 - \sin \alpha - \cos \alpha) + 2|a_k|t^{k+1} \cos \alpha + |a_n|t^{n+1}(1 - \sin \alpha - \cos \alpha) + 2 \sum_{j=\mu}^n |a_j|t^{j+1} \sin \alpha.$$

For $t = 1$, we get the following:

Corollary 1.1. *Let $P(z) := a_0 + \sum_{j=\mu}^n a_j z^j$, $1 \leq \mu \leq n - 1$, $a_0 \neq 0$ be a polynomial of degree n . If for some real α and β*

$$|\arg a_j - \beta| \leq \alpha \leq \frac{\pi}{2}, \quad \mu \leq j \leq n$$

and some k with

$$|a_\mu| \leq \dots \leq |a_{k-1}| \leq |a_k| \geq |a_{k+1}| \geq \dots \geq |a_{n-1}| \geq |a_n|$$

then the number of zeros of $P(z)$ in $|z| \leq \frac{1}{2}$ does not exceed

$$\frac{1}{\log 2} \log \frac{M}{|a_0|},$$

where

$$M = 2|a_0| + |a_\mu|(1 - \sin \alpha - \cos \alpha) + 2|a_k| \cos \alpha + |a_n|(1 - \sin \alpha - \cos \alpha) + 2 \sum_{j=\mu}^n |a_j| \sin \alpha.$$

With $k = n$ in Corollary 1.1, we get:

Corollary 1.2. Let $P(z) := a_0 + \sum_{j=\mu}^n a_j z^j$, $1 \leq \mu \leq n-1$, $a_0 \neq 0$ be a polynomial of degree n . If for some real α and β

$$|\arg a_j - \beta| \leq \alpha \leq \frac{\pi}{2}, \quad \mu \leq j \leq n$$

such that

$$|a_\mu| \leq \cdots \leq |a_{n-1}| \leq |a_n|$$

then the number of zeros of $P(z)$ in $|z| \leq \frac{1}{2}$ does not exceed

$$\frac{1}{\log 2} \log \frac{M}{|a_0|},$$

where

$$M = 2|a_0| + |a_\mu|(1 - \sin \alpha - \cos \alpha) + |a_n|(1 - \sin \alpha + \cos \alpha) + 2 \sum_{j=\mu}^n |a_j| \sin \alpha.$$

Choosing $k = \mu$ in Corollary 1.1, we get:

Corollary 1.3. Let $P(z) := a_0 + \sum_{j=\mu}^n a_j z^j$, $1 \leq \mu \leq n-1$, $a_0 \neq 0$ be a polynomial of degree n . If for some real α and β

$$|\arg a_j - \beta| \leq \alpha \leq \frac{\pi}{2}, \quad \mu \leq j \leq n$$

such that

$$|a_\mu| \geq \cdots \geq |a_{n-1}| \geq |a_n|$$

then the number of zeros of $P(z)$ in $|z| \leq \frac{1}{2}$ does not exceed

$$\frac{1}{\log 2} \log \frac{M}{|a_0|},$$

where

$$M = 2|a_0| + |a_\mu|(1 - \sin \alpha + \cos \alpha) + |a_n|(1 - \sin \alpha - \cos \alpha) + 2 \sum_{j=\mu}^n |a_j| \sin \alpha.$$

Taking $\mu = 1$ in Corollary 1.3, we have

Corollary 1.4. Let $P(z) := \sum_{j=0}^n a_j z^j$, $a_0 \neq 0$ be a polynomial of degree n . If for some real α and β

$$|\arg a_j - \beta| \leq \alpha \leq \frac{\pi}{2}, \quad \mu \leq j \leq n$$

such that

$$|a_1| \geq \dots \geq |a_{n-1}| \geq |a_n|$$

then the number of zeros of $P(z)$ in $|z| \leq \frac{1}{2}$ does not exceed

$$\frac{1}{\log 2} \log \frac{M}{|a_0|},$$

where

$$M = 2|a_0| + |a_1|(1 - \sin \alpha + \cos \alpha) + |a_n|(1 - \sin \alpha - \cos \alpha) + 2 \sum_{j=1}^n |a_j| \sin \alpha.$$

Next, we put restriction on the real part of coefficients of a polynomial and proved:

Theorem 2. Let $P(z) := a_0 + \sum_{j=\mu}^n a_j z^j$, $1 \leq \mu \leq n - 1$, $a_0 \neq 0$, be a polynomial of degree n with $Re a_j = \alpha_j$ and $Im a_j = \beta_j$ for $\mu \leq j \leq n$. Suppose that for some $t > 0$ and some k with $\mu \leq k \leq n$ we have

$$t^\mu \alpha_\mu \leq \dots \leq t^{k-1} \alpha_{k-1} \leq t^k \alpha_k \geq t^{k+1} \alpha_{k+1} \geq \dots \geq t^{n-1} \alpha_{n-1} \geq t^n \alpha_n$$

then the number of zeros of $P(z)$ in $|z| \leq \frac{t}{2}$ does not exceed

$$\frac{1}{\log 2} \log \frac{M}{|a_0|},$$

where

$$M = 2(|\alpha_0| + |\beta_0|)t + (|\alpha_\mu| - \alpha_\mu)t^{\mu+1} + 2\alpha_k t^{k+1} + (|\alpha_n| - \alpha_n)t^{n+1} + 2 \sum_{j=\mu}^n |\beta_j| t^{j+1}.$$

For $t = 1$ in Theorem 2, we obtain

Corollary 2.1. Let $P(z) := a_0 + \sum_{j=\mu}^n a_j z^j$, $1 \leq \mu \leq n - 1$, $a_0 \neq 0$, be a polynomial of degree n with $Re a_j = \alpha_j$ and $Im a_j = \beta_j$ for $\mu \leq j \leq n$. Suppose that for some k with $\mu \leq k \leq n$ we have

$$\alpha_\mu \leq \dots \leq \alpha_{k-1} \leq \alpha_k \geq \alpha_{k+1} \geq \dots \geq \alpha_{n-1} \geq \alpha_n$$

then the number of zeros of $P(z)$ in $|z| \leq \frac{1}{2}$ does not exceed

$$\frac{1}{\log 2} \log \frac{M}{|a_0|},$$

where

$$M = 2(|\alpha_0| + |\beta_0|) + (|\alpha_\mu| - \alpha_\mu) + 2\alpha_k + (|\alpha_n| - \alpha_n) + 2 \sum_{j=\mu}^n |\beta_j|.$$

For $k = n$ in Corollary 2.1, we get:

Corollary 2.2. Let $P(z) := a_0 + \sum_{j=\mu}^n a_j z^j$, $1 \leq \mu \leq n-1$, $a_0 \neq 0$, be a polynomial of degree n with $\operatorname{Re} a_j = \alpha_j$ and $\operatorname{Im} a_j = \beta_j$ for $\mu \leq j \leq n$ such that

$$\alpha_\mu \leq \cdots \leq \alpha_{n-1} \leq \alpha_n$$

then the number of zeros of $P(z)$ in $|z| \leq \frac{1}{2}$ does not exceed

$$\frac{1}{\log 2} \log \frac{M}{|a_0|},$$

where

$$M = 2(|\alpha_0| + |\beta_0|) + (|\alpha_\mu| - \alpha_\mu) + 2 \sum_{j=\mu}^n |\beta_j|.$$

For $k = \mu$, in Corollary 2.1, we get:

Corollary 2.3. Let $P(z) := a_0 + \sum_{j=\mu}^n a_j z^j$, $1 \leq \mu \leq n-1$, $a_0 \neq 0$, be a polynomial of degree n with $\operatorname{Re} a_j = \alpha_j$ and $\operatorname{Im} a_j = \beta_j$ for $\mu \leq j \leq n$ such that

$$\alpha_\mu \geq \cdots \geq \alpha_{n-1} \geq \alpha_n$$

then the number of zeros of $P(z)$ in $|z| \leq \frac{1}{2}$ does not exceed

$$\frac{1}{\log 2} \log \frac{M}{|a_0|},$$

where

$$M = 2(|\alpha_0| + |\beta_0|) + (|\alpha_\mu| + \alpha_\mu) + (|\alpha_\mu| - \alpha_\mu) + 2 \sum_{j=\mu}^n |\beta_j|.$$

For $\beta_j = 0$, $1 \leq j \leq n$ in Theorem 2, we have the following:

Corollary 2.4. Let $P(z) := a_0 + \sum_{j=\mu}^n a_j z^j$, $\mu \leq n-1$, where $a_0 \neq 0$. Suppose that for some $t > 0$ and some k we have

$$t^\mu a_\mu \leq \cdots \leq t^{k-1} a_{k-1} \leq t^k a_k \geq t^{k+1} a_{k+1} \geq \cdots \geq t^{n-1} a_{n-1} \geq t^n a_n$$

then the number of zeros of $P(z)$ in $|z| \leq \frac{t}{2}$ does not exceed

$$\frac{1}{\log 2} \log \frac{M}{|a_0|},$$

where

$$M = 2|a_0|t + (|a_\mu| - a_\mu)t^{\mu+1} + 2a_k t^{k+1} + (|a_n| - a_n)t^{n+1}.$$

Finally, we prove the following result:

Theorem 3. Let $P(z) := a_0 + \sum_{j=\mu}^n a_j z^j$, $1 \leq \mu \leq n-1$, where $a_0 \neq 0$, $Re a_j = \alpha_j$ and $Im a_j = \beta_j$ for $\mu \leq j \leq n$. Suppose that for some $t > 0$ and some k with $\mu \leq k \leq n$ we have

$$t^\mu \alpha_\mu \leq \dots \leq t^{k-1} \alpha_{k-1} \leq t^k \alpha_k \geq t^{k+1} \alpha_{k+1} \geq \dots \geq t^{n-1} \alpha_{n-1} \geq t^n \alpha_n$$

and for some $\mu \leq l \leq n$ we have

$$t^\mu \beta_\mu \leq \dots \leq t^{l-1} \beta_{l-1} \leq t^l \beta_l \geq t^{l+1} \beta_{l+1} \geq \dots \geq t^{n-1} \beta_{n-1} \geq t^n \beta_n$$

then the number of zeros of $P(z)$ in $|z| \leq \frac{t}{2}$ does not exceed

$$\frac{1}{\log 2} \log \frac{M}{|a_0|},$$

where

$$M = 2(|\alpha_0| + |\beta_0|)t + (|\alpha_\mu| - \alpha_\mu + |\beta_\mu| - \beta_\mu)t^{\mu+1} + 2(\alpha_k t^{k+1} + \beta_l t^{l+1})t^{n+1} + (|\alpha_n| - \alpha_n + |\beta_n| - \beta_n)t^{n+1}.$$

If we take $t = 1$, in Theorem 3 we obtain:

Corollary 3.1. Let $P(z) = a_0 + \sum_{j=\mu}^n a_j z^j$, $1 \leq \mu \leq n-1$, where $a_0 \neq 0$, $Re a_j = \alpha_j$ and $Im a_j = \beta_j$ for $\mu \leq j \leq n$. Suppose that for some k with $\mu \leq k \leq n$ we have

$$\alpha_\mu \leq \dots \leq \alpha_{k-1} \leq \alpha_k \geq \alpha_{k+1} \geq \dots \geq \alpha_{n-1} \geq \alpha_n$$

and for some $\mu \leq l \leq n$ we have

$$\beta_\mu \leq \dots \leq \beta_{l-1} \leq \beta_l \geq \beta_{l+1} \geq \dots \geq \beta_{n-1} \geq \beta_n.$$

Then the number of zeros of $P(z)$ in $|z| \leq \frac{1}{2}$ does not exceed

$$\frac{1}{\log 2} \log \frac{M}{|a_0|},$$

where

$$M = 2(|\alpha_0| + |\beta_0|) + (|\alpha_\mu| - \alpha_\mu + |\beta_\mu| - \beta_\mu) + 2(\alpha_k + \beta_l) + (|\alpha_n| - \alpha_n + |\beta_n| - \beta_n).$$

For $k = l = n$ in Corollary 3.1, we get the following:

Corollary 3.2. Let $P(z) = a_0 + \sum_{j=\mu}^n a_j z^j$, $1 \leq \mu \leq n-1$, where $a_0 \neq 0$, $Re a_j = \alpha_j$ and $Im a_j = \beta_j$ for $\mu \leq j \leq n$ such that

$$\alpha_\mu \leq \dots \leq \alpha_{n-1} \leq \alpha_n$$

and

$$\beta_\mu \leq \cdots \leq \beta_{n-1} \leq \beta_n$$

then the number of zeros of $P(z)$ in $|z| \leq \frac{1}{2}$ does not exceed

$$\frac{1}{\log 2} \log \frac{M}{|a_0|},$$

where

$$M = 2(|\alpha_0| + |\beta_0|) + (|\alpha_\mu| - \alpha_\mu + |\beta_\mu| - \beta_\mu) + (|\alpha_n| + \alpha_n + |\beta_n| + \beta_n).$$

In Corollary 3.1, if we choose $k = l = \mu$ we get:

Corollary 3.3. Let $P(z) = a_0 + \sum_{j=\mu}^n a_j z^j$, $1 \leq \mu \leq n-1$, where $a_0 \neq 0$, $\operatorname{Re} a_j = \alpha_j$ and $\operatorname{Im} a_j = \beta_j$ for $\mu \leq j \leq n$ such that

$$\alpha_\mu \geq \cdots \geq \alpha_{n-1} \leq \alpha_n$$

and

$$\beta_\mu \geq \cdots \geq \beta_{n-1} \geq \beta_n$$

then the number of zeros of $P(z)$ in $|z| \leq \frac{1}{2}$ does not exceed

$$\frac{1}{\log 2} \log \frac{M}{|a_0|},$$

where

$$M = 2(|\alpha_0| + |\beta_0|) + (|\alpha_\mu| + \alpha_\mu + |\beta_\mu| + \beta_\mu) + (|\alpha_n| - \alpha_n + |\beta_n| - \beta_n).$$

2. Lemma

For the proof of some these results we need the following lemma which is due to Govil and Rahman [4].

Lemma 2.1. For any two complex numbers b_0 and b_1 such that $|b_0| \geq |b_1|$ and

$$|\arg b_j - \beta| \leq \alpha \leq \frac{\pi}{2}, \quad j = 0, 1$$

for some β , then

$$|b_0 - b_1| \leq (|b_0| - |b_1|) \cos \alpha + (|b_0| + |b_1|) \sin \alpha.$$

An application of the Maximum modulus theorem shown in (p.171, [14]) we have the following interesting result:

Lemma 2.2. *Let $f(z)$ be regular and $|f(z)| \leq M$, in the circle $|z| \leq R$ and suppose that $f(0) \neq 0$, then the number of zeros of $f(z)$ in the circle $|z| \leq \frac{1}{2}R$ does not exceed $\frac{1}{\log 2} \log \left[\frac{M}{|f(0)|} \right]$.*

3. Proofs of Theorems

Proof of Theorem 1. Consider the polynomial

$$\begin{aligned} F(z) &= (t - z)P(z) \\ &= (t - z) \left(a_0 + \sum_{j=\mu}^n a_j z^j \right) \\ &= a_0 t + \sum_{j=\mu}^n a_j t z^j - a_0 z - \sum_{j=\mu}^n a_j z^{j+1} \\ &= a_0(t - z) + \sum_{j=\mu}^n a_j t z^j - \sum_{j=\mu+1}^{n+1} a_{j-1} z^j \\ &= a_0(t - z) + a_\mu t z^\mu + \sum_{j=\mu+1}^n (a_j t - a_{j-1}) z^j - a_n z^{n+1}. \end{aligned}$$

For $|z| = t$, we have

$$\begin{aligned} |F(z)| &\leq 2|a_0|t + |a_\mu|t^{\mu+1} + \sum_{j=\mu+1}^n |a_j t - a_{j-1}|t^j + |a_n|t^{n+1} \\ &= 2|a_0|t + |a_\mu|t^{\mu+1} + \sum_{j=\mu+1}^k |a_j t - a_{j-1}|t^j + \sum_{j=k+1}^n |a_{j-1} - a_j t|t^j + |a_n|t^{n+1}. \end{aligned}$$

Using Lemma 2.1 with $b_0 = a_j t$ and $b_1 = a_{j-1}$ when $1 \leq j \leq k$ and with $b_0 = a_{j-1}$ and $b_1 = a_j t$ when $k + 1 \leq j \leq n$,

$$\begin{aligned} |F(z)| &\leq 2|a_0|t + |a_\mu|t^{\mu+1} + \sum_{j=\mu+1}^k \{(|a_j|t - |a_{j-1}|) \cos \alpha + (|a_j|t + |a_{j-1}|) \sin \alpha\} t^j \\ &\quad + \sum_{j=k+1}^n \{(|a_{j-1}| - |a_j|t) \cos \alpha + (|a_j|t + |a_{j-1}|) \sin \alpha\} t^j + |a_n|t^{n+1} \end{aligned}$$

$$\begin{aligned}
&= 2|a_0|t + |a_\mu|t^{\mu+1} + \sum_{j=\mu+1}^k |a_j|t^{j+1} \cos \alpha - \sum_{j=\mu+1}^k |a_{j-1}|t^j \cos \alpha + \sum_{j=\mu+1}^k |a_j|t^{j+1} \sin \alpha \\
&\quad + \sum_{j=\mu+1}^k |a_{j-1}|t^j \sin \alpha + \sum_{j=k+1}^n |a_{j-1}|t^j \cos \alpha - \sum_{j=k+1}^n |a_j|t^{j+1} \cos \alpha \\
&\quad + \sum_{j=k+1}^n |a_{j-1}|t^j \sin \alpha + \sum_{j=k+1}^n |a_j|t^{j+1} \sin \alpha + |a_n|t^{n+1} \\
&= 2|a_0|t + |a_\mu|t^{\mu+1} - |a_\mu|t^{\mu+1} \cos \alpha + |a_k|t^{k+1} \cos \alpha + |a_\mu|t^{\mu+1} \sin \alpha + |a_k|t^{k+1} \sin \alpha \\
&\quad + 2 \sum_{j=\mu+1}^{k-1} |a_j|t^{j+1} \sin \alpha + |a_k|t^{k+1} \cos \alpha - |a_n|t^{n+1} \cos \alpha + |a_k|t^{k+1} \sin \alpha \\
&\quad + |a_n|t^{n+1} \sin \alpha + 2 \sum_{j=k+1}^{n-1} |a_j|t^{j+1} \sin \alpha + |a_n|t^{n+1} \\
&= 2|a_0|t + |a_\mu|t^{\mu+1} + |a_\mu|t^{\mu+1}(\sin \alpha - \cos \alpha) + 2 \sum_{j=\mu+1}^{n-1} |a_j|t^{j+1} \sin \alpha + 2|a_k|t^{k+1} \cos \alpha \\
&\quad + (\sin \alpha - \cos \alpha + 1)|a_n|t^{n+1} \\
&= 2|a_0|t + |a_\mu|t^{\mu+1}(1 - \sin \alpha - \cos \alpha) + 2|a_k|t^{k+1} \cos \alpha + |a_n|t^{n+1}(1 - \sin \alpha - \cos \alpha) \\
&\quad + 2 \sum_{j=\mu}^n |a_j|t^{j+1} \sin \alpha \\
&= M(\text{say}).
\end{aligned}$$

Now $F(z)$ is analytic in $|z| \leq t$ and $F(z) \leq M$ for $|z| = t$. Applying Lemma 2.2 to the polynomial $F(z)$, we get the number of zeros of $F(z)$ in $|z| \leq \frac{t}{2}$ does not exceed

$$\frac{1}{\log 2} \log \frac{M}{|f(0)|}.$$

Thus, the number of zeros of $F(z)$ in $|z| \leq \frac{t}{2}$ does not exceed

$$\frac{1}{\log 2} \log \left\{ \frac{2|a_0|t + |a_\mu|t^{\mu+1}(1 - \sin \alpha - \cos \alpha) + 2|a_k|t^{k+1} \cos \alpha}{|a_0|} + \frac{|a_n|t^{n+1}(1 - \sin \alpha - \cos \alpha) + 2 \sum_{j=\mu}^n |a_j|t^{j+1} \sin \alpha}{|a_0|} \right\}.$$

As the number of zeros of $P(z)$ in $|z| \leq \frac{1}{2}$ is also equal to the number of zeros $F(z)$ the theorem follows. \square

Proof of Theorem 2. Consider the polynomial

$$\begin{aligned}
 F(z) &= (t - z)P(z) \\
 &= (t - z) \left(a_0 + \sum_{j=\mu}^n a_j z^j \right) \\
 &= a_0 t + \sum_{j=\mu}^n a_j t z^j - a_0 z - \sum_{j=\mu}^n a_j z^{j+1} \\
 &= a_0(t - z) + \sum_{j=\mu}^n a_j t z^j - \sum_{j=\mu+1}^{n+1} a_{j-1} z^j
 \end{aligned}$$

and therefore

$$\begin{aligned}
 F(z) &= (\alpha_0 + i\beta_0)(t - z) + (\alpha_\mu + i\beta_\mu)t z^\mu + \sum_{j=\mu+1}^n (\alpha_j t - \alpha_{j-1})z^j \\
 &\quad + i \sum_{j=\mu+1}^n (\beta_j t - \beta_{j-1})z^j - (\alpha_n + i\beta_n)z^{n+1}.
 \end{aligned}$$

For $|z| = t$, we have

$$\begin{aligned}
 |F(z)| &\leq 2(|\alpha_0| + |\beta_0|)t + (|\alpha_\mu| + |\beta_\mu|)t^{\mu+1} + \sum_{j=\mu+1}^n |\alpha_j t - \alpha_{j-1}|t^j \\
 &\quad + \sum_{j=\mu+1}^n (|\beta_j|t + |\beta_{j-1}|)t^j + (|\alpha_n| + |\beta_n|)t^{n+1} \\
 &= 2(|\alpha_0| + |\beta_0|)t + (|\alpha_\mu| + |\beta_\mu|)t^{\mu+1} + \sum_{j=\mu+1}^k (\alpha_j t - \alpha_{j-1})t^j \\
 &\quad + \sum_{j=k+1}^n (\alpha_{j-1} - \alpha_j t)t^j + |\beta_\mu|t^{\mu+1} + 2 \sum_{j=\mu+1}^{n-1} |\beta_j|t^{j+1} + |\beta_n|t^{n+1} \\
 + (|\alpha_n| + |\beta_n|)t^{n+1} &= 2(|\alpha_0| + |\beta_0|)t + (|\alpha_\mu| + |\beta_\mu|)t^{\mu+1} - \alpha_\mu t^{\mu+1} + 2\alpha_k t^{k+1} \\
 - \alpha_n t^{n+1} + |\beta_\mu|t^{\mu+1} &+ 2 \sum_{j=\mu+1}^n |\beta_j|t^{j+1} + |\alpha_n|t^{n+1} = 2(|\alpha_0| + |\beta_0|)t \\
 + (|\alpha_\mu| - \alpha_\mu)t^{\mu+1} + 2\alpha_k t^{k+1} &+ (|\alpha_n| - \alpha_n)t^{n+1} + 2 \sum_{j=\mu}^n |\beta_j|t^{j+1} = M.
 \end{aligned}$$

Proceedings on the same lines of the proof of Theorem 1, the proof of this result follows.

Proof of Theorem 3. Consider the polynomial

$$F(z) = (t - z)P(z) = a_0(t - z) + a_\mu tz^\mu + \sum_{j=\mu+1}^n (a_j t - a_{j-1})z^j - a_n z^{n+1},$$

and so

$$\begin{aligned} F(z) &= (\alpha_0 + i\beta_0)(t - z) + (\alpha_\mu + i\beta_\mu)tz^\mu \\ &+ \sum_{j=\mu+1}^n ((\alpha_j + i\beta_j)t - (\alpha_{j-1} + i\beta_{j-1}))z^j - (\alpha_n + i\beta_n)z^{n+1} \\ &= (\alpha_0 + i\beta_0)(t - z) + (\alpha_\mu + i\beta_\mu)tz^\mu + \sum_{j=\mu+1}^n (\alpha_j t - \alpha_{j-1})z^j \\ &+ i \sum_{j=\mu+1}^n (\beta_j t - \beta_{j-1})z^j - (\alpha_n + i\beta_n)z^{n+1}. \end{aligned}$$

For $|z| = t$, we have

$$\begin{aligned} |F(z)| &\leq 2(|\alpha_0| + |\beta_0|)t + (|\alpha_\mu| + |\beta_\mu|)t^{\mu+1} + \sum_{j=\mu+1}^n |\alpha_j t - \alpha_{j-1}|t^j \\ &+ \sum_{j=\mu+1}^n |\beta_j t - \beta_{j-1}|t^j + (|\alpha_n| + |\beta_n|)t^{n+1} = 2(|\alpha_0| + |\beta_0|)t \\ &+ (|\alpha_\mu| + |\beta_\mu|)t^{\mu+1} + \sum_{j=\mu+1}^k (\alpha_j t - \alpha_{j-1})t^j + \sum_{j=k+1}^n (\alpha_{j-1} - \alpha_j t)t^j \\ &+ \sum_{j=\mu+1}^l (\beta_j t - \beta_{j-1})t^j + \sum_{j=l+1}^n (\beta_{j-1} - \beta_j t)t^j + (|\alpha_n| + |\beta_n|)t^{n+1} \\ &= 2(|\alpha_0| + |\beta_0|)t + (|\alpha_\mu| + |\beta_\mu|)t^{\mu+1} - \alpha_\mu t^{\mu+1} + 2\alpha_k t^{k+1} \\ &- \alpha_n t^{n+1} - \beta_\mu t^{\mu+1} + 2\beta_l t^{l+1} - \beta_n t^{n+1} + (|\alpha_n| + |\beta_n|)t^{n+1} \\ &= 2(|\alpha_0| + |\beta_0|)t + (|\alpha_\mu| - \alpha_\mu + |\beta_\mu| - \beta_\mu)t^{\mu+1} \\ &+ 2(\alpha_k t^{k+1} + \beta_l t^{l+1}) + (|\alpha_n| - \alpha_n + |\beta_n| - \beta_n)t^{n+1} = M. \end{aligned}$$

The result now follows as in the proof of Theorem 1. \square

References

- [1] I. Ahmad, T. Rasool, A. Liman, *Zeros of certain polynomials and analytic functions with restricted coefficients*, Journal of Classical Analysis 2 (2014) 149–157.
- [2] T. Chan, M. Malik, *On the Erdős-Lax theorem*, Proceeding of Indian Academy of Sciences 92 (1983) 191–193.
- [3] K.K. Dewan, *Extremal properties and coefficient estimates for polynomials with restricted zeros and on location of zeros of polynomials*, Ph.D Thesis, IIT Delhi, 1980.
- [4] N.K. Govil, Q.I. Rahman, *On the Eneström-Keakeya theorem*, Tohoku Math. J. 20 (1968) 126–136.
- [5] E. Landau, *Über den Picardschen Satz*, Vierteljahrsschrift Naturforsch. Gesellschaft Zürcher & Furrer 51 (1906) 252–318.
- [6] E. Landau, *Sur quelques generalisations du theoreme de M. Picard*, Ann. Ecole Norm. 24 (3) (1907) 179–201.
- [7] A. Liman, W.M. Shah, T. Rasool, *Some new generalizations of Eneström-Keakeya theorem*, Nonlinear Functional Analysis and Applications 17 (2) (2012) 235–248.
- [8] A. Liman, I.A. Faiq, *Bounds for the zeros of polynomial*, International Journal of Modern Mathematical Sciences 13 (1) (2015) 12–16.
- [9] M. Marden, *Geometry of Polynomials*, Math. Surveys No. 3, Amer. Math. Soc. Providence R. I., 1949.
- [10] G.V. Milovanovic, D.S. Mitrinovic, Th. M. Rassias, *Topics in Polynomials, Extermal Problems, Inequalities, Zeros*, World Scientific, Singapore, 1994.
- [11] Q.G. Mohammad, *On the zeros of the polynomials*, Amer. Math. Monthly 72 (6) (1965) 631–633.
- [12] Q.I. Rahman, S. Schmeisser, *Analytic Theory of Polynomials*, Oxford University press, New York, 2002.
- [13] W.M. Shah, A. Liman, *On Eneström-Keakeya theorem and related analytic functions*, Proc. Indian Acad. Sci.(Math Sci.) 117 (3) (2007) 359–370.
- [14] E.C. Titchmarsh, *The Theory of Functions*, 2nded. Oxford Univ. Press, London, 1939.

DOI: 10.7862/rf.2018.13

Idrees Qasim

email: idreesf3@gmail.com
ORCID: 0000-0001-5838-6522
Department of Mathematics
National Institute of Technology
Srinagar - 190006
INDIA

Tawheeda Rasool

email: tawheedrasool@gmail.com
Department of Mathematics
National Institute of Technology
Srinagar - 190006
INDIA

Abdul Liman

email: abliman22@rediffmail.com
Department of Mathematics
National Institute of Technology
Srinagar - 190006
INDIA

Received 12.08.2017

Accepted 14.08.2018