

dr n. tech. Andrzej Antoni CZAJKOWSKI^{a,b}, dr inż. Wojciech Kazimierz OLESZAK^b, mgr inż., naucz. dypl. Sławomir Wawrzyniec SNASTIN^a, naucz. dypl. Lech Adam REWKOWSKI^c

^a Higher School of Technology and Economics in Szczecin, Faculty of Automotive Systems
Wyższa Szkoła Techniczno-Ekonomiczna w Szczecinie, Wydział Systemów Automotywowych

^b Higher School of Humanities of Common Knowledge Society in Szczecin
Wyższa Szkoła Humanistyczna Towarzystwa Wiedzy Powszechniej w Szczecinie

^c 18 High School in the Complex School No. 5 in Szczecin
18 Liceum Ogólnokształcące w Zespole Szkół Nr 5 w Szczecinie

ANALYTICAL AND NUMERICAL SOLVING OF RIGHT TRIANGLES WITH GIVEN A SUM OF TWO SIDES LENGTH AND THE ACUTE ANGLE

Abstract

Introduction and aims: The paper shows the analytical models of solving right triangles with appropriate discussion. For right triangles have been discussed six cases taking into account the acute angle and the sum of two side length in the right triangle. The main aim of this paper is not only to create some analytical algorithms for solving right triangle, but also their implementation in programs *MS-Excel*, *MathCAD* and *Mathematica*.

Material and methods: Elaboration of six analytical cases of solving right triangles has been made on the basis of the relevant trigonometric properties occurring in a right triangle. In the paper have been used some analytical and numerical methods by using *MS-Excel*, *MathCAD* and *Mathematica* programs.

Results: As some results have been obtained numerical algorithms in the programs *MS-Excel*, *MathCAD* and *Mathematica* for six analytical cases of solving right triangles taking into account the acute angle and the sum of two side length in the right triangle.

Conclusion: Created numerical algorithms of solving the right triangles in the programs *MS-Excel*, *MathCAD* and *Mathematica* allow for faster significant performance calculations than the traditional way of using logarithms and logarithmic tables.

Keywords: Trigonometry, right triangle, acute angle, sum of two sides length of a triangle, numerical algorithms, *MS-Excel*, *MathCAD*, *Mathematica*.

(Received: 01.05.2018; Revised: 13.05.2018; Accepted: 20.05.2018)

ANALITYCZNO-NUMERYCZNE ROZWIĄZYWANIE TRÓJKĄTÓW PROSTOKĄTNYCH GDZIE DANA JEST SUMA DŁUGOŚCI DWÓCH BOKÓW I KĄT OSTRY

Streszczenie

Wstęp i cele: W pracy pokazano analityczne modele rozwiązywania trójkątów prostokątnych wraz z odpowiednią dyskusją. Dla trójkątów prostokątnych omówiono sześć przypadków z uwzględnieniem kąta ostrego oraz sumy długości dwóch boków trójkąta. Głównym celem jest pracy jest nie tylko utworzenie algorytmów analitycznych rozwiązywania takich trójkątów lecz również ich implementacja w programach *MS-Excel*, *MathCAD* i *Mathematica*.

Materiał i metody: Opracowanie sześciu analitycznych przypadków rozwiązywania trójkątów prostokątnych wykonyano opierając się odpowiednich własnościach trygonometrycznych występujących w trójkącie prostokątnym. Zastosowano metodę analityczną i numeryczną wykorzystując programy *MS-Excel*, *MathCAD* i *Mathematica*.

Wyniki: Otrzymano algorytmy numeryczne w programach *MS-Excel*, *MathCAD* i *Mathematica* dla sześciu analitycznych przypadków rozwiązywania trójkątów prostokątnych z uwzględnieniem kąta ostrego oraz sumy długości dwóch boków trójkąta.

Wniosek: Utworzono algorytmy numeryczne rozwiązywania trójkątów prostokątnych w programach *MS-Excel*, *MathCAD* oraz *Mathematica*, pozwalają na znaczne szybsze wykonanie obliczeń niż drogą tradycyjną z użyciem logarytmów i tablic logarytmicznych.

Słowa kluczowe: Trygonometria, trójkąt prostokątny, kąt ostry, suma długości dwóch boków trójkąta, algorytmy numeryczne, *MS-Excel*, *MathCAD*, *Mathematica*.

(Otrzymano: 01.05.2018; Zrecenzowano: 13.05.2018; Zaakceptowano: 20.05.2018)

1. Introduction to trigonometry

1.1. Definitions of trigonometric functions

Let be given the right triangle ABC (angle ACB is the right angle) (Fig. I).




Fig. I. The right triangle with legs a and b and hypotenuse c
Source: Elaboration of the Authors

where a, b, c mean respectively the length of shorter leg, longer leg and hypotenuse. The sides a, b and c satisfy the condition:

$$a^2 + b^2 = c^2 \quad (1)$$

and acute angles α and β satisfying the following condition [4], [6], [7]:

$$\alpha + \beta = \frac{\pi}{2}. \quad (2)$$

Definition 1.

The sine of acute angle is the ratio of the length of leg opposite this angle to the length of hypotenuse, which express the formulae [4], [6], [7]:

$$\sin(\alpha) = \frac{a}{c}, \quad (3)$$

$$\sin(\beta) = \frac{b}{c}. \quad (4)$$

Definition 2.

The cosine of acute angle is the ratio of the length of leg adjacent this angle to the length of hypotenuse, which express the formulae [4], [6], [7]:

$$\cos(\alpha) = \frac{b}{c}, \quad (5)$$

$$\cos(\beta) = \frac{a}{c}. \quad (6)$$

Definition 3.

The tangent of acute angle is the ratio of the length of leg opposite this angle to the length of leg adjacent to that angle, which express the formulae [4], [6], [7]:

$$\operatorname{tg}(\alpha) = \frac{a}{b}, \quad (7)$$

$$\operatorname{tg}(\beta) = \frac{b}{a}. \quad (8)$$

Definition 4.

The cotangent of acute angle is the ratio of the length of leg adjacent this angle to the length of leg opposite to that angle, which express the formulae [4], [6], [7]:

$$\operatorname{ctg}(\alpha) = \frac{b}{a}, \quad (9)$$

$$\operatorname{ctg}(\beta) = \frac{a}{b}. \quad (10)$$

Theorem 1. (Basic relationships in the triangle)

In any length of each side of a triangle is less than the sum of the length of the other side and higher than the absolute value of the difference in length of the sides [10], [11], [13]-[16]:

$$|b - c| < a < b + c, \quad (11)$$

$$|a - c| < b < a + c, \quad (12)$$

$$|a - b| < c < a + b. \quad (13)$$

1.2. Solving right triangles with given the length of two legs and acute angle of triangle

For the right triangle ABC shown in figure I, there are six cases of solving which are given below (Tab. 1).

Tab. 1. The discussion of solutions for the right triangles

Case:	Data:	Solution:			
1	$a+c=k$ α	$\beta = \frac{\pi}{2} - \alpha$	$a = \frac{k \cdot \sin(\alpha)}{1 + \sin(\alpha)}$	$b = \frac{k \cdot \cos(\alpha)}{1 + \sin(\alpha)}$	$c = \frac{k}{1 + \sin(\alpha)}$
2	$a+c=k$ β	$\alpha = \frac{\pi}{2} - \beta$	$a = \frac{k \cdot \cos(\beta)}{1 + \cos(\beta)}$	$b = \frac{k \cdot \sin(\beta)}{1 + \cos(\beta)}$	$c = \frac{k}{1 + \cos(\beta)}$
3	$b+c=p$ α	$\beta = \frac{\pi}{2} - \alpha$	$a = \frac{p \cdot \sin(\alpha)}{1 + \cos(\alpha)} = p \cdot \operatorname{tg}\left(\frac{\alpha}{2}\right)$	$a = \frac{p \cdot \cos(\alpha)}{1 + \cos(\alpha)}$	$a = \frac{p}{1 + \cos(\alpha)}$
4	$b+c=p$ β	$\alpha = \frac{\pi}{2} - \beta$	$a = \frac{p \cdot \cos(\beta)}{1 + \cos(\beta)}$	$a = \frac{p \cdot \sin(\beta)}{1 + \cos(\beta)}$	$a = \frac{p}{1 + \sin(\beta)}$
5	$a+b=q$ α	$\beta = \frac{\pi}{2} - \alpha$	$a = \frac{q \cdot \operatorname{tg}(\alpha)}{1 + \operatorname{tg}(\alpha)}$	$b = \frac{q}{1 + \operatorname{tg}(\alpha)}$	$c = \frac{q \cdot \sec(\alpha)}{1 + \operatorname{tg}(\alpha)} \text{ } ^1$
6	$a+b=q$ β	$\alpha = \frac{\pi}{2} - \beta$	$a = \frac{q \cdot \operatorname{ctg}(\beta)}{1 + \operatorname{ctg}(\beta)}$	$b = \frac{q}{1 + \operatorname{ctg}(\beta)}$	$c = \frac{q \cdot \operatorname{cosec}(\beta)}{1 + \operatorname{ctg}(\beta)} \text{ } ^2$

Source: Elaborated by the Authors

¹ Note: $\sec(\alpha) \equiv 1/\cos(\alpha)$.

² Note: $\operatorname{cosec}(\alpha) \equiv 1/\sin(\alpha)$.

2. Solving of right triangles

2.1. Case 1: Sum of two sides length $a+c = k$ and a measure of the angle α .

2.1.1. Theoretical analysis

➤ Let be the right triangle as shown on the figure 1.




Fig. 1. Right triangle ABC with given a sum of the sides length $a+c=k$ and a measure of the angle α
Source: Elaboration of the Authors

➤ *Data in $\triangle ABC$:* Sum of the sides length $a+c=k$ and a measure of the angle α .

➤ *Unknown in $\triangle ABC$:* Length of the sides a, b, c and a measure of the angle β .

➤ *Solution:*

◆ From some properties of acute angles in a right triangle we obtain:

$$\beta = \frac{\pi}{2} - \alpha. \quad (14)$$

◆ From the sum of the triangle sides we have:

$$a + c = k, \quad (15)$$

$$a = k - c. \quad (16)$$

◆ From the formulae (3) and (5) we obtain:

$$a = c \cdot \sin(\alpha), \quad (17)$$

$$b = c \cdot \cos(\alpha). \quad (18)$$

◆ From the formulae (16) and (17) we determine the triangle side c :

$$k - c = c \cdot \sin(\alpha), \quad (19)$$

$$k = c \cdot [1 + \sin(\alpha)], \quad (20)$$

$$c = \frac{k}{1 + \sin(\alpha)}. \quad (21)$$

◆ Substituting (21) into the formulae (17) and (18) we determine the triangle sides a, b :

$$a = \frac{k \cdot \sin(\alpha)}{1 + \sin(\alpha)}, \quad (22)$$

$$b = \frac{k \cdot \cos(\alpha)}{1 + \sin(\alpha)}. \quad (23)$$

➤ *Answer:* $\beta = \frac{\pi}{2} - \alpha$, $a = \frac{k \cdot \sin(\alpha)}{1 + \sin(\alpha)}$, $b = \frac{k \cdot \cos(\alpha)}{1 + \sin(\alpha)}$, $c = \frac{k}{1 + \sin(\alpha)}$.

2.1.2. Numerical algorithms in MS-Excel, MathCAD and Mathematica programs

For numerical analysis we take a sum of the sides $a+c = 43.6398$ and the angle $\alpha = 31^\circ$.

- ◆ Program *MS-Excel 7.0* [3],[8]

<i>Algorithm:</i>	<i>Commentary:</i>
A6=43,6398	<i>Given the sum of the sides $a+c$</i>
A7=31	<i>Given a measure of the angle $\alpha [^\circ]$</i>
A8=A7*PI()/180	<i>Command for the angle $\alpha [rad]$, $\alpha = 0.541052$</i>
A9=90-A7	<i>Command for the angle $\beta [^\circ]$</i>
A10=(A6*SIN(A8))/(1+SIN(A8))	<i>Command for calculation of the side a</i>
A11=(A6*COS(A8))/(1+SIN(A8))	<i>Command for calculation of the side b</i>
A12=A6/(1+SIN(A8))	<i>Command for calculation of the side c</i>
59	<i>Result: measure of the angle $\beta [^\circ]$</i>
14,8354	<i>Result: length of the side a</i>
24,6902	<i>Result: length of the side b</i>
28,8044	<i>Result: length of the side c</i>

- ◆ Program *MathCAD 15* [9],[12]

<i>Algorithm:</i>	<i>Commentary:</i>
k:=43.6398	<i>Given the sum of the sides $a+c$</i>
α0:= 31	<i>Given measure of the angle $\alpha [^\circ]$</i>
α := $\frac{\alpha_0 \cdot \pi}{180} = 0.541052$	<i>Command for the angle $\alpha [rad]$ $\alpha = 0.541052$</i>
β0:=90-α0=59	<i>Command for the angle $\beta [^\circ]$</i>
a := $\frac{k \cdot \sin(\alpha)}{1 + \sin(\alpha)} = 14.835375$	<i>Command and result: length of the side a</i>
b := $\frac{k \cdot \cos(\alpha)}{1 + \sin(\alpha)} = 24.690211$	<i>Command and result: length of the side b</i>
a := $\frac{k}{1 + \sin(\alpha)} = 28.804425$	<i>Command and result: length of the side c</i>

- ◆ Program *Mathematica 7.0* [1],[2],[5]

<i>Algorithm:</i>	<i>Commentary:</i>
k:=43.6398	<i>Given the sum of the sides $a+c$</i>
A0:=31	<i>Given measure of the angle $\alpha [^\circ]$</i>
A:=A0*N[Pi]/180	<i>Command for the angle $\alpha [rad]$ $\alpha = 0.541052$</i>
B=90-A0	<i>Command for the angle $\beta [^\circ]$</i>
a=(k*Sin[A])/(1+Sin[A])	<i>Command: length of the side a</i>
b=(k*Cos[A])/(1+Sin[A])	<i>Command: length of the side b</i>
c=k/(1+Sin[A])	<i>Command: length of the side c</i>
59	<i>Result: measure of the angle $\beta [^\circ]$</i>
14.8354	<i>Result: length of the side a</i>
24.6902	<i>Result: length of the side b</i>
28.8044	<i>Result: length of the side c</i>

Numerical results: $a = 14.8354$, $b = 24.6902$, $c = 28.8044$, $\beta = 59^\circ$.

2.2. Case 2: Sum of two sides length $a+c = k$ and a measure of the angle β

2.2.1. Theoretical analysis

➤ Let be a right triangle as shown on the figure 2.




Fig. 2. Right triangle ABC with given a sum of the sides length $a+c=k$ and a measure of the angle β

Source: Elaboration of the Authors

➤ *Data in $\triangle ABC$:* Sum of the sides length $a+c=k$ and a measure of the angle β .

➤ *Unknown in $\triangle ABC$:* Length of the sides a, b, c and a measure of the angle α .

➤ *Solution:*

◆ From some properties of acute angles in a right triangle we obtain:

$$\alpha = \frac{\pi}{2} - \beta. \quad (24)$$

◆ From the sum of the triangle sides we have:

$$a + c = k, \quad (25)$$

$$a = k - c. \quad (26)$$

◆ From the formulae (6) and (4) we obtain:

$$a = c \cdot \cos(\beta), \quad (27)$$

$$b = c \cdot \sin(\beta). \quad (28)$$

◆ From the formulae (26) and (27) we determine the triangle side c :

$$k - c = c \cdot \cos(\beta), \quad (29)$$

$$k = c \cdot [1 + \cos(\beta)], \quad (30)$$

$$c = \frac{k}{1 + \cos(\beta)}. \quad (31)$$

◆ Substituting (31) into the formulae (27) and (28) we determine the triangle sides a, b :

$$a = \frac{k \cdot \cos(\beta)}{1 + \cos(\beta)}, \quad (32)$$

$$b = \frac{k \cdot \sin(\beta)}{1 + \cos(\beta)}. \quad (33)$$

➤ *Answer:* $\alpha = \frac{\pi}{2} - \beta$, $a = \frac{k \cdot \cos(\beta)}{1 + \cos(\beta)}$, $b = \frac{k \cdot \sin(\beta)}{1 + \cos(\beta)}$, $c = \frac{k}{1 + \cos(\beta)}$.

2.2.2. Numerical algorithms in MS-Excel, MathCAD and Mathematica programs

For numerical analysis we take a sum of the sides $a+c = 43.6398$ and the angle $\beta = 59^\circ$.

- ◆ Program *MS-Excel 7.0* [3],[8]

<i>Algorithm:</i>	<i>Commentary:</i>
A6=43,6398	<i>Given the sum of the sides $a+c$</i>
A7=59	<i>Given a measure of the angle $\beta [^\circ]$</i>
A8=A7*PI()/180	<i>Command for the angle $\beta [\text{rad}]$, $\beta = 1.029744$</i>
A9=90-A7	<i>Command for the angle $\alpha [^\circ]$</i>
A10=(A6*COS(A8))/(1+COS(A8))	<i>Command for calculation of the side a</i>
A11=(A6*SIN(A8))/(1+COS(A8))	<i>Command for calculation of the side b</i>
A12=A6/(1+COS(A8))	<i>Command for calculation of the side c</i>
31	<i>Result: measure of the angle $\alpha [^\circ]$</i>
14,8354	<i>Result: length of the side a</i>
24,6902	<i>Result: length of the side b</i>
28,8044	<i>Result: length of the side c</i>

- ◆ Program *MathCAD 15* [9],[12]

<i>Algorithm:</i>	<i>Commentary:</i>
k:=43.6398	<i>Given the sum of the sides $a+c$</i>
β0:=59	<i>Given measure of the angle $\beta [^\circ]$</i>
β := β0 · π / 180 = 1.029744	<i>Command for the angle $\beta [\text{rad}]$, $\beta = 1.029744$</i>
α0 := 90 - β0 = 31	<i>Command for the angle $\alpha [^\circ]$</i>
a := k · cos(β) / (1 + cos(β)) = 14.835375	<i>Command and result: length of the side a</i>
b := k · sin(β) / (1 + cos(β)) = 24.690211	<i>Command and result: length of the side b</i>
a := k / (1 + cos(β)) = 28.804425	<i>Command and result: length of the side c</i>

- ◆ Program *Mathematica 7.0* [1],[2],[5]

<i>Algorithm:</i>	<i>Commentary:</i>
k := 43.6398	<i>Given the sum of the sides $a+c$</i>
B0 := 59	<i>Given measure of the angle $\beta [^\circ]$</i>
B := A0*N[Pi]/180	<i>Command for the angle $\beta [\text{rad}]$, $\beta = 1.029744$</i>
A = 90 - B0	<i>Command for the angle $\alpha [^\circ]$</i>
a = (k*Cos[A]) / (1+Cos[A])	<i>Command: length of the side a</i>
b = (k*Sin[A]) / (1+Cos[A])	<i>Command: length of the side b</i>
c = k / (1+Cos[A])	<i>Command: length of the side c</i>
31	<i>Result: measure of the angle $\alpha [^\circ]$</i>
14.8354	<i>Result: length of the side a</i>
24.6902	<i>Result: length of the side b</i>
28.8044	<i>Result: length of the side c</i>

Numerical results: $a = 14.8354$, $b = 24.6902$, $c = 28.8044$, $\alpha = 31^\circ$.

2.3. Case 3: Sum of two sides length $b+c=p$ and a measure of the angle α .

2.3.1. Theoretical analysis

➤ Let be a right triangle as shown on the figure 3.




Fig. 3. Right triangle ABC with given a sum of the sides length $b+c=p$ and a measure of the angle α

Source: Elaboration of the Authors

➤ *Data in $\triangle ABC$:* Sum of the sides length $b+c=p$ and a measure of the angle α .

➤ *Unknown in $\triangle ABC$:* Length of the sides a, b, c and a measure of the angle β .

➤ *Solution:*

◆ From some properties of acute angles in a right triangle we obtain:

$$\beta = \frac{\pi}{2} - \alpha. \quad (34)$$

◆ From the sum of the triangle sides we have:

$$b + c = p, \quad (35)$$

$$b = p - c. \quad (36)$$

◆ From the formulae (3) and (5) we obtain:

$$a = c \cdot \sin(\alpha), \quad (37)$$

$$b = c \cdot \cos(\alpha). \quad (38)$$

◆ From the formulae (36) and (38) we determine the triangle side c :

$$p - c = c \cdot \cos(\alpha), \quad (39)$$

$$p = c \cdot [1 + \cos(\alpha)], \quad (40)$$

$$c = \frac{p}{1 + \cos(\alpha)}. \quad (41)$$

◆ Substituting (41) into the formulae (37) and (38) we determine the triangle sides a, b :

$$a = \frac{p \cdot \sin(\alpha)}{1 + \cos(\alpha)} \equiv p \cdot \operatorname{tg}\left(\frac{\alpha}{2}\right), \quad (42)$$

$$b = \frac{p \cdot \cos(\alpha)}{1 + \cos(\alpha)}. \quad (43)$$

➤ *Answer:* $\beta = 90^\circ - \alpha$, $a = \frac{p \cdot \sin(\alpha)}{1 + \cos(\alpha)} = p \cdot \operatorname{tg}\left(\frac{\alpha}{2}\right)$, $b = \frac{p \cdot \cos(\alpha)}{1 + \cos(\alpha)}$, $c = \frac{p}{1 + \cos(\alpha)}$.

2.3.2. Numerical algorithms in MS-Excel, MathCAD and Mathematica programs

For numerical analysis we take a sum of the sides $b+c = 53.4946$ and the angle $\alpha = 31^\circ$.

◆ Program *MS-Excel 7.0* [3],[8]

<i>Algorithm:</i>	<i>Commentary:</i>
A6=53,4946	<i>Given the sum of the sides $b+c$</i>
A7=31	<i>Given a measure of the angle $\alpha [^\circ]$</i>
A8=A7*PI()/180	<i>Command for the angle $\alpha [rad]$, $\alpha = 0.541052$</i>
A9=90-A7	<i>Command for the angle $\beta [^\circ]$</i>
A10=(A6*SIN(A8))/(1+COS(A8))	<i>Command for calculation of the side a</i>
A11=(A6*COS(A8))/(1+COS(A8))	<i>Command for calculation of the side b</i>
A12=A6/(1+COS(A8))	<i>Command for calculation of the side c</i>
59	<i>Result: measure of the angle $\beta [^\circ]$</i>
14,8354	<i>Result: length of the side a</i>
24,6902	<i>Result: length of the side b</i>
28,8044	<i>Result: length of the side c</i>

◆ Program *MathCAD 15* [9],[12]

<i>Algorithm:</i>	<i>Commentary:</i>
p:=53.4946	<i>Given the sum of the sides $b+c$</i>
α0:=31	<i>Given measure of the angle $\alpha [^\circ]$</i>
$\alpha := \frac{\alpha_0 \cdot \pi}{180} = 0.541052$	<i>Command for the angle $\alpha [rad]$ $\alpha = 0.541052$</i>
$\beta_0 := 90 - \alpha_0 = 59$	<i>Command for the angle $\beta [^\circ]$</i>
$a := \frac{p \cdot \sin(\alpha)}{1 + \cos(\alpha)} = 14.835375$	<i>Command and result: length of the side a</i>
$b := \frac{p \cdot \cos(\alpha)}{1 + \cos(\alpha)} = 24.690211$	<i>Command and result: length of the side b</i>
$c := \frac{p}{1 + \cos(\alpha)} = 28.804425$	<i>Command and result: length of the side c</i>

◆ Program *Mathematica 7.0* [1],[2],[5]

<i>Algorithm:</i>	<i>Commentary:</i>
p:=53.4946	<i>Given the sum of the sides $b+c$</i>
α0:=31	<i>Given measure of the angle $\alpha [^\circ]$</i>
A:=A0*N[Pi]/180	<i>Command for the angle $\alpha [rad]$ $\alpha = 0.541052$</i>
B=90-A0	<i>Command for the angle $\beta [^\circ]$</i>
a=(p*Sin[A])/(1+Cos[A])	<i>Command: length of the side a</i>
b=(p*Cos[A])/(1+Cos[A])	<i>Command: length of the side b</i>
c=p/(1+Cos[A])	<i>Command: length of the side c</i>
59	<i>Result: measure of the angle $\beta [^\circ]$</i>
14.8354	<i>Result: length of the side a</i>
24.6902	<i>Result: length of the side b</i>
28.8044	<i>Result: length of the side c</i>

Numerical results: $a = 14.8354$, $b = 24.6902$, $c = 28.8044$, $\beta = 59^\circ$.

2.4. Case 4: Sum of two sides length $b+c = p$ and a measure of the angle β

2.4.1. Theoretical analysis

➤ Let be a triangle as shown on the figure 4.




Fig. 4. Right triangle ABC with given a sum of the sides length $b+c=p$ and a measure of the angle β

Source: Elaboration of the Authors

➤ *Data in $\triangle ABC$:* Sum of the sides length $b+c=p$ and a measure of the angle β .

➤ *Unknown in $\triangle ABC$:* Length of the sides a, b, c and a measure of the angle α .

➤ *Solution:*

◆ From some properties of acute angles in a right triangle we obtain:

$$\alpha = \frac{\pi}{2} - \beta. \quad (44)$$

◆ From the sum of the triangle sides we have:

$$b + c = p, \quad (45)$$

$$b = p - c. \quad (46)$$

◆ From the formulae (4) and (6) we obtain:

$$a = c \cdot \cos(\beta), \quad (47)$$

$$b = c \cdot \sin(\beta). \quad (48)$$

◆ From the formulae (46) and (48) we determine the triangle side c :

$$p - c = c \cdot \sin(\beta), \quad (49)$$

$$p = c \cdot [1 + \sin(\beta)], \quad (50)$$

$$c = \frac{p}{1 + \sin(\beta)}. \quad (51)$$

◆ Substituting (51) into the formulae (47) and (48) we determine the triangle sides a, b :

$$a = \frac{p \cdot \cos(\beta)}{1 + \sin(\beta)}, \quad (52)$$

$$b = \frac{p \cdot \sin(\beta)}{1 + \sin(\beta)}. \quad (53)$$

➤ *Answer:* $\alpha = \frac{\pi}{2} - \beta$, $a = \frac{p \cdot \cos(\beta)}{1 + \sin(\beta)}$, $b = \frac{p \cdot \sin(\beta)}{1 + \sin(\beta)}$, $c = \frac{p}{1 + \sin(\beta)}$.

2.4.2. Numerical algorithms in MS-Excel, MathCAD and Mathematica programs

For numerical analysis we take a sum of the sides $b+c = 53.4946$ and the angle $\beta = 59^\circ$.

- ◆ Program *MS-Excel 7.0* [3],[8]

<i>Algorithm:</i>	<i>Commentary:</i>
A6=53,4946	<i>Given the sum of the sides $b+c$</i>
A7=59	<i>Given a measure of the angle $\beta [^\circ]$</i>
A8=A7*PI()/180	<i>Command for the angle $\beta [\text{rad}]$, $\beta = 1.029744$</i>
A9=90-A7	<i>Command for the angle $\alpha [^\circ]$</i>
A10=(A6*COS(A8))/(1+SIN(A8))	<i>Command for calculation of the side a</i>
A11=(A6*SIN(A8))/(1+SIN(A8))	<i>Command for calculation of the side b</i>
A12=A6/(1+SIN(A8))	<i>Command for calculation of the side c</i>
31	<i>Result: measure of the angle $\alpha [^\circ]$</i>
14,8354	<i>Result: length of the side a</i>
24,6902	<i>Result: length of the side b</i>
28,8044	<i>Result: length of the side c</i>

- ◆ Program *MathCAD 15* [9],[12]

<i>Algorithm:</i>	<i>Commentary:</i>
p:=53.4946	<i>Given the sum of the sides $b+c$</i>
β0:=59	<i>Given measure of the angle $\beta [^\circ]$</i>
β := β0 · π / 180 = 1.029744	<i>Command for the angle $\beta [\text{rad}]$, $\beta = 1.029744$</i>
α0 := 90 - β0 = 31	<i>Command for the angle $\alpha [^\circ]$</i>
a := p · cos(β) / (1 + sin(β)) = 14.835375	<i>Command and result: length of the side a</i>
b := p · sin(β) / (1 + sin(β)) = 24.690211	<i>Command and result: length of the side b</i>
a := p / (1 + sin(β)) = 28.804425	<i>Command and result: length of the side c</i>

- ◆ Program *Mathematica 7.0* [1],[2],[5]

<i>Algorithm:</i>	<i>Commentary:</i>
p := 53.4946	<i>Given the sum of the sides $b+c$</i>
A0 := 59	<i>Given measure of the angle $\beta [^\circ]$</i>
A := A0 * N[π] / 180	<i>Command for the angle $\beta [\text{rad}]$, $\beta = 1.029744$</i>
B = 90 - A0	<i>Command for the angle $\alpha [^\circ]$</i>
a = (p * Cos[A]) / (1 + Sin[A])	<i>Command: length of the side a</i>
b = (p * Sin[A]) / (1 + Sin[A])	<i>Command: length of the side b</i>
c = p / (1 + Sin[A])	<i>Command: length of the side c</i>
31	<i>Result: measure of the angle $\alpha [^\circ]$</i>
14.8354	<i>Result: length of the side a</i>
24.6902	<i>Result: length of the side b</i>
28.8044	<i>Result: length of the side c</i>

Numerical results: $a = 14.8354$, $b = 24.6902$, $c = 28.8044$, $\alpha = 31^\circ$.

2.5. Case 5: Sum of two sides length $a+b=q$ and a measure of the angle α .

2.5.1. Theoretical analysis

➤ Let be a triangle as shown on the figure 5.




Fig. 5. Right triangle ABC with given a sum of the sides length $a+b=q$ and a measure of the angle α
Source: Elaboration of the Authors

➤ *Data in $\triangle ABC$:* Sum of the sides length $a+b=q$ and a measure of the angle α .

➤ *Unknown in $\triangle ABC$:* Length of the sides a, b, c and a measure of the angle β .

➤ *Solution:*

◆ From some properties of acute angles in a right triangle we obtain:

$$\beta = \frac{\pi}{2} - \alpha. \quad (54)$$

◆ From the sum of the triangle sides we have:

$$a + b = q, \quad (55)$$

$$a = q - b. \quad (56)$$

◆ From the formulae (7) i (5) we obtain:

$$a = b \cdot \operatorname{tg}(\alpha), \quad (57)$$

$$b = c \cdot \cos(\alpha). \quad (58)$$

◆ From the formula (58) we determine the triangle side c :

$$c = \frac{b}{\cos(\alpha)}. \quad (59)$$

◆ From the formula (56) and (57) we determine the triangle side b :

$$q - b = b \cdot \operatorname{tg}(\alpha), \quad (60)$$

$$q = b \cdot [1 + \operatorname{tg}(\alpha)], \quad (61)$$

$$b = \frac{q}{1 + \operatorname{tg}(\alpha)}. \quad (62)$$

◆ From the formulae (57), (59) and (62) we finally determine the triangle side a and c :

$$a = \frac{q \cdot \operatorname{tg}(\alpha)}{1 + \operatorname{tg}(\alpha)}, \quad (63)$$

$$c = \frac{q}{\cos(\alpha)[1 + \operatorname{tg}(\alpha)]} \equiv \frac{q \cdot \sec(\alpha)}{1 + \operatorname{tg}(\alpha)}. \quad (64)$$

➤ *Answer:* $\beta = \frac{\pi}{2} - \alpha$, $a = \frac{q \cdot \operatorname{tg}(\alpha)}{1 + \operatorname{tg}(\alpha)}$, $b = \frac{q}{1 + \operatorname{tg}(\alpha)}$, $c = \frac{q \cdot \sec(\alpha)}{1 + \operatorname{tg}(\alpha)}$.

2.5.2. Numerical algorithms in MS-Excel, MathCAD and Mathematica programs

For numerical analysis we take a sum of the sides $a+b = 39.5256$ and the angle $\alpha = 31^\circ$.

◆ Program *MS-Excel 7.0* [3],[8]

<i>Algorithm:</i>	<i>Commentary:</i>
A6=39,5256	<i>Given the sum of the sides $a+b$</i>
A7=31	<i>Given a measure of the angle $\alpha [^\circ]$</i>
A8=A7*PI()/180	<i>Command for the angle $\alpha [rad]$, $\alpha = 0.541052$</i>
A9=90-A7	<i>Command for the angle $\beta [^\circ]$</i>
A10=(A6*TAN(A8))/(1+TAN(A8))	<i>Command for calculation of the side a</i>
A11=A6/(1+TAN(A8))	<i>Command for calculation of the side b</i>
A12=A6/(COS(A8)*(1+TAN(A8)))	<i>Command for calculation of the side c</i>
59	<i>Result: measure of the angle $\beta [^\circ]$</i>
14,8354	<i>Result: length of the side a</i>
24,6902	<i>Result: length of the side b</i>
28,8044	<i>Result: length of the side c</i>

◆ Program *MathCAD 15* [9],[12]

<i>Algorithm:</i>	<i>Commentary:</i>
q:=39.5256	<i>Given the sum of the sides $a+b$</i>
α0:=31	<i>Given measure of the angle $\alpha [^\circ]$</i>
α := $\frac{\alpha_0 \cdot \pi}{180} = 0.541052$	<i>Command for the angle $\alpha [rad]$ $\alpha = 0.541052$</i>
β0 := $90 - \alpha_0 = 59$	<i>Command for the angle $\beta [^\circ]$</i>
a := $\frac{q \cdot \tan(\alpha)}{1 + \tan(\alpha)} = 14.835375$	<i>Command and result: length of the side a</i>
b := $\frac{q}{1 + \tan(\alpha)} = 24.690211$	<i>Command and result: length of the side b</i>
a := $\frac{q \cdot \sec(\alpha)}{1 + \tan(\alpha)} = 28.804425$	<i>Command and result: length of the side c</i>

◆ Program *Mathematica 7.0* [1],[2],[5]

<i>Algorithm:</i>	<i>Commentary:</i>
q := 39.5256	<i>Given the sum of the sides $a+b$</i>
A0 := 31	<i>Given measure of the angle $\alpha [^\circ]$</i>
A := A0*N[Pi]/180	<i>Command for the angle $\alpha [rad]$ $\alpha = 0.541052$</i>
B = 90 - A0	<i>Command for the angle $\beta [^\circ]$</i>
a = (q*Tan[A])/(1+Tan[A])	<i>Command: length of the side a</i>
b = q/(1+Tan[A])	<i>Command: length of the side b</i>
c = (q*Sec[A])/(1+Tan[A])	<i>Command: length of the side c</i>
59	<i>Result: measure of the angle $\beta [^\circ]$</i>
14.8354	<i>Result: length of the side a</i>
24.6902	<i>Result: length of the side b</i>
28.8044	<i>Result: length of the side c</i>

Numerical results: $a = 14.8354$, $b = 24.6902$, $c = 28.8044$, $\beta = 59^\circ$.

2.6. Case 6: Sum of two sides length $a+b=q$ and a measure of the angle β

2.6.1. Theoretical analysis

➤ Let be a triangle as shown on the figure 6.




Fig. 6. Right triangle ABC with given a sum of the sides length $a+b=q$ and a measure of the angle β

Source: Elaboration of the Authors

➤ *Data in $\triangle ABC$:* Sum of the sides length $a+b=q$ and a measure of the angle β .

➤ *Unknown in $\triangle ABC$:* Length of the sides a, b, c and a measure of the angle α .

➤ *Solution:*

◆ From some properties of acute angles in a right triangle we obtain:

$$\alpha = \frac{\pi}{2} - \beta. \quad (65)$$

◆ From the sum of the triangle sides we have:

$$a + b = q, \quad (66)$$

$$a = q - b. \quad (67)$$

◆ From the formulae (10) i (4) we obtain:

$$a = b \cdot \operatorname{ctg}(\beta), \quad (68)$$

$$b = c \cdot \sin(\beta). \quad (69)$$

◆ From the formula (69) we determine the triangle side c :

$$c = \frac{b}{\sin(\beta)}. \quad (70)$$

◆ From the formula (67) and (68) we determine the triangle side b :

$$q - b = b \cdot \operatorname{ctg}(\beta), \quad (71)$$

$$q = b \cdot [1 + \operatorname{ctg}(\beta)], \quad (72)$$

$$b = \frac{q}{1 + \operatorname{ctg}(\beta)}. \quad (73)$$

◆ From the formulae (68), (70) and (73) we finally determine the triangle side a and c :

$$a = \frac{q \cdot \operatorname{ctg}(\beta)}{1 + \operatorname{ctg}(\beta)}, \quad (74)$$

$$c = \frac{q}{\sin(\beta)[1 + \operatorname{ctg}(\beta)]} \equiv \frac{q \cdot \operatorname{cosec}(\beta)}{1 + \operatorname{ctg}(\beta)}. \quad (75)$$

➤ *Answer:* $\alpha = \frac{\pi}{2} - \beta$, $a = \frac{q \cdot \operatorname{ctg}(\beta)}{1 + \operatorname{ctg}(\beta)}$, $b = \frac{q}{1 + \operatorname{ctg}(\beta)}$, $c = \frac{q \cdot \operatorname{cosec}(\beta)}{1 + \operatorname{ctg}(\beta)}$.

2.6.2. Numerical algorithms in MS-Excel, MathCAD and Mathematica programs

For numerical analysis we take a sum of the sides $a+b = 39.5256$ and the angle $\beta = 59^\circ$.

◆ Program *MS-Excel 7.0* [3],[8]

Algorithm:	Commentary:
A6=39,5256	<i>Given the sum of the sides $a+b$</i>
A7=59	<i>Given a measure of the angle $\beta [^\circ]$</i>
A8=A7*PI()/180	<i>Command for the angle $\beta [\text{rad}]$, $\beta = 1,030$</i>
A9=90-A7	<i>Command for the angle $\alpha [^\circ]$</i>
A10=(A6/TAN(A8))/(1+(1/TAN(A8)))	<i>Command for calculation of the side a</i>
A11=A6/(1+(1/TAN(A8)))	<i>Command for calculation of the side b</i>
A12=A6/((SIN(A8))*(1+TAN(A8)))	<i>Command for calculation of the side c</i>
31	<i>Result: measure of the angle $\beta [^\circ]$</i>
14,8354	<i>Result: length of the side a</i>
24,6902	<i>Result: length of the side b</i>
28,8044	<i>Result: length of the side c</i>

◆ Program *MathCAD 15* [9],[12]

Algorithm:	Commentary:
q:=39.5256	<i>Given the sum of the sides $a+b$</i>
β0:=59	<i>Given measure of the angle $\beta [^\circ]$</i>
β := β0 · π / 180 = 1.029744	<i>Command for the angle $\beta [\text{rad}]$</i>
α0 := 90 - β0 = 31	<i>Command for the angle $\alpha [^\circ]$</i>
a := q · cot(β) / (1 + cot(β)) = 14.835375	<i>Command and result: length of the side a</i>
b := q / (1 + cot(β)) = 24.690211	<i>Command and result: length of the side b</i>
a := q / (sin(β) · (1 + cot(β))) = 28.804425	<i>Command and result: length of the side c</i>

◆ Program *Mathematica 7.0* [1],[2],[5]

Algorithm:	Commentary:
q:=39.5256	<i>Given the sum of the sides $a+b$</i>
A0:=31	<i>Given measure of the angle $\beta [^\circ]$</i>
A:=A0*Pi/180	<i>Command for the angle $\beta [\text{rad}]$ $\beta = 1.02974$</i>
B=90-A0	<i>Command for the angle $\alpha [^\circ]$</i>
a=(q*Cot[A])/(1+Cot[A])	<i>Command: length of the side a</i>
b=q/(1+Cot[A])	<i>Command: length of the side b</i>
c=q/((Sin[A])*(1+Cot[A]))	<i>Command: length of the side c</i>
59	<i>Result: measure of the angle $\alpha [^\circ]$</i>
14.8354	<i>Result: length of the side a</i>
24.6902	<i>Result: length of the side b</i>
28.8044	<i>Result: length of the side c</i>

Numerical results: $a = 14.8354$, $b = 24.6902$, $c = 28.8044$, $\alpha = 31^\circ$.

3. Conclusions

- Discussed analytical models of solving right triangles allow to perform the other theoretical considerations in each of the six basic cases, taking into account the definition of trigonometric functions and the corresponding properties and formulae.
- Created numerical algorithms of solving the right triangles in *MS-Excel*, *MathCAD*, and *Mathematica* programs allow for significant execution of faster calculations than the traditional way of using logarithms and logarithmic tables.

Literature

- [1] Abell M.L., Braselton J.P.: *Mathematica by example. Revised edition*. AP Professional. A Division of Harcourt Brace & Company. Boston San Diego New York London Sydney Tokyo Toronto 1994.
- [2] Blachman N.: *Mathematica: A Practical Approach*. Prentice-Hall, 1992.
- [3] Bourg D.M.: *Excel in science and technology. Recipes*. HELION Pub. House, Gliwice 2006 (*in Polish*).
- [4] Bronsztejn I.N., Siemiendajew K.A., Musiol G., Mühlig H.: *Modern compendium of mathematics*. Polish Scientific Publishers, Warsaw 2004 (*in Polish*).
- [5] Crandall R.E.: *Mathematica for sciences*. Addison-Wesley, 1991.
- [6] Dolciani M.P., Berman S.L., Wooton W., Meder A.E.: *Modern algebra and trigonometry. Structure and Method. Book Two*. Houghton Mifflin Company, Boston New York Atlanta Geneva, Ill. Dallas Palo Alto 1965.
- [7] Gabszewicz Z.: *Trigonometry. Handbook for trainees in the field of secondary schools course. Set of problems*. Gebethner & Wolff Publishing House, Cracov, Gebethner & Company, Warsaw 1907 (*in Polish*).
- [8] Gonet M.: *Excel in scientific computing and engineering*. HELION Publishing House, Gliwice 2010 (*in Polish*).
- [9] Jakubowski K.: *MathCAD 2000 Professional*, EXIT Publishing House, Warsaw 2000.
- [10] Neill Hugh: *Trigonometry. A complete introduction*. Teach Yourself, 2013.
- [11] Nowosiłow S.I.: *Special lecture of trigonometry*. Polish Scientific Publishers, Warsaw 1956 (*in Polish*).
- [12] Paleczek W.: *MathCAD 12, 11, 2001i, 2001, 2000 in algorithms*, EXIT Publishing House, Warsaw 2005 (*in Polish*).
- [13] Pokorny E.J.: *Trigonometry for the self-taught*. Polish Betting School Publishing (PZWS), Warsaw 1962 (*in Polish*).
- [14] Wojtowicz Wł.: *Trigonometry. The 5th edition*. Polish Betting School Publishing (PZWS), Warsaw 1948, (*in Polish*).
- [15] Wojtowicz Wł., Bielecki B., Czyżkowski M.: *Trigonometry for classes X-XI. The 16th edition*. Polish Betting School Publishing (PZWS), Warsaw 1964, (*in Polish*).
- [16] Young J.W., Morgan F.M.: *Plane trigonometry and numerical computation*, The MacMillan Company, New York 1919.