

Tetiana FARTUSHOK<sup>1\*</sup>, Orysia KOVALYSHYN<sup>1\*</sup>, Yuri FEDEVYCH<sup>1</sup>, Igor TANCHYN<sup>1</sup>  
and Volodymyr ZHYKOVSKIY<sup>1</sup>

## EPIGENETICS IS PROMISING DIRECTION IN MODERN SCIENCE

**Abstract:** Epigenetics studies the inherited changes in a phenotype or in expression of genes caused by other mechanisms, without changing the nucleotide sequence of DNA. The most distinguished epigenetic tools are: modifications of histones, enzymatic DNA methylation, and gene silencing mediated by small RNAs (miRNA, siRNA). The resulting m5C residues in DNA substantially affect the cooperation of proteins with DNA. It is organized by hormones and aging-related alterations, one of the mechanisms controlling sex and cellular differentiation. DNA methylation regulates all genetic functions: repair, recombination, DNA replication, as well as transcription. Distortions in DNA methylation and other epigenetic signals lead to diabetes, premature aging, mental dysfunctions, and cancer.

**Keywords:** epigenetics, 5-methylcytosine, DNA methylation, histones

### Introduction

Recent scientific studies have highlighted the important role of fine-tuning mechanisms in the processes of genome functioning and its interaction with the environment.

The obtained results proved that a significant part of these data lies in those processes that are described by studies of hereditary properties of the organism, which are not directly related to the changes of the nucleotide sequence in a DNA molecule [1-3].

Despite the successes of molecular biology and genetics, it is still unclear how the usual growth of an organism occurs, as cells containing the same genetic data from the beginning, precisely and correctly implement the peculiarities of certain parts of the genome in a specific phenotype. Epigenetics helps to answer the following questions: carcinogenesis, plasticity of stem cells, cell identity (specificity), regeneration of cells and tissues in animals and plants, aging, programmed death [4, 5].

There are a number of factors that regulate the expression of genes: methyl CpG DNA-binding and other proteins, histone modifications, hormone-receptor complexes, DNA methylation, nucleosome remodeling and chromatin overall rearrangement, and interaction with short non-encoding RNAs. Gene expression requires complex complexes with more than 100 proteins that are involved in initiating and prolonging transcription

---

<sup>1</sup> Danylo Halytsky Lviv National Medical University, Pekarska, 69, 79010, Ukraine, phone +38 032 275-76-32, email: office@meduniv.lviv.ua, ORCID: TF 0000-0001-6571-0108, OK 0000-0002-9710-0694, YF 0000-0002-2536-8376, IT 0000-0002-0863-4732, VZ 0000-0002-0594-5316

\* Corresponding author: fartushok1@ukr.net, orusia75@gmail.com

from a single promoter and iRNA processing. Epigenetic control can amplify the initial signal (the promoter stimulation) or carry out the silencing of the gene. Epigenetic memory is sometimes related to particular alterations of histones in chromatin [6, 7].

In particular, this can occur with gene silencing this prohibition (conformational "lock") in a specific area of the chromosome may become substantial as a consequence of induced (allowed) methylation of DNA by this chromatin state [8].

The factors related to the environment can have a noticeable effect on the enzymes' activity, which carry out alterations of DNA and histones. These cofactors incorporate: acetylcoenzyme A for acetylase, ATP for kinases, S-adenosylmethionine (SAM) for methyltransferases. The level of those cofactors in the cell can differ remarkably and rely upon the environment. Adequate epigenetic control is formed on the basis of many factors' balance [9].

How data and information concerning the chromatin's structure is realized to the daughter from the mother cell? This can happen in this way. This is clear that the of "core" histone proteins' synthesis is clearly organized in the cycle of cell. In the process of chromatin formation, certain proteins, relying upon the nature and degree of their enzyme alterations, can bind to the DNA to shape corresponding sites for transcription [10].

Changes in the character of histone and DNA methylation have been detected; structural alterations in the nucleosome are not resulted from a gene that has undergone mutations directly, therefore, it can be regarded as epigenetic aberration. That also is applicable to mutations in the synthesis and utilization of SAM genes: the absence of SAM results in a violation of trans-methylation responses in the cell and inactivation of a number of enzymes for which SAM is an allosteric effector. However, although epigenetic changes are transmitted by heredity, this is not infinitely [11]. The central biology dogma of DNA  $\leftrightarrow$  RNA  $\rightarrow$  protein is currently supplemented with knowledge regarding prions proteins that are capable of replication, inherited without the participation of DNA and RNA matrices [12].

Chromatine proteins histones are inhibitors of transcription: the DNA free of them is much better transcribed than chromatin bound to them. There was an impression that for effective transcription it seems vital to "undress" the DNA, releasing it from the histones. Nonetheless, Allfrey and Mirsky demonstrated that histones can be proacetylated to activate transcription of inactive chromatin, which is accompanied by a significant weakening of the binding of these proteins to DNA [13]. Widely known modifications of histones include acetylation, phosphorylation, methylation, ubiquitination [14].

There are many different epigenetic signals in the cell and in the organism. Among these signals are currently known [15, 16]:

- Methylation of DNA;
- Several histones' enzymatic modifications;
- Chromosomal and genomic rearrangements;
- Small non-coding RNA (siRNA, or so-called small interfering RNA).

Epigenetics is the study of epigenetic signals, their nature, their interaction and the physico-chemical effects that they cause, their biological action in the cell at several functional levels of the organism and in different conditions of the external and internal environment.

## Methylation of DNA - an epigenetic mechanism for monitoring the genetic functions of the body

Scientists have discovered that the features of all higher plants' DNA is somewhat high content of their additional base - 5-methylcytosine ( $m^5C$ ). Later in plant DNA, as in bacteria, was found  $N^6$ -methyladenin ( $m^6A$ ) (Table 1) [17].

Table 1

Minor methylated bases in DNA [11, 13]

Organisms	Minor bases [%]	
	$m^5C$	$m^6A$
Bacteria	0.01-1.53	0.02-0.70
Algae	0.20-3.50	0.10-0.60
Mushrooms	+	0-0.5
The simplest		0.3-1.0
Plants	2.0-10.0	0.5-1.0
Invertebrates	0.1-2.5	?
Vertebrates	0.7-3.5	+

For several years, these bases' origin in the DNA stayed unrevealed. First in bacteria and then in eukaryotes in 1963, enzymes were detected, which in the existence of a donor of methyl groups of S-adenosylmethionine selectively methylated individual residues of cytosine and adenine in the DNA chains. Scientists have investigated that the "minor" bases ( $m^5C$  and  $m^6A$ ) found in the DNA molecule are not embedded in them in the finished form. They are formed as a consequence of enzymatic alteration (methylation) of the corresponding bases (Fig. 1) in the shaped chains of DNA [18, 19].




Fig. 1. Modifications (methylation) of the corresponding bases in the DNA chains (arrows show the locations of the methylated points) [20]

In this case, the enzyme DNA-methyltransferase forms a covalently linked complex with DNA extracted from the double-stranded DNA of the modified base DNA and methylates it (Fig. 2).




Fig. 2. Complex of cytosine DNA-methyltransferase from DNA [21]

After that, the covalent bond between the DNA and the enzyme is broken, the complex breaks down, and the methylated base ( $m^5C$ ) gets back to its position in the DNA structure [22]. Methylation of DNA has a substantial effect on its relation (binding) with different proteins. Particularly, proteins that specially bind to the regulatory elements of rRNA genes have been detected in plant nuclei and indicated that some of those nuclear proteins binding is modulated by methylation of in vitro cytosine DNA residues [23, 24].

In several cases, the methylation of DNA by cytosine residues stop its binding to specifically reacting with proteins that carry out several genetic procedures, such as replication, transcription, as well as DNA repair [25, 26]. The so-called  $m^5SrG$  DNA-binding proteins, which are specifically located on the DNA of complex protein complexes, control and express genes [27].

If protein-free DNA is incubated with S-adenosylmethionine labeled in the methyl group, then after a while this label is already present in the DNA in the form of the formed 5-methylcytosine and thymine. Thus, non-enzyme methylation of DNA was discovered (Fig. 3) [28].




Fig. 3. Non-enzymatic methylation of DNA [28]

Regarding that case, the labeled thymine in DNA was more than m<sup>5</sup>C. Therefore, it was discovered that the non-enzyme methylation of DNA in the solution is accompanied by rapid oxidative deamination of the formed residues m<sup>5</sup>C with conversion to the remainders of thymine. This proved that the cytosine residues methylation in DNA can result in C → T transitions (GC-pair of bases is substituted for AT-pair), and the residues of 5-methylcytosine act as mutational points [29].

## Specificity of enzyme methylation of DNA

DNA-methyltransferase enzymes have used the possibility of DNA methylation, they modify cytosolic or adenine residues in certain nucleotide sequences [30]. Methylation of DNA is also carried out in animal cells, where has a significant biological significance. Methylation of cytosine residues is mainly observed in the methylation of DNA, which is induced by small RNAs [31].

Methylation of DNA in animals and plants, carried out by site-specific enzymes with cytosine DNA-methyltransferases, leads to the occurrence of 5-methylcytosine residues (m<sup>5</sup>C) in the sequences of CG, CNG and SNN. Adenine methylation of DNA has been discovered in plants. The appearance of m<sup>5</sup>C residues in DNA significantly affects the interaction of DNA with proteins. DNA methylation often blocks the binding of DNA to such proteins and prevents gene transcription, and on the other hand, it is a prerequisite for binding proteins [31]. There are special m<sup>5</sup>CrG DNA binding proteins. Linking proteins with DNA arranges the entire ensemble of proteins of the transcription machinery and is required for its activity [32].

In animals and plants, along with the linking proteins species, there is also tissue (cellular) subcellular (organoid) and age-specific DNA methylation. Methylation of DNA in animals decreases with age [33].

It has been proven that in the mitochondria and nucleus of the same cell DNA is methylated in different ways. In the mitochondrial DNA from the heart of the bull, 5-methylcytosine is detected. Cytosine DNA-methyltransferase has been isolated from animal mitochondria and it has been shown that this enzyme has another site specificity compared to nuclear DNA-methyl transferase. Unlike animals, 5-methylcytosine was not found in plants in DNA of mitochondria, but N<sup>6</sup>-methylenadenine was detected in them [34].

Three cytosine DNA-methyltransferases types and above ten genes encoding DNA-methyltransferases have been identified in plants. This is much more than in the eukaryotes. These enzymes carry out the methylation of the CpG sites for DNA replication, generally ensuring the conservation and transfer of the inherent inheritance of the overall picture of the methylation of these sites in the genome [35]. In plants, methylation of DNA is controlled by phytohormones and specific regulators of plant growth and development. Due to the influence of various phytohormones in plants, methylation of DNA in the cell cycle significantly decreases [36].

In humans, three enzymes, the so-called DNA-methyltransferases 1, 3a and 3b (DNMT1, DNMT3a and DNMT3b), correspond to the DNA methylation process. It is anticipated that DNMT3a and DNMT3b are de novo synthesized methyltransferases that create the formation of DNA methylation in the early stages of development, while DNMT1 carries out DNA methylation on subsequent stages of the body's life. The enzyme DNMT1 has a high affinity to 5-methylcytosine. When DNMT1 finds a "semi-methylated site" (a site where methylated cytosine is only in one DNA strand), it methylates cytosine

of the second thread on the same site. The function of methylation is to activate / inactivate the gene. In most cases, methylation of the promoter regions of the gene leads to inhibition of gene activity. It is shown that even minor changes in the degree of methylation of DNA can significantly alter the genetic expression [37].

DNA-methyltransferases are redox sensitive enzymes. The DNMT1 enzyme *in vivo* is responsible for the proper development of the embryo, imprinting, inactivation of the X chromosome and restoring the DNA structure after replication. It is present in most cases in somatic cells and is localized in the forks of DNA replication, interacting with the nuclear antigen of proliferating cells (PCNA). Enzymes DNMT3a and DNMT3b are involved in *de novo* methylation. Expressed in embryonic and somatic cells of an adult. DNMT 3L is present in the germ cells and interacts with DNMT3a and DNMT3b in imprinting of parent genes. DNMT2 is detected in all cell types, but does not exhibit enzyme activity, so its function is not yet established [38].

A number of diseases, in particular cancer, are accompanied by abnormal hypomethylation of DNA and hypermethylation of CpG-islands in the promoter portions of proapoptotic genes (suppressors of oncogenes), which leads to sustained repression of transcription [39, 40]. The transcriptional repression in this case is mediated by proteins that can bind to methylated CpG dinucleotides. Methylcytosine-binding proteins activate histone deacetylase (HDAC) and other factors involved in chromatin remodeling [39].

The formed complex may modify histones, forming a condensed transcriptionally inactive structure of heterochromatin. The effect of methylation of DNA on chromatin structure is of great importance for the development and functioning of a living organism. In particular, the absence of methylcytosine-binding protein 2 (MeCP2) due to mutations in the corresponding gene, leads to the development of Rett syndrome in humans [40].

The inactivation of methylcytosine-binding protein domain 2 (Methyl-CpG binding domain protein 2 - MBD2), which is involved in the repression of transcription of hypermethylated genes, is observed in oncological diseases [41].

Methylation of DNA in animals and plants has a lot in common. Nevertheless, in plants it has several specific features. The methylated CNG proportion and asymmetric DNA sequences in plant genomes is more substantial compared to that of in animals. Plants own a significantly more complex system of methylation of genomes than animals [42].

Contrary to animals, plants own particular organelles - plastids (amyloplasts, leukoplasts, chloroplasts, chromoplasts), which have their own, different from nuclear, system of modification (methylation) of DNA. Those systems can have a substantial role in the functioning and differentiation of plastids. DNA Methylation (mtDNA) in plant mitochondria is different from that in the nucleus. N<sup>6</sup>-methyladenine was found in plant mtDNA, but not 5-methylcytosine, which is inherent in animal mtDNA [43].

In plant there is a system of restriction - modification of the genome. The plants have S-adenosylmethionine-dependent endonuclease, which are sensitive to the status of methylation of DNA. Plant endonuclease is to some extent similar to a typical bacterial restriction endonuclease [44].

Three components of this complex system: substrate DNA, enzyme (DNA methyltransferase) and donor methyl groups S-adenosylmethionine are involved in genome modification (Fig. 4). Control over the modification of DNA and the effectiveness of this process is carried out at the level of all these components, and with the participation of other diverse components of cellular metabolism. In addition, the methylation of the

genome in a differentiated cell at a certain stage of ontogeny also depends on the activity of the enzymes that demethylate the DNA.




Fig. 4. Methylation and demethylation of DNA [44]

However, often in the presence of these active enzymes, a sufficient amount of S-adenosylmethionine (a donor of methyl groups and an enzyme activity modulator) and in the absence of appropriate inhibitors, these reactions in the nucleus are not possible due to the inaccessibility of the substrate-DNA in the chromatin for enzymes. Here it is important to organize the chromatin itself [45].

In addition to the already mentioned multiple modifications of histones that modulate the chromatin organization and the availability of DNA for enzymes, many other proteins compete for the binding and interaction of DNA-methyltransferases with DNA. In particular, they can include proteins of hormone-receptor complexes.

The cytosine methylation of DNA controls plant growth and development (Vanyushin B.) and animals (Pugh), is involved in the regulation of all genetic processes, including transcription, replication, DNA repair, cell differentiation, genomic imprinting, and gene transcription [46].

The discovery of specific, so-called small RNAs (siRNAs, miRNAs) played an important role in representing the molecular mechanisms of gene expression regulation. These RNAs are encoded in the inverted sequences of the genome, which, as a result of reading and subsequent processing, appear in the form of short 12-14-member oligonucleotides.

## Modification of histones

In the cell there are many other systems of epigenetic signals, they are different. In some of them the important role belongs not to DNA, but to proteins, including histones of chromatin. Due to one or another modification of histones the chromatin structure changes, which leads to inherited changes in the transcription of genes. Modification (methylation, phosphorylation, acetylation, ubiquitination) often determines whether the genes are active or not. The modification is also accompanied by multiple special non-histone regulatory proteins that form complexes with DNA that silence genes or, conversely, trigger them. It is known that there is a correlation between the methylation of DNA and the modification of histones. The histones may act as a carrier of signals for the methylation of the genome [47].

Modification of N-tails is much more frequent, despite the fact that modifications of amino acids in histones occur throughout the protein molecule. Modifications include: acetylation, phosphorylation, methylation.

## Acetylation of histones

Acetylation is the most studied modification of histones (Fig. 5). Thus, acetylation with the acetyltransferase of the 14th and 9th lysines histone H3 (H3K14ac and H3K9ac, respectively) correlates with the transcriptional activity in this region of the chromosome. This is because the acetylation of lysine changes the positive charge to neutral, which makes it impossible to connect it with negatively charged phosphate groups in the DNA. As a result, the histones are disconnected from the DNA. Histones are able to maintain a modified state and act as a matrix for the synthesis of new histones that bind to DNA after replication [48].




Fig. 5. Acetylation and deacetylation of histones [48]

Modifications of histones are indicators of active or suppressed chromatin. Degree of acetylation of histones is determined by activity of two types of enzymes - histone acetyltransferase (HAT) and deacetylase (HDAC) [48].



## Phosphorylation of histones

Phosphorylation of histones occurs mainly on the remnants of serine and threonine, and affects chromatin changes, increasing the negative charge on the histone tail (Fig. 6). This process is subject to dynamic regulation of the kinases (for example, phosphatase). Phosphorylation of histones plays a functional role in regulating the early response to expression, as well as in the dynamics of chromosomes during mitosis [49].




Fig. 6. Histone phosphorylation [50]

Contrary to phosphorylation and acylation, histone methylation doesn't influence the nucleosome charge and can have a beneficial or repressive function in the expression of gene. Methyltransferases and dimethylases act on arginine and lysine, and control the dynamic methylation process. [51, 52]. Modification of histones leads to the induction or enhancement of methylation of CNG sites (Fig. 7).




Fig. 7. Activation of chromatin as a result of demethylation of DNA and acetylation of histones [53]

In a number of these transformations, methylation of DNA can be a reason and a result of "silencing" genes [54, 55].

## RNA-directed DNA methylation and gene silencing

Of particular interest today is the study mechanisms and biological role of methylated RNA-directed DNA methylation, which carries out specific exclusion of genes (gene silencing) [56, 57].

It is believed that site-specific DNA methyltransferase in the presence of a small signal RNA (Fig. 8) is carried out de novo DNA methylation at CNG and other sites in the nucleotide sequence of DNA, recognizing small RNA; this is the methylation of the mobile lysates the corresponding enzymes, which, in particular, and modify histones.




Fig. 8. Gene silencing by small interfering RNAs (siRNAs) [58]




Fig. 9. Epigenetic bases of ontogenesis [59]

Modification of histones leads to induction enhancing or enhancing the methylation of CNG sites, which is then maintained without the participation of RNA trigger (Fig. 7). DNA methylation in the RNA-DNA triplex plays a decisive role in gene silencing small RNAs; in organisms or mutants with defective cytosine DNA methylation this RNA gene silencing is ineffective.

The discovery of specific so-called small RNAs (siRNA, miRNA) have exploded the understanding of molecular mechanisms regulation of gene expression and played an exception significant role in the recognition and strengthening of the epigenetics.

The development of epigenetics as a separate branch of molecular biology began in the 1940s. Then the English geneticist Conrad Waddington formulated the concept of "epigenetic landscape", which explains the process of formation of the organism. The development of epigenetics is presented in Figure 9.

## Conclusion

The essential epigenetic signals in the cell is the methylation of DNA In higher plants, DNA is methylated with cytosine residues; 5-methylcytosine is localized predominantly in the CG and CNG sequences. The ontogeny of plants and animals is generally impossible without the methylation of the genome.

In the nucleus, there are numerous DNA-methyltransferases, the methylation of DNA at different stages of replication may be carried out in different enzymes specificity. The state of methylation of the genome in a differentiated cell at a certain stage of ontogeny also depends on the activity of the enzymes that demethylated DNA. The degree and character of the methylation of the genome can be determined by the ratio of the activity of the enzymes that are methylating and demethyling the DNA.

## References

- [1] Bird A. Perceptions of epigenetics. *Nature*. 2007;447(7143):396-8. DOI: 10.1038/nature05913.
- [2] Ashapkin VV, Kutueva LI, Vanyushin BF. The gene for domains rearranged methyltransferase (DRM2) in *Arabidopsis thaliana* plants is methylated at both cytosine and adenine residues. *FEBS Lett*. 2002;532(3):367-72. DOI: 10.1016/s0014-5793(02)03711-0.
- [3] Chow J, Heard E. X inactivation and the complexities of silencing a sex chromosome. *Current Opinion Cell Biol*. 2009;21:359-66. DOI: 10.1016/j.ceb.2009.04.012.
- [4] Singh RK, Prasad M. Delineating the epigenetic regulation of heat and drought response in plants. *Critical Rev Biotechnol*. 2021;1:4. DOI: 10.1080/07388551.2021.1946004.
- [5] Jiang W, Xia T, Liu C, Li J, Zhang W, Sun C. Remodeling the epigenetic landscape of cancer - Application potential of flavonoids in the prevention and treatment of cancer. *Frontiers Oncol*. 2021;11:2331. DOI: 10.3389/fonc.2021.705903.
- [6] Bartholdy B, Lajugie J, Yan Z, Zhang S, Mukhopadhyay R, Grealley J, et al. Mechanisms of establishment and functional significance of DNA demethylation during erythroid differentiation. *Blood Adv*. 2018;2(15):1833-52. DOI: 10.1182/bloodadvances.2018015651.
- [7] de Oliveira DT, Guerra-Sa R. Uncovering epigenetic landscape: a new path for biomarkers identification and drug development. *Molecular Biol Reports*. 2020;47:9097-122. DOI: 10.1007/s11033-020-05916-3.
- [8] Ehrlich M. DNA hypermethylation in disease: mechanisms and clinical relevance. *J Epigenetics*. 2019;14:1141-63. DOI: 10.1080/15592294.2019.1638701.
- [9] Wilson AS, Power BE, Molloy PL. DNA hypomethylation and human diseases. *Biochim Biophys Acta*. 2007;1775:138-62. DOI: 10.1016/j.bbcan.2006.08.007.
- [10] Le Hellard S, Keller MC, Andreassen OA, Deary IJ, Glahn DC, Malhotra AK. GWAS meta-analysis reveals novel loci and genetic correlates for general cognitive function: a report from the COGENT consortium. *Mol Psychiatry*. 2017;22(3):336-45. DOI: 10.1038/mp.2016.244.
- [11] Ashapkin VV, Kutueva LI, Vanyushin BF. Aging as an epigenetic phenomenon. *Curr Genomics*. 2017;18(5):385-407. DOI: 10.2174/1389202918666170412112130.

- [12] Ryabukha OI, Fedorenko VI. To the structural and functional preconditions of the emergence of thyroid pathology (literature review). *Medicni Perspektivi*. 2021;26(3):169-78. DOI: 10.26641/2307-0404.2021.3.242253.
- [13] Ashapkin VV, Kutueva LI, Vanyushin BF. Epigenetic clock: Just a convenient marker or an active driver of aging? *Adv Exp Med Biol*. 2019;1178:175-206. DOI: 10.1007/978-3-030-25650-0\_10.
- [14] Gangisetty O, Cabrera MA, Murugan S. Impact of epigenetics in aging and age related neurodegenerative diseases. *Front Biosci (Landmark Ed)*. 2018;23:1445-64. DOI: 10.2741/4654.
- [15] Korda Z, Pishva E, van den Hove DL. Epigenetics in Drug Discovery: Achievements and Challenges. In: *Modern CNS Drug Discovery*. Cham: Springer; 2021;57-75. DOI: 10.1007/978-3-030-62351-7\_4.
- [16] Oda M, Glass JL, Thompson RF, Mo Y, Olivier EN, Figueroa ME et al. High-resolution genome-wide cytosine methylation profiling with simultaneous copy number analysis and optimization for limited cell numbers. *Nucleic Acids Res*. 2009;37:3829-39. DOI: 10.1093/nar/gkp260.
- [17] Yi SV, Goodisman MA. The impact of epigenetic information on genome evolution. *Phil Trans Royal Soc B*. 2021;376(1826):20200114. DOI: 10.1098/rstb.2020.0114.
- [18] Baribault C, Ehrlich KC, Ponnaluri VKC. Developmentally linked human DNA hypermethylation is associated with down-modulation, repression, and upregulation of transcription. *Epigenetics*. 2018;2:1-15. DOI: 10.1080/15592294.2018.1445900.
- [19] Kundakovic M, Jiang Y, Kavanagh DH, Dincer A, Brown L, Pothula V, et al. Practical guidelines for high-resolution epigenomic profiling of nucleosomal histones in postmortem human brain tissue. *Biol Psychiatry*. 2017;81(2):162-70. DOI: 10.1016/j.biopsych.2016.03.1048.
- [20] Aref-Eshghi E, Rodenhiser DI, Schenkel LC. Genomic DNA methylation signatures enable concurrent diagnosis and clinical genetic variant classification in neurodevelopmental syndromes. *Am J Hum Genet*. 2018;102(1):156-74. DOI: 10.1016/j.ajhg.2017.12.008.
- [21] Mazzone R, Zwergel C, Artico M. The emerging role of epigenetics in human autoimmune disorders. *Clin Epigenetics*. 2019;11(1):34. DOI: 10.1186/s13148-019-0632-2.
- [22] Evans LM, Tahmasbi R, Vrieze SI, Abecasis GR, Das S, Gazal S, et al. Comparison of methods that use whole genome data to estimate the heritability and genetic architecture of complex traits. *Nat Genet*. 2018; 50(5):737-45. DOI: 10.1038/s41588-018-0108-x.
- [23] Collings CK, Anderson JN. Links between DNA methylation and nucleosome occupancy in the human genome. *Epigenetics Chromatin*. 2017;10(18):1-19. DOI: 10.1186/s13072-017-0125-5.
- [24] Kane AE, Sinclair DA. Epigenetic changes during aging and their reprogramming potential. *Crit Rev Biochem Mol Biol*. 2019;54(1):61-83. DOI: 10.1080/10409238.2019.1570075.
- [25] Ummarino S, Hausman C, Gaggi G, Rinaldi L, Bassal MA, Zhang Y, et al. NAD modulates DNA methylation and cell differentiation. *Cells*. 2021;10(11):2986:1-17. DOI: 10.3390/cells10112986.
- [26] Mochizuki K, Hariya N, Kubota T. Novel models of epigenetic gene regulation in the nutritional environment. *Adv Exp Med Biol*. 2018;1012:11-8. DOI: 10.1007/978-981-10-5526-3\_2.
- [27] Hitchler MJ, Domann FE. The epigenetic and morphogenetic effects of molecular oxygen and its derived reactive species in development. *Free Radic Biol Med*. 2021;170:70-84. DOI: 10.1016/j.freeradbiomed.2021.01.008.
- [28] Domann FE, Hitchler MJ. Aberrant redox biology and epigenetic reprogramming: Co-conspirators across multiple human diseases. *Free Radic Biol Med*. 2021;170:2-5. DOI: 10.1016/j.freeradbiomed.2021.04.020.
- [29] Lamadema N, Burr S, Brewer AC. Dynamic regulation of epigenetic demethylation by oxygen availability and cellular redox. *Free Radic Biol Med*. 2019;131:282-98. DOI: 10.1016/j.freeradbiomed.2018.12.009.
- [30] Kumar SRM, Wang Y, Zhang X, Cheng H, Sun L, He S, et al. Redox components: key regulators of epigenetic modifications in plants. *Int J Mol Sci*. 2020;21(4):1419. DOI: 10.3390/ijms21041419.
- [31] Kumar S., Chinnusamy V., Mohapatra T. Epigenetics of modified DNA bases: 5-Methylcytosine and beyond. *Front Genet*. 2018;9:640-54. DOI: 10.3389/fgene.2018.00640.
- [32] Kumar S, Cheng X, Klimasauskas S. The DNA (cytosine-5) methyltransferases. *Nucleic Acids Res*. 1994;22(1):1-10. DOI: 10.1093/nar/22.1.1.
- [33] Rocha MS, Castro R, Rivera I, Kok RM, Smulders YM, Jakobs C. Global DNA methylation: comparison of enzymatic- and non-enzymatic-based methods. *Clinical Chem Lab Medicine*. 2010;48:1793-8. DOI: 10.1515/CCLM.2010.346.
- [34] Neri F, Incarnato D, Oliviero S. DNA methylation and demethylation dynamics. *Oncotarget*. 2015;6(33):34049-50. DOI: 10.18632/oncotarget.6039.
- [35] Ito K, Barnes PJ, Adcock IM. Histone acetylation and deacetylation. *Methods Mol Med*. 2000;44:309-19. DOI: 10.1385/1-59259-072-1:309.
- [36] Rossetto D, Avvakumov N, Côté J. Histone phosphorylation. *Epigenetics*. 2012;7(10):1098-108. DOI: 10.4161/epi.21975.

- [37] Osta AE, Wolffe AP. DNA Methylation and histone deacetylation in the control of gene expression: Basic biochemistry to human development and disease. *Gene Expr.* 2000;9(1-2):63-75. DOI: 10.3727/00000001783992731.
- [38] Matzke MA, Matzke AJ. Homology dependent gene silencing in transgenic plants: what does it really tell us. *Trends Genet.* 1995;11(1):1-2. DOI: 10.1016/s0168-9525(00)88973-8.
- [39] Santos JH. Mitochondria signaling to the epigenome: A novel role for an old organelle. *Free Radic Biol Med.* 2021;170:59-69. DOI: 10.1016/j.freeradbiomed.2020.11.016. PMID: 33271282
- [40] Stefanowicz D, Ullah J, Lee K, Shaheen F, Olumese E, Fishbane N, et al. Epigenetic modifying enzyme expression in asthmatic airway epithelial cells and fibroblasts. *BMC Pulm Med.* 2017;17(1):1-11. DOI: 10.1186/s12890-017-0371-0.
- [41] Coppedè F. One-carbon epigenetics and redox biology of neurodegeneration. *Free Radic Biol Med.* 2021;170:19-33. DOI: 10.1016/j.freeradbiomed.2020.12.002.
- [42] Hamza M, Halayem S, Mrad R, Bourgou S, Charfi F, Belhadj A. Epigenetics' implication in autism spectrum disorders: A review. *Encephale.* 2017;43(4):374-81. DOI: 10.1016/j.encep.2016.07.007
- [43] Brewer AC. Physiological interrelationships between NADPH oxidases and chromatin remodeling. *Free Radic Biol Med.* 2021;170:109-15. DOI: 10.1016/j.freeradbiomed.2021.01.052.
- [44] Xiao X, Liu X, Jiao B. Epigenetics: Recent advances and its role in the treatment of Alzheimer's disease. *Front Neurol.* 2020;11:538301. DOI: 10.3389/fneur.2020.538301.
- [45] Xu M, Zhu J, Liu XD, Luo MY, Xu NJ. Roles of physical exercise in neurodegeneration: reversal of epigenetic clock. *Transl Neurodegener.* 2021;10:30-45. DOI: 10.1186/s40035-021-00254-1.
- [46] Wang L, Yu CC, Liu XY, Deng XN, Tian Q, Du YJ. Epigenetic modulation of microglia function and phenotypes in neurodegenerative diseases. *Neural Plast.* 2021;2021:1-13. DOI: 10.1155/2021/9912686.
- [47] Pierandrei S, Truglio G, Ceci F, Del Porto P, Bruno SM, Castellani S, et al. M. DNA methylation patterns correlate with the expression of SCNN1A, SCNN1B, and SCNN1G (epithelial sodium channel, ENaC) genes. *Int J Mol Sci.* 2021;22:3754. DOI: 10.3390/ijms22073754.
- [48] Hogan R, Flamier A, Nardini E, Bernier G. The role of BMI1 in late-onset sporadic Alzheimer's disease. *Genes (Basel).* 2020;11:E825. DOI: 10.3390/genes11070825.
- [49] Murshid NM, Lubis FA, Makpol S. Epigenetic changes and its intervention in age-related neurodegenerative diseases. *Cellular Molecular Neurobiol.* 2020. DOI: 10.1007/s10571-020-00979-z.
- [50] Titcombe P, Murray R, Hewitt M, Antoun E, Cooper C, Inskip HM, et al. Human non-CpG methylation patterns display both tissue-specific and inter-individual differences suggestive of underlying function. *Epigenetics.* 2021;1-12. DOI: 10.1080/15592294.2021.1950990.
- [51] Coppedè F. One-carbon epigenetics and redox biology of neurodegeneration. *Free Radical Biol Med.* 2021;170:19-33. DOI: 10.1016/j.freeradbiomed.2020.12.002.
- [52] Domann FE, Hitchler MJ. Introduction to the special issue on 'epigenetics and redox signaling'. *Free Radical Biol Med.* 2021;170. DOI: 10.1016/j.freeradbiomed.2021.04.015.
- [53] García-Giménez J-L, Garcés C, Romá-Mateo C, Pallardó FV. Oxidative stress-mediated alterations in histone post-translational modifications. *Free Radical Biol Med.* 2021;170:6-18. DOI: 10.1016/j.freeradbiomed.2021.02.027.
- [54] Fernandez A, O'Leary C, O'Byrne KJ, Burgess J, Richard DJ, Suraweera A. Epigenetic mechanisms in DNA double strand break repair: A clinical review. *Front Mol Biosci.* 2021;8:685440. DOI: 10.3389/fmolb.2021.685440.
- [55] Yang B, Chen Q. Cross-talk between oxidative stress and m<sup>6</sup>A RNA methylation in cancer. *Oxid Med Cell Longev.* 2021;1-26. DOI: 10.1155/2021/6545728.
- [56] Börner JH, Rawashdeh O, Rami A. Exacerbated age-related hippocampal alterations of microglia morphology,  $\beta$ -amyloid and lipofuscin deposition and presenilin overexpression in per1<sup>-/-</sup>-mice. *Antioxidants (Basel).* 2021;10(9):1330. DOI: 10.3390/antiox10091330.
- [57] Ebrahimi SO, Reisi S, Shareef S. miRNAs, oxidative stress, and cancer: A comprehensive and updated review. *J Cell Physiol.* 2020;235(11):8812-25. DOI: 10.1002/jcp.29724.
- [58] Dionisio PA, Amaral JD, Rodrigues CMP. Oxidative stress and regulated cell death in Parkinson's disease. *Ageing Res Rev.* 2021;67. DOI: 10.1016/j.arr.2021.101263.
- [59] Barbieri M. A new theory of development: the generation of complexity in ontogenesis. *Phil Trans Royal Soc Math Phys Eng Sci.* 2016;1-13. DOI: 10.1098/rsta.2015.0148.