

Dariusz Jakóbczak

Katedra Podstaw Informatyki i Zarządzania
Wydział Elektroniki i Informatyki
Politechnika Koszalińska

The Zero of Function and Interpolation by the Method of Hurwitz-Radon Matrices

1. Introduction

A significant problem in mathematics and computer sciences [1] is to solve any nonlinear equation $g(x) = h(x)$. This means the issue of approximation x in equation $f(x)=0$ where $f=g-h$. Two-dimensional data can be treated as points on the curve. Many numerical methods for nonlinear equations are known as iterative methods: bisection, *regula falsi*, Newton's method (also called as the Newton-Raphson method), Steffensen's method, Brent's method, Broyden's method, fixed-point iterations, inverse interpolation, the secant method [2]. These methods can be used for any function, but sometimes there are troubles. For example in Newton's method we may find difficulties in calculating derivative of a function or troubles with bad starting point of iteration. Generally iterative methods need many assumptions about function (monotonicity, convexity, derivative, starting point). Some methods are used only for polynomials (Muller's, Laguerre's, Bairstow's, Jenkins-Traub's methods). Nonlinear systems are still opened for researchers [3].

This paper is dealing with novel method of root's approximation by using a family of Hurwitz-Radon matrices. Method of Hurwitz-Radon Matrices (MHR) does not need any assumptions about function. The only information about curve is the set of at least five interpolation nodes and a zero of the function between them. Proposed method of Hurwitz-Radon Matrices (MHR) is used in data interpolation and then calculations to solve the nonlinear equation are introduced. MHR connects two significant problems in mathematics: interpolation of the function and the solution of nonlinear equation [4]. MHR method uses two-dimensional data for knowledge representation [5] and computational foundations [6]. Also medicine [7], industry and manufacturing are looking for the methods connected with geometry of the curves [8]. So suitable data representation and precise solving of any equation [9] are key factors in many applications of artificial intelligence and numerical methods [10].

2. Information about Function

Each nonlinear equation is represented by $f(x) = 0$ and succeeding points $(x_i, y_i) \in \mathbf{R}^2$ of function f (interpolation nodes) as follows in proposed MHR method:

1. first node (x_1, y_1) and last node (x_L, y_L) must fulfill a condition $y_1 \cdot y_L < 0$;
2. at least three nodes (x_2, y_2) , (x_3, y_3) , (x_4, y_4) , for example equidistant between first and last node, have to be calculated ($L = 5$) if MHR method is used with matrices of dimension $N = 2$.

Condition 1 is well known in numerical methods for existing a zero of the function. Condition 2 is connected with important features of MHR method: MHR version with matrices of dimension $N = 2$ (as MHR-2) requires at least five nodes ($L = 5$), MHR version with matrices of dimension $N = 4$ (as MHR-4) requires at least nine nodes ($L=9$) and MHR version with matrices of dimension $N = 8$ (as MHR-8) requires at least 17 nodes ($L = 17$).




Fig. 1. Five nodes of function and a root between first and second node (MS Excel graph).

Fig.1 presents the graph of function $f(x) = x^3 + x^2 - x + 1$ with nodes: first (-2;-1), (-1.75; 0.453125), (-1.5;1.375), (-1.25;1.859375) and last (-1;2). All five nodes will be applied in MHR calculations, but we will search a root of function only between nodes (-2;-1) and (-1.75;0.453125). The approximation of a zero point of the function is possible using novel MHR method.

3. Interpolation of the Points

The key question exists in many branches of science: is it possible to find a method of nonlinear equation solution without iterations of numerical methods [11]? This paper aims at giving the positive answer to this question. Method of Hurwitz – Radon Matrices (MHR), described in this paper, is computing points between two successive nodes for searching a root of the function. The curve or function in MHR method is parameterized for real number $\alpha \in [0;1]$ in the range of two successive interpolation nodes.

3.1. The Operator of Hurwitz-Radon

Adolf Hurwitz (1859-1919) and Johann Radon (1887-1956) published the papers about specific class of matrices in 1923, working on the problem of quadratic forms. Matrices $A_i, i = 1, 2, \dots, m$ satisfying

$$A_i A_k + A_k A_i = 0, A_i^2 = -I \text{ for } j \neq k; j, k = 1, 2, \dots, m$$

are called a family of Hurwitz - Radon matrices. A family of Hurwitz - Radon (HR) matrices has important features [12]: HR matrices are skew-symmetric ($A_i^T = -A_i$) and reverse matrices are easy to find ($A_i^{-1} = -A_i$). Only for dimension $N = 2, 4$ or 8 the family of HR matrices consists of $N - 1$ matrices. For $N = 2$ there is one matrix:

$$A_1 = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}.$$

For $N = 4$ there are three HR matrices with integer entries:

$$A_1 = \begin{bmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \end{bmatrix}, \quad A_2 = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{bmatrix}, \quad A_3 = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{bmatrix}.$$

For $N = 8$ we have seven HR matrices with elements $0, \pm 1$. So far HR matrices are applied in electronics [13]: in Space-Time Block Coding (STBC) and orthogonal design [14], also in signal processing [15] and Hamiltonian Neural Nets [16].

If one curve is described by a set of following points $\{(x_i, y_i), i = 1, 2, \dots, n\}$ then HR matrices combined with the identity matrix I_N are used to build the orthogonal and discrete Hurwitz - Radon Operator (OHR). For nodes (x_1, y_1) and (x_2, y_2) OHR M of dimension $N = 2$ is constructed:

$$M = \frac{1}{x_1^2 + x_2^2} \begin{bmatrix} x_1 y_1 + x_2 y_2 & x_2 y_1 - x_1 y_2 \\ x_1 y_2 - x_2 y_1 & x_1 y_1 + x_2 y_2 \end{bmatrix}. \quad (1)$$

For nodes $(x_1, y_1), (x_2, y_2), (x_3, y_3)$ and (x_4, y_4) OHR of dimension $N = 4$ is constructed:

$$M = \frac{1}{x_1^2 + x_2^2 + x_3^2 + x_4^2} \begin{bmatrix} u_0 & u_1 & u_2 & u_3 \\ -u_1 & u_0 & -u_3 & u_2 \\ -u_2 & u_3 & u_0 & -u_1 \\ -u_3 & -u_2 & u_1 & u_0 \end{bmatrix} \quad (2)$$

where

$$u_0 = x_1y_1 + x_2y_2 + x_3y_3 + x_4y_4, \quad u_1 = -x_1y_2 + x_2y_1 + x_3y_4 - x_4y_3, \\ u_2 = -x_1y_3 - x_2y_4 + x_3y_1 + x_4y_2, \quad u_3 = -x_1y_4 + x_2y_3 - x_3y_2 + x_4y_1.$$

For nodes $(x_1, y_1), (x_2, y_2), \dots$ and (x_8, y_8) OHR of dimension $N = 8$ is built [17] similarly as (1) and (2):

$$M = \frac{1}{\sum_{i=1}^8 x_i^2} \begin{bmatrix} u_0 & u_1 & u_2 & u_3 & u_4 & u_5 & u_6 & u_7 \\ -u_1 & u_0 & u_3 & -u_2 & u_5 & -u_4 & -u_7 & u_6 \\ -u_2 & -u_3 & u_0 & u_1 & u_6 & u_7 & -u_4 & -u_5 \\ -u_3 & u_2 & -u_1 & u_0 & u_7 & -u_6 & u_5 & -u_4 \\ -u_4 & -u_5 & -u_6 & -u_7 & u_0 & u_1 & u_2 & u_3 \\ -u_5 & u_4 & -u_7 & u_6 & -u_1 & u_0 & -u_3 & u_2 \\ -u_6 & u_7 & u_4 & -u_5 & -u_2 & u_3 & u_0 & -u_1 \\ -u_7 & -u_6 & u_5 & u_4 & -u_3 & -u_2 & u_1 & u_0 \end{bmatrix} \quad (3)$$

where

$$u = \begin{bmatrix} y_1 & y_2 & y_3 & y_4 & y_5 & y_6 & y_7 & y_8 \\ -y_2 & y_1 & -y_4 & y_3 & -y_6 & y_5 & y_8 & -y_7 \\ -y_3 & y_4 & y_1 & -y_2 & -y_7 & -y_8 & y_5 & y_6 \\ -y_4 & -y_3 & y_2 & y_1 & -y_8 & y_7 & -y_6 & y_5 \\ -y_5 & y_6 & y_7 & y_8 & y_1 & -y_2 & -y_3 & -y_4 \\ -y_6 & -y_5 & y_8 & -y_7 & y_2 & y_1 & y_4 & -y_3 \\ -y_7 & -y_8 & -y_5 & y_6 & y_3 & -y_4 & y_1 & y_2 \\ -y_8 & y_7 & -y_6 & -y_5 & y_4 & y_3 & -y_2 & y_1 \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \\ x_6 \\ x_7 \\ x_8 \end{bmatrix}. \quad (4)$$

The components of the vector $\mathbf{u} = (u_0, u_1, \dots, u_7)^T$, appearing in the matrix M (3), are defined by (4) in the similar way to (1)-(2) but in terms of the coordinates of the above 8 nodes. Note that OHR operators M (1)-(3) satisfy the condition of interpolation

$$M \cdot \mathbf{x} = \mathbf{y} \quad (5)$$

for $\mathbf{x} = (x_1, x_2, \dots, x_N)^T \in \mathbf{R}^N$, $\mathbf{x} \neq \mathbf{0}$, $\mathbf{y} = (y_1, y_2, \dots, y_N)^T \in \mathbf{R}^N$, $N = 2, 4$ or 8 .

3.2. The Method of Hurwitz-Radon Matrices

Key question looks as follows: how can we compute coordinates of points settled between the interpolation nodes [18]? The answer is connected with novel MHR method [19]. On a segment of a line every number “ c ” situated between “ a ” and “ b ” is described by a linear (convex) combination $c = \alpha \cdot a + (1 - \alpha) \cdot b$ for

$$\alpha = \frac{b - c}{b - a} \in [0; 1]. \quad (6)$$

The average OHR operator M_2 of dimension $N = 2, 4$ or 8 is constructed as follows:

$$M_2 = \alpha \cdot M_0 + (1 - \alpha) \cdot M_1 \quad (7)$$

with the operator M_0 built (1)-(3) by “odd” nodes $(x_1=a, y_1), (x_3, y_3), \dots, (x_{2N-1}, y_{2N-1})$ and M_1 built (1)-(3) by “even” nodes $(x_2=b, y_2), (x_4, y_4), \dots, (x_{2N}, y_{2N})$. Having the operator M_2 it is possible to reconstruct the second coordinates of points (x, y) in terms of the vector C defined with

$$c_i = \alpha \cdot x_{2i-1} + (1 - \alpha) \cdot x_{2i}, \quad i = 1, 2, \dots, N \quad (8)$$

as $C = [c_1, c_2, \dots, c_N]^T$. The required formula is similar to (5):

$$Y(C) = M_2 \cdot C \quad (9)$$

in which components of vector $Y(C)$ give the second coordinate of the points (x,y) corresponding to the first coordinate, given in terms of components of the vector C .

Calculations of unknown coordinates for curve points using (6)-(9) are called by author the method of Hurwitz - Radon Matrices (MHR) [20]. Here is the application of MHR method (Fig.2) for function $f(x) = x^3 + x^2 - x + 1$ with nodes as Fig.1 and computed 99 points between each pair of nodes ($\alpha = 0.01, 0.02, \dots, 0.99$). MHR-2 means MHR version with matrices of dimension $N = 2$.




Fig. 2. Function $f(x) = x^3 + x^2 - x + 1$ with 396 interpolated points using MHR-2 method with 5 nodes.

Solving the equation $x^3 + x^2 - x + 1 = 0$ via MHR interpolation, as it was said under Fig.1, we will search a root of the function only between nodes $(-2;-1)$ and $(-1.75;0.453125)$. Points calculated between other pairs of nodes are useless in the process of root approximation and they do not have to be computed. Considering points between nodes $(-2;-1)$ and $(-1.75;0.453125)$, coordinate y is near zero at $(-1.835;0.00184)$. Solution of equation $x^3 + x^2 - x + 1 = 0$ via MHR-2 method is approximated by $x = -1.835$. True value is $x = -1.839$.

The same equation for nodes $(-2;-1)$, $(-1.95;-0.662)$, $(-1.9;-0.349)$, $(-1.85;-0.059)$ and $(-1.8;0.208)$, solved by MHR-2 method, gives better result $x = -1.839$. So shorter distance between first and last node is of course very significant.

4. The Zero of Function via MHR method

Example 1

MHR calculations are done for function $f(x) = x^3 + \ln(7-x)$ with nodes: $(-2;-5.803)$, $(-1.75;-3.190)$, $(-1.5;-1.235)$, $(-1.25;0.1571)$ and $(-1;1.0794)$. So a root of this function is situated between 3rd and 4th node. MHR-2 interpolation gives the graph of function (Fig.3):




Fig. 3. Function $f(x) = x^3 + \ln(7-x)$ with 396 interpolated points using MHR-2 method with 5 nodes.

Considering points between nodes (-1.5;-1.235) and (-1.25;0.1571), coordinate y is near zero at (-1.2825;0.00194). Solution of equation $x^3 + \ln(7-x) = 0$ via MHR method is approximated by $x = -1.2825$. True value is hardly approximated (even for MathCad) by $x = -1.28347$.

Example 2

MHR calculations are done for function $f(x) = x^3 + 2x - 1$ with nodes: (0;-1), (0.25;-0.484), (0.5;0.125), (0.75;0.9219) and (1;2). So a zero of this function is situated between 2nd and 3rd node. MHR-2 interpolation gives the graph of function (Fig.4):




Fig. 4. Function $f(x) = x^3 + 2x - 1$ with 396 interpolated points using MHR-2 method with 5 nodes.

Considering points between nodes (0.25;-0.484) and (0.5;0.125), coordinate y is near zero at (0.4625;0.00219). Solution of equation $x^3 + 2x - 1 = 0$ via MHR-2 method is approximated by $x = 0.4625$. The only one real solution of this equation is $x = 0.453$.

Now MHR calculations are done for the same equation $x^3 + 2x - 1 = 0$ with seven nodes between (0;-1) and (1;2) for $x_i = 0; 0.125; 0.25; 0.375; 0.5; 0.625; 0.75; 0.875$ and 1. The solution is approximated by MHR-4 method with nine nodes. MHR-4 interpolation gives the graph of function (Fig.5):




Fig. 5. Function $f(x) = x^3 + 2x - 1$ with interpolated points using MHR-4 method with 9 nodes.

Considering points between nodes (0.375;-0.197) and (0.5;0.125), coordinate y is near zero at (0.45625;0.00018). Solution of equation $x^3 + 2x - 1 = 0$ via MHR-4 method is approximated by $x = 0.45625$. This is better result than MHR-2: greater number of nodes (with the same distance between first and last) means better approximation. And seventeen nodes in MHR-8 guarantee more precise results than MHR-4.

Example 3

MHR calculations are done for equation $3 - 2^x = 0$ with nodes: (1;1), (1.2;0.7026), (1.4;0.361), (1.6;-0.031) and (1.8;-0.482). MHR-2 interpolation gives the graph of function (Fig.6):




Fig. 6. Function $f(x) = 3 - 2^x$ with 396 interpolated points using MHR-2 method with 5 nodes.

Considering points between nodes (1.4;0.361) and (1.6;-0.031), second coordinate is near zero at (1.586;-0.000311). Solution of equation $3 - 2^x = 0$ via MHR-2 method is approximated by $x = 1.586$. Precise solution $x = \log_2 3$ is approximated by 1.585.

Interpolated values, calculated by MHR method, are applied in the process of solving the nonlinear equations. Shorter distance between first and last node or greater number of nodes guarantee better approximation. Approximated solutions of nonlinear equations are used in many branches of science. MHR joins two important problems in computer sciences: interpolation of the function with the solution of nonlinear equation.

5. Conclusions

The method of Hurwitz-Radon Matrices leads to curve interpolation [21] and approximation of nonlinear equation solution depending on the number and location of nodes. No characteristic features of function are important in MHR method: polynomial or not, monotonicity, convexity, derivative, starting point. These features are very significant for iterative numerical methods. MHR method gives the possibility of reconstruction a curve and searching for a root of the function. The only condition is to have a set of nodes according to assumptions in MHR method. Main features of MHR method are: accuracy of the equation solution depends on the number of nodes and the distance between first and last node (MHR-4 is more precise than MHR-2 and MHR-8 is more

precise than MHR-4); interpolation of a curve consists of L points is connected with the computational cost of rank $O(L)$; MHR is well-conditioned method (orthogonal matrices); MHR is dealing with local operators: average OHR operators are built by successive 4, 8 or 16 data points, what means smaller computational costs then using all nodes; MHR is not an affine interpolation [22].

Future works are connected with: possibility to apply MHR method to three-dimensional curves (3D data), computing the interpolation error, implementation of MHR in object recognition [23], MHR extrapolation method [24] and curve parameterization [25].

References

1. Brachman, R.J., Levesque, H.J.: Knowledge Representation and Reasoning. Morgan Kaufman, San Francisco (2004)
2. Yang, W.Y., Cao, W., Chung, T.-S., Morris, J.: Applied Numerical Methods Using MATLAB®. John Wiley & Sons, New York (2005)
3. Khalil, H.K.: Nonlinear Systems. Prentice Hall, New York (2001)
4. Kelley, C.T.: Iterative Methods for Linear and Nonlinear Equations. Society for Industrial and Applied Mathematics, Philadelphia (1995)
5. Markman, A.B.: Knowledge Representation. Lawrence Erlbaum Associates (1998)
6. Sowa, J.F.: Knowledge Representation: Logical, Philosophical and Computational Foundations. Brooks/Cole, New York (2000)
7. Soussen, C., Mohammad-Djafari, A.: Polygonal and Polyhedral Contour Reconstruction in Computed Tomography. IEEE Transactions on Image Processing 11(13), 1507-1523 (2004)
8. Tang, K.: Geometric Optimization Algorithms in Manufacturing. Computer – Aided Design & Applications 2(6), 747-757 (2005)
9. Kozera, R.: Curve Modeling via Interpolation Based on Multidimensional Reduced Data. Silesian University of Technology Press, Gliwice (2004)
10. Dahlquist, G., Björck, A.: Numerical Methods. Prentice Hall, New York (1974)
11. Ralston, A.: A First Course in Numerical Analysis. McGraw-Hill Book Company, New York (1965)
12. Eckmann, B.: Topology, Algebra, Analysis- Relations and Missing Links. Notices of the American Mathematical Society 5(46), 520-527 (1999)
13. Citko, W., Jakóbczak, D., Sieńko, W.: On Hurwitz - Radon Matrices Based Signal Processing. Workshop Signal Processing at Poznan University of Technology (2005)
14. Tarokh, V., Jafarkhani, H., Calderbank, R.: Space-Time Block Codes from Orthogonal Designs. IEEE Transactions on Information Theory 5(45), 1456-1467 (1999)
15. Sieńko, W., Citko, W., Wilamowski, B.: Hamiltonian Neural Nets as a Universal Signal Processor. 28th Annual Conference of the IEEE Industrial Electronics Society IECON (2002)
16. Sieńko, W., Citko, W.: Hamiltonian Neural Net Based Signal Processing. The International Conference on Signal and Electronic System ICSES (2002)
17. Jakóbczak, D.: 2D and 3D Image Modeling Using Hurwitz-Radon Matrices. Polish Journal of Environmental Studies 4A(16), 104-107 (2007)
18. Jakóbczak, D.: Shape Representation and Shape Coefficients via Method of Hurwitz-Radon Matrices. Lecture Notes in Computer Science 6374 (Computer Vision and Graphics: Proc. ICCVG 2010, Part I), Springer-Verlag Berlin Heidelberg, 411-419 (2010)
19. Jakóbczak, D.: Curve Interpolation Using Hurwitz-Radon Matrices. Polish Journal of Environmental Studies 3B(18), 126-130 (2009)
20. Jakóbczak, D.: Application of Hurwitz-Radon Matrices in Shape Representation. In: Banaszak, Z., Świć, A. (eds.) Applied Computer Science: Modelling of Production Processes 1(6), pp. 63-74. Lublin University of Technology Press, Lublin (2010)
21. Jakóbczak, D.: Object Modeling Using Method of Hurwitz-Radon Matrices of Rank k . In: Wolski, W., Borawski, M. (eds.) Computer Graphics: Selected Issues, pp. 79-90. University of Szczecin Press, Szczecin (2010)

22. Jakóbczak, D.: Implementation of Hurwitz-Radon Matrices in Shape Representation. In: Choraś, R.S. (ed.) *Advances in Intelligent and Soft Computing 84, Image Processing and Communications: Challenges 2*, pp. 39-50. Springer-Verlag, Berlin Heidelberg (2010)
23. Jakóbczak, D.: Object Recognition via Contour Points Reconstruction Using Hurwitz-Radon Matrices. In: Józefczyk, J., Orski, D. (eds.) *Knowledge-Based Intelligent System Advancements: Systemic and Cybernetic Approaches*, pp. 87-107. IGI Global, Hershey PA, USA (2011)
24. Jakóbczak, D.: Data Extrapolation and Decision Making via Method of Hurwitz-Radon Matrices. *Lecture Notes in Computer Science/LNAI 6922 (Computational Collective Intelligence, Technologies and Applications: Proc. ICCCI 2011, Part 1)*, Springer-Verlag Berlin Heidelberg, 173-182 (2011)
25. Jakóbczak, D.: Curve Parameterization and Curvature via Method of Hurwitz-Radon Matrices. *Image Processing & Communications- An International Journal* 1-2(16), 49-56 (2011)

Abstract

Mathematics need suitable methods to approximate a zero of the function. Coordinate x for $f(x)=0$ is crucial in a large number of calculations because each equation can be transformed into $f(x)=0$. A novel method of Hurwitz-Radon Matrices (MHR) can be used in approximation of a root of function in the plane. The paper contains a way of data approximation via MHR method to solve any equation. Proposed method is based on the family of Hurwitz-Radon (HR) matrices. The matrices are skew-symmetric and possess columns composed of orthogonal vectors. The operator of Hurwitz-Radon (OHR), built from these matrices, is described. Two-dimensional data are represented by discrete set of curve f points. It is shown how to create the orthogonal OHR operator and how to use it in a process of data interpolation. MHR method is interpolating the curve point by point without using any formula or function.

Streszczenie

Matematyka wymaga odpowiednich metod przybliżania miejsca zerowego funkcji. Współrzędna x w równaniu $f(x)=0$ jest kluczowa w wielu przypadkach, ponieważ dowolne równanie nieliniowe może zostać przedstawione jako $f(x)=0$. Nowa metoda Macierzy Hurwitza-Radona (MHR) może zostać użyta w rozwiązywaniu dowolnego równania z jedną niewiadomą. Artykuł zawiera sposób przybliżania pierwiastka funkcji. Metoda ta jest oparta na rodzinie macierzy Hurwitza-Radona (HR). Macierze HR są skośnie-symetryczne i składają się z kolumn tworzących ortogonalne wektory. W pracy pokazano jak konstruować Operator Hurwitza-Radona (OHR) oraz jak wykorzystać go w procesie rozwiązywania równania. Krzywa płaska opisana jest za pomocą punktów węzłowych. Metoda MHR interpoluje funkcję punkt po punkcie bez użycia wzoru opisującego krzywą.