

## ON MEAN-VALUE PROPERTIES FOR THE DUNKL POLYHARMONIC FUNCTIONS

Grzegorz Łysik

*Communicated by Semyon B. Yakubovich*

**Abstract.** We derive differential relations between the Dunkl spherical and solid means of continuous functions. Next we use the relations to give inductive proofs of mean-value properties for the Dunkl polyharmonic functions and their converses.

**Keywords:** Dunkl Laplacian, Dunkl polyharmonic functions, mean-values, Pizzetti formula.

**Mathematics Subject Classification:** 31A30, 31B30, 33C52.

### 1. INTRODUCTION

During the last few years there is a growing interest in the study of Dunkl harmonic functions, i.e., solutions to  $\Delta_\kappa u = 0$ , where  $\Delta_\kappa$  is a second order differential-difference operator invariant under the action of a discrete reflection group, see (2.1). The operator  $\Delta_\kappa$  was introduced by Dunkl in [2, 3], in the context of the theory of orthogonal polynomials in several variables. Afterwards the whole theory related to  $\Delta_\kappa$  was elaborated including analogues of Fourier analysis, special functions connected with root systems, algebraic approaches and an application to the solution of quantum Calogero-Sutherland models (see [5] for an excellent survey). In particular, Mejjali and Triméche proved in [12] that the operator  $\Delta_\kappa$  is hypoelliptic on  $\mathbb{R}^n$  and that smooth Dunkl harmonic functions on  $\mathbb{R}^n$  can be characterized by the Dunkl spherical mean value property. Furthermore, they derived a Pizzetti type formula for smooth functions on  $\mathbb{R}^n$ . Maslouhi and Yousfi solved in [10] the Dirichlet problem for  $\Delta_\kappa$  on the unit ball  $B$  and derived a characterization of  $C^2$  Dunkl harmonic functions on  $B$  by the Dunkl spherical mean value property. Recently, Hassine has obtained in [7] the characterization without smoothness assumptions. Maslouhi and Daher proved in [11] Weil's lemma for  $\Delta_\kappa$  and gave a characterization of Dunkl harmonic functions in a class of tempered distributions in terms of invariance under Dunkl convolution with suitable kernels. The Pizzetti series associated with  $\Delta_\kappa$  was studied by Salem and

Touahri in [15], they proved its convergence for a real analytic function and derived some Liouville type results for Dunkl polyharmonic functions. Some other results related to the Dunkl spherical mean value operator were also derived in [9, 14, 16].

In this paper we first derive differential relations between the Dunkl spherical and solid means of functions. Next, we use the relations to give a short proof of an analogue of the Beckenbach-Read theorem stating that equality of the Dunkl spherical and solid means of a continuous function implies its Dunkl harmonicity. Taking full advantage of the relations we also give simple inductive proofs of the Dunkl solid and spherical mean-value properties for the Dunkl polyharmonic functions and their converses in arbitrary dimension. The paper is a continuation of [8], where analogous results were obtained for polyharmonic functions.

## 2. PRELIMINARIES

Recall that for a nonzero vector  $\alpha \in \mathbb{R}^n \setminus \{0\}$  the reflection with respect to the orthogonal to the  $\alpha$  hyperplane  $H_\alpha$  is given by

$$\sigma_\alpha(x) = x - \frac{2\langle \alpha, x \rangle}{\|\alpha\|^2} \alpha, \quad x \in \mathbb{R}^n,$$

where  $\langle \cdot, \cdot \rangle$  is the euclidian scalar product on  $\mathbb{R}^n$  and  $\|\cdot\|$  the associated norm. A finite set  $R$  of nonzero vectors is called a root system if  $R \cap \mathbb{R}\alpha = \{\pm\alpha\}$  and  $\sigma_\alpha R = R$  for all  $\alpha \in R$ . The reflections  $\sigma_\alpha$  with  $\alpha$  in a given root system  $R$  generate a finite group  $W \subset O(n)$ , called the reflection group associated with  $R$ . For a fixed  $\beta \in \mathbb{R}^n \setminus \bigcup_{\alpha \in R} H_\alpha$  one can decompose  $R = R_+ \cup R_-$  where  $R_\pm = \{\alpha \in R : \pm\langle \alpha, \beta \rangle > 0\}$ ; vectors in  $R_+$  are called positive roots. A function  $\kappa : R \rightarrow \mathbb{R}$  is called a multiplicity function if it is invariant under the action of the associated reflection group  $W$ . Its index  $\gamma$  is defined by

$$\gamma = \sum_{\alpha \in R_+} \kappa(\alpha).$$

Throughout the paper we shall assume that  $\kappa \geq 0$  and  $\gamma > 0$ .

The Dunkl operators  $T_j$ ,  $j = 1, \dots, n$ , associated with a root system  $R$  and a multiplicity function  $\kappa$  were introduced by C. Dunkl [3] as

$$T_j f(x) = \frac{\partial}{\partial x_j} f(x) + \sum_{\alpha \in R_+} \kappa(\alpha) \frac{f(x) - f(\sigma_\alpha(x))}{\langle \alpha, x \rangle} \alpha_j \quad \text{for } f \in C^1(\mathbb{R}^n).$$

Clearly,  $T_j f$  is well defined for  $f \in C^1(\Omega)$  where  $\Omega$  is a  $W$ -invariant open subset of  $\mathbb{R}^n$  and it reduces to  $\frac{\partial}{\partial x_j} f$  if  $f$  is  $W$ -invariant. The Dunkl Laplacian  $\Delta_\kappa$  is defined as a sum of squares of the operators  $T_j$ ,  $j = 1, \dots, n$ , i.e.,

$$\Delta_\kappa f = \sum_{j=1}^n T_j^2 f \quad \text{for } f \in C^2(\Omega).$$

A simple computation leads to

$$\Delta_\kappa f(x) = \Delta f(x) + \sum_{\alpha \in R_+} \kappa(\alpha) \left( \frac{2\langle \nabla f(x), \alpha \rangle}{\langle \alpha, x \rangle} - \|\alpha\| \frac{f(x) - f(\sigma_\alpha(x))}{\langle \alpha, x \rangle^2} \right). \tag{2.1}$$

Here  $\Delta$  and  $\nabla$  denote the usual Laplacian and gradient, respectively.

The Dunkl intertwining operator  $V_\kappa$  acting on polynomials was defined in [4] by

$$T_j V_\kappa f = V_\kappa \frac{\partial}{\partial x_j} f \quad \text{for } j = 1, \dots, n \quad \text{and} \quad V_\kappa 1 = 1.$$

The operator  $V_\kappa$  extends to a topological isomorphism of  $C^\infty(\mathbb{R}^n)$  onto itself [17]. In general there is no explicit description of  $V_\kappa$ , but Rösler has shown [13, Th. 1.2, Cor. 5.3] that for any  $x \in \mathbb{R}^n$  there exists a unique probability measure  $\mu_x$  such that

$$V_\kappa f(x) = \int_{\mathbb{R}^n} f(y) d\mu_x(y). \tag{2.2}$$

Moreover, the support of  $\mu_x$  is contained in  $\text{ch}(Wx)$  – the convex hull of the set  $\{gx : g \in W\}$ ,  $\mu_{rx}(U) = \mu_x(r^{-1}U)$  and  $\mu_{gx}(U) = \mu_x(g^{-1}U)$  for  $r > 0, g \in W$  and a Borel set  $U \subset \mathbb{R}^n$ . Note that by (2.2),  $V_\kappa$  can be extended to continuous functions and  $|V_\kappa(f)(x)| \leq \sup_{y \in \text{ch}(Wx)} |f(y)|$ ; the extension is a topological isomorphism of  $C(\mathbb{R}^n)$ .

The Dunkl translation operators  $\tau_x, x \in \mathbb{R}^n$ , are defined on  $C(\mathbb{R}^n)$  by

$$\tau_x f(y) = (V_\kappa)_x (V_\kappa)_y [V_\kappa^{-1} f(x + y)] \quad \text{for } y \in \mathbb{R}^n.$$

A more suggestive notation  $f(x *_\kappa y) := \tau_x f(y)$  is also used. Note that  $\tau_0 f = f$  and  $\tau_y f(x) = \tau_x f(y)$  for  $x, y \in \mathbb{R}^n$ .

### 3. THE DUNKL MEAN VALUE PROPERTY

The Poisson kernel for the Dunkl Laplacian  $\Delta_\kappa$  is defined in [6] by<sup>1)</sup>

$$P_\kappa(x, y) = V_\kappa \left[ \frac{1 - \|x\|^2}{(1 - 2\langle x, \cdot \rangle + \|x\|)^{\gamma+n/2}} \right] (y) \quad \text{for } \|x\| < 1, \|y\| \leq 1.$$

The kernel  $P_\kappa(x, y)$  is non-negative, bounded by 1 and it has the reproducing property for Dunkl harmonic functions on the unit ball  $B = B(0, 1)$ . Furthermore it is used as a tool to solve the Dirichlet problem for the Dunkl Laplacian. Namely it holds

**Theorem 3.1** ([10, Theorem A, Prop. 2.1]). *Let  $u$  be a continuous function on the unit sphere  $S(0, 1)$ . Set*

$$P_\kappa[u](x) = \frac{1}{d_\kappa} \int_{S(0,1)} P_\kappa(x, y) u(y) \omega_\kappa(y) dS(y) \quad \text{for } \|x\| < 1,$$

<sup>1)</sup>  $P_\kappa(x, y) = P(h_\kappa^2; y, x)$  where  $P(h_\kappa^2; \cdot, \cdot)$  is defined in [6, p. 190].

where

$$d_\kappa = \int_{S(0,1)} \omega_\kappa(y) dS(y) \quad \text{and} \quad \omega_\kappa(y) = \prod_{\alpha \in R_+} |\langle \alpha, y \rangle|^{2\kappa(\alpha)}.$$

Then  $P_\kappa[u]$  is  $\Delta_\kappa$ -harmonic on the unit ball  $B$ , extends continuously to  $\overline{B}$  and  $P_\kappa[u] = u$  on  $S(0, 1)$ . Furthermore,  $P_\kappa[u]$  is the unique  $\Delta_\kappa$ -harmonic function on  $B$  which extends continuously to  $u$  on  $S(0, 1)$ .

Since  $P_\kappa(0, y) = 1$  for  $\|y\| \leq 1$  for a function  $u$  continuous on  $\overline{B}$  and Dunkl harmonic in  $B$  we get

$$u(0) = \frac{1}{d_\kappa} \int_{S(0,1)} u(y) \omega_\kappa(y) dS(y).$$

More generally, if a function  $u$  is continuous on  $\overline{B}$  and  $\Delta_\kappa$ -harmonic in  $B$ , then for any  $x \in B$  and  $0 < r < 1 - \|x\|$  the spherical mean value formula holds (see [10, Theorem C])

$$u(x) = \frac{1}{d_\kappa} \int_{S(0,1)} \tau_x u(ry) \omega_\kappa(y) dS(y) \tag{3.1}$$

The converse statement was also stated ([10, Theorem C]) under the assumption that  $u$  is a  $C^2$  function. Recently Hassine has proved it without that assumption.

**Theorem 3.2** ([7, Theorem 3.1]). *Let  $u$  be a bounded function on the closed unit ball  $\overline{B}$ . If for any  $x \in B$  and  $0 < r < 1 - \|x\|$  the formula (3.1) holds, then  $u$  is  $\Delta_\kappa$ -harmonic in  $B$ .*

#### 4. RELATIONS BETWEEN THE DUNKL SPHERICAL AND SOLID MEANS

Let  $u$  be a smooth function on the ball  $\overline{B}$ . For any  $x \in B$  and  $0 < R < 1 - \|x\|$  we denote by  $N^D(u; x, R)$  the Dunkl spherical integral mean of  $u$  over the sphere  $S(x, R)$ ,

$$N^D(u; x, R) = \frac{1}{d_\kappa} \int_{S(0,1)} \tau_x u(Ry) \omega_\kappa(y) dS(y). \tag{4.1}$$

It was proved in [14, Theorem 4.1] that the Dunkl spherical mean operator  $u \mapsto N^D(u; x, R)$  can be represented in the form

$$N^D(u; x, R) = \int_{\mathbb{R}^n} u(y) d\mu_{x,R}^\kappa(y),$$

where  $\mu_{x,R}^\kappa$  is a probability measure with support in  $\bigcup_{g \in W} \{y \in \mathbb{R}^n : \|y - gx\| \leq R\}$ . Hence  $N^D(u; x, R)$  is well defined for a continuous function  $u$ . Since  $\omega_\kappa$  is homogenous of degree  $2\gamma$ , we also have

$$N^D(u; x, R) = \frac{1}{d_\kappa R^{2\gamma+n-1}} \int_{S(0,R)} \tau_x u(z) \omega_\kappa(z) dS(z).$$

Note that using the spherical coordinates by homogeneity of  $\omega_\kappa$  we get

$$\int_{B(0,1)} \omega_\kappa(x) dx = \int_0^1 \left( \int_{S(0,1)} \omega_\kappa(y) dS(y) \right) t^{2\gamma+n-1} dt = \frac{d_\kappa}{2\gamma+n}.$$

So we can define the Dunkl solid integral mean of  $u$  over the ball  $\overline{B}(x, R)$  by

$$\begin{aligned} M^D(u; x, R) &= \frac{2\gamma+n}{d_\kappa} \int_{B(0,1)} \tau_x u(Ry) \omega_\kappa(y) dy \\ &= \frac{2\gamma+n}{d_\kappa R^{2\gamma+n}} \int_{B(0,R)} \tau_x u(z) \omega_\kappa(z) dz. \end{aligned} \tag{4.2}$$

For the convenience of the reader recall the Green formula for the Dunkl Laplacian.

**Theorem 4.1** (Green formula for  $\Delta_\kappa$ , [12, Theorem 4.11]). *Let  $\Omega$  be a bounded  $W$ -invariant regular open set in  $\mathbb{R}^n$  containing the origin and  $u \in C^2(\Omega)$ . Then for any closed ball  $\overline{B}(0, R) \subset \Omega$  it holds*

$$\int_{B(0,R)} \Delta_\kappa u(z) \omega_\kappa(z) dz = \int_{S(0,R)} \frac{\partial u(z)}{\partial \eta} \omega_\kappa(z) dS(z), \tag{4.3}$$

where  $\frac{\partial u}{\partial \eta}$  denotes the external normal derivative of  $u$ .

The relations between  $M^D(u; x, R)$  and  $N^D(u; x, R)$  are given in the following lemma.

**Lemma 4.2.** *Let  $u$  be a continuous function on the ball  $\overline{B}$ . Then for any  $x \in B$  and  $0 < R < 1 - \|x\|$  it holds*

$$\left( \frac{R}{2\gamma+n} \frac{\partial}{\partial R} + 1 \right) M^D(u; x, R) = N^D(u; x, R). \tag{4.4}$$

If we further assume that  $u$  has continuous derivatives up to second order, then

$$\frac{2\gamma+n}{R} \frac{\partial}{\partial R} N^D(u; x, R) = M^D(\Delta_\kappa u; x, R). \tag{4.5}$$

*Proof.* By (4.2) using the spherical coordinates, homogeneity of  $\omega_\kappa$  and (4.1) we compute

$$\begin{aligned} M^D(u; x, R) &= \frac{2\gamma+n}{d_\kappa R^{2\gamma+n}} \int_0^R \left( \int_{S(0,s)} \tau_x u(z) \omega_\kappa(z) dS(z) \right) ds \\ &= \frac{2\gamma+n}{R^{2\gamma+n}} \int_0^R N^D(u; x, s) s^{2\gamma+n-1} ds. \end{aligned}$$

Hence, by the Leibniz rule

$$\frac{\partial}{\partial R} M^D(u; x, R) = \frac{2\gamma + n}{R} \left( N^D(u; x, R) - M^D(u; x, R) \right),$$

which proves (4.4).

To show (4.5) we differentiate (4.1) under the integral sign to get

$$\begin{aligned} \frac{\partial}{\partial R} N^D(u; x, R) &= \frac{1}{d_\kappa} \int_{S(0,1)} \langle \nabla(\tau_x u)(Ry), y \rangle \omega_\kappa(y) dS(y) \\ &= \frac{1}{d_\kappa R^{2\gamma+n-1}} \int_{S(0,R)} \langle \nabla(\tau_x u)(z), \frac{z}{R} \rangle \omega_\kappa(z) dS(z). \end{aligned}$$

Note that the external normal vector to  $S(0, R)$  at a point  $z \in S(0, R)$  is  $\eta = \frac{z}{R}$  and  $\langle \nabla(\tau_x u), \eta \rangle = \frac{\partial(\tau_x u)}{\partial \eta}$ . So applying the Green formula (4.3) we get

$$\frac{\partial}{\partial R} N^D(u; x, R) = \frac{1}{d_\kappa R^{2\gamma+n-1}} \int_{B(0,R)} \Delta_\kappa(\tau_x u)(z) \omega_\kappa(z) dz = \frac{R}{2\gamma + n} M^D(\Delta_\kappa u; x, R),$$

since  $\Delta_\kappa \tau_x u(z) = \tau_x \Delta_\kappa u(z)$ , which implies (4.5). □

By (4.4) and (4.5), we obtain the following corollary.

**Corollary 4.3.** *Let  $u \in C^2(B)$ . Then for any  $x \in B$  and  $0 < R < 1 - \|x\|$  it holds that*

$$M^D(\Delta_\kappa u; x, R) = \left( \frac{\partial^2}{\partial R^2} + \frac{2\gamma + n + 1}{R} \frac{\partial}{\partial R} \right) M^D(u; x, R) \tag{4.6}$$

and

$$N^D(\Delta_\kappa u; x, R) = \left( \frac{\partial^2}{\partial R^2} + \frac{2\gamma + n - 1}{R} \frac{\partial}{\partial R} \right) N^D(u; x, R). \tag{4.7}$$

Let us point out that formula (4.7) was established in [12, Proposition 4.16].

By the first part of Lemma 4.2 we get an analogue of the Beckenbach-Read theorem ([1]) for the Dunkl harmonic functions.

**Corollary 4.4.** *Let  $u \in C^2(B)$ . If for any  $x \in B$  and  $0 < R < 1 - \|x\|$  it holds*

$$M^D(u; x, R) = N^D(u; x, R), \tag{4.8}$$

then  $u$  is Dunkl harmonic on  $B$ .

*Proof.* The assumption (4.8) and (4.4) imply that  $\frac{\partial}{\partial R} M^D(u; x, R) = 0$ . So for any  $x \in B$ ,  $M^D(u; x, R)$  is a constant equal to  $u(x)$  and the converse to the mean-value property for Dunkl harmonic functions ([10, Theorem C]) implies that  $u$  is Dunkl harmonic on  $B$ . □

5. MEAN-VALUE PROPERTIES FOR DUNKL POLYHARMONIC FUNCTIONS

Let  $m \in \mathbb{N}$ . A function  $u \in C^{2m}(\Omega)$  defined on a  $W$ -invariant open set  $\Omega \subset \mathbb{R}^n$  is called an  $m$ -Dunkl harmonic if it is a solution of the  $m$ -times iteration of the Dunkl operator, i.e.,  $\Delta_\kappa^m u = 0$ . One of the most trivial examples is given by an even power of the Euclidean distance from the origin.

**Example 5.1.** Let  $u(x) = r^{2m}(x)$  with  $m \in \mathbb{N}_0$ , where  $r(x) = (\sum_{i=1}^n x_i^2)^{1/2}$  is the radius function. Since  $u$  is  $W$ -invariant  $\Delta_\kappa u$  reduces to

$$\Delta_\kappa u(x) = \Delta u(x) + 2 \sum_{\alpha \in R_+} \kappa(\alpha) \frac{\langle \nabla u(x), \alpha \rangle}{\langle \alpha, x \rangle}.$$

Since  $\Delta u = 2m(n + 2m - 2)r^{2m-2}$  and  $\nabla u = 2mx \cdot r^{2m-2}$ , we get  $\Delta_\kappa u = 2m(n + 2m + 2\gamma - 2)r^{2m-2}$ . So  $u$  is  $(m + 1)$ -Dunkl harmonic,  $\Delta^i u(0) = 0$  for  $i = 0, 1, \dots, m - 1$  and

$$\begin{aligned} \Delta_\kappa^m u(0) &= 2m(2m - 2) \cdots 2 \times (n + 2m + 2\gamma - 2) \cdots (n + 2\gamma) r^0(0) \\ &= 4^m \left(\gamma + \frac{n}{2}\right)_m m!, \end{aligned}$$

where for  $a \in \mathbb{R}$ ,  $(a)_0 = 1$  and  $(a)_i = a(a + 1) \cdots (a + i - 1)$  for  $i \in \mathbb{N}$ . On the other hand using the spherical coordinates and the fact that  $\omega$  is homogeneous of degree  $2\gamma$  we get

$$\begin{aligned} M^D(u; 0, R) &= \frac{2\gamma + n}{d_\kappa R^{2\gamma+n}} \int_0^R \int_{S(0,s)} \|y\|^{2m} \omega_\kappa(y) dS(y) ds \\ &= \frac{2\gamma + n}{d_\kappa R^{2\gamma+n}} \int_0^R d_\kappa s^{2m+2\gamma+n-1} ds = \frac{2\gamma + n}{2m + 2\gamma + n} R^{2m}. \end{aligned} \tag{5.1}$$

Hence

$$M^D(u; 0, R) = \frac{\Delta_\kappa^m u(0)}{4^m \left(\gamma + \frac{n}{2} + 1\right)_m m!} \cdot R^{2m}.$$

The above example suggests a form of an expansion of  $M(u; x, R)$  for a polyharmonic function  $u$  into powers of the radius  $R$  of the ball  $B(x, R)$ .

**Theorem 5.2** (Mean-value property for solid means, [16, formula (1.1)]). *Let  $m \in \mathbb{N}_0$ . If  $u \in C^{2m+2}(B)$  and  $\Delta_\kappa^{m+1} u = 0$  in  $B$ , then for any  $x \in B$  and  $0 < R < 1 - \|x\|$  it holds*

$$M^D(u; x, R) = \sum_{k=0}^m \frac{\Delta_\kappa^k u(x)}{4^k \left(\gamma + \frac{n}{2} + 1\right)_k k!} \cdot R^{2k}. \tag{5.2}$$

*Proof.* It was pointed out in [16, p. 120] that the mean value formula (5.2) for solid means can be derived from an analogous one for spherical means by integration. Here

we give a proof based on a simple inductive arguments. Clearly, by the mean-value property for the Dunkl harmonic functions, which follows from [10, Theorem C], the formula (5.2) holds for  $m = 0$ . Inductively assume that Theorem 5.2 holds for a fixed  $m \in \mathbb{N}_0$ . Let  $v \in C^{2m+4}(B)$  and  $\Delta_\kappa^{m+2}v = 0$ . Then  $u = \Delta_\kappa v \in C^{2m+2}(B)$  satisfies  $\Delta_\kappa^{m+1}u = 0$  and so (5.2) holds. But, by (4.6),

$$\frac{2\gamma + n}{R} \frac{\partial}{\partial R} \left( \frac{R}{2\gamma + n} \frac{\partial}{\partial R} + 1 \right) M^D(v; x, R) = M^D(\Delta_\kappa v; x, R) = M^D(u; x, R).$$

So after one integration

$$\left( \frac{R}{2\gamma + n} \frac{\partial}{\partial R} + 1 \right) M^D(v; x, R) = \sum_{k=0}^m \frac{\Delta_\kappa^k u(x)}{4^k (2\gamma + n) (\gamma + \frac{n}{2} + 1)_k k!} \cdot \frac{R^{2k+2}}{2k + 2} + c. \tag{5.3}$$

Note that the general solution of  $\left( \frac{R}{2\gamma + n} \frac{\partial}{\partial R} + 1 \right) M^D(v; x, R) = 0$  is  $CR^{-2\gamma-n}$  and a particular solution of

$$\left( \frac{R}{2\gamma + n} \frac{\partial}{\partial R} + 1 \right) M^D(v; x, R) = \frac{\Delta_\kappa^k u(x)}{4^k (2\gamma + n) (\gamma + \frac{n}{2} + 1)_k k!} \cdot \frac{R^{2k+2}}{2k + 2}$$

is  $A_k R^{2k+2}$ , where

$$A_k \left( \frac{2k + 2}{2\gamma + n} + 1 \right) = \Delta_\kappa^k u(x) \cdot [4^k (2\gamma + n) (2k + 2) (\gamma + \frac{n}{2} + 1)_k k!]^{-1}.$$

So

$$\begin{aligned} A_k &= \frac{\Delta_\kappa^k u(x)}{4^k (2k + 2) (2\gamma + n + 2k + 2) (\gamma + \frac{n}{2} + 1)_k k!} \\ &= \frac{\Delta_\kappa^k u(x)}{4^{k+1} (\gamma + \frac{n}{2} + 1)_{k+1} (k + 1)!}. \end{aligned}$$

Hence, the general solution of (5.3) is

$$M^D(v; x, R) = CR^{-2\gamma-n} + \sum_{k=0}^m \frac{\Delta_\kappa^k u(x)}{4^{k+1} (\gamma + \frac{n}{2} + 1)_{k+1} (k + 1)!} \cdot R^{2k+2} + c.$$

Finally, note that  $\lim_{R \rightarrow 0} M^D(v; x, R) = v(x)$  and  $\lim_{R \rightarrow 0} R^{2\gamma+n} M^D(v; x, R) = 0$ . So  $c = v(x)$ ,  $C = 0$  and

$$\begin{aligned} M^D(v; x, R) &= v(x) + \sum_{k=0}^m \frac{\Delta_\kappa^{k+1} v(x)}{4^{k+1} (\gamma + \frac{n}{2} + 1)_{k+1} (k + 1)!} \cdot R^{2k+2} \\ &= \sum_{k=0}^{m+1} \frac{\Delta_\kappa^k v(x)}{4^k (\gamma + \frac{n}{2} + 1)_k k!} \cdot R^{2k} \end{aligned}$$

which proves Theorem 5.2 □



By Theorem 5.2 and the relation (4.4), we get the following corollary.

**Corollary 5.3** (Mean-value property for spherical means, [15, Proposition 3.1] and [12, Theorem 4.17]). *Under the assumptions of Theorem 5.2 for any  $x \in B$  and  $0 < R < 1 - \|x\|$  it holds*

$$N^D(u; x, R) = \sum_{k=0}^m \frac{\Delta_\kappa^k u(x)}{4^k (\gamma + \frac{n}{2})_k k!} \cdot R^{2k}. \tag{5.4}$$

**Theorem 5.4** (Converse to the mean value property for spherical means). *Let  $m \in \mathbb{N}_0$ . If  $u \in C^{2m}(B)$  and the formula (5.4) holds for any  $x \in B$  and  $0 < R < 1 - \|x\|$ , then  $\Delta_\kappa^{m+1}u = 0$  in  $B$ .*

*Proof.* Clearly, if  $m = 0$ , Theorem 5.4 follows from Theorem 3.2. Fix  $p \in \mathbb{N}$  and assume that Theorem 5.4 holds for  $m < p$ . We shall prove that it holds for  $m = p$ . To this end take  $v \in C^{2p}(B)$  and assume that for any  $x \in B$  and  $R$  small enough (5.4) holds with  $m = p$  and  $u = v$ . Set  $u = \Delta_\kappa v$ . Then  $u \in C^{2p-2}(B)$ . By (4.5) and (5.4) with  $m = p$  and  $u = v$ , we get

$$M^D(u; x, R) = \sum_{k=1}^p \frac{2k(2\gamma + n)\Delta_\kappa^k v(x)}{4^k (\gamma + \frac{n}{2})_k k!} \cdot R^{2k-2} = \sum_{k=0}^{p-1} \frac{\Delta_\kappa^k u(x)}{4^k (\gamma + \frac{n}{2} + 1)_k k!} \cdot R^{2k}.$$

So for any  $x \in B$  and  $R$  small enough, by (4.4) we derive

$$N^D(u; x, R) = \sum_{k=0}^{p-1} \left( \frac{2k}{2\gamma + n} + 1 \right) \frac{\Delta_\kappa^k u(x)}{4^k (\gamma + \frac{n}{2} + 1)_k k!} \cdot R^{2k} = \sum_{k=0}^{p-1} \frac{\Delta_\kappa^k u(x)}{4^k (\gamma + \frac{n}{2})_k k!} \cdot R^{2k}.$$

Hence, by the inductive assumption,  $\Delta_\kappa^p u = \Delta_\kappa^{p+1}v = 0$ . □

By Theorem 5.4 and the relation (4.4), we get the following corollary.

**Corollary 5.5** (Converse to the mean value property for solid means). *Under the assumptions of Theorem 5.2 if  $u \in C^{2m}(B)$  and for all  $x \in B$  and  $R$  small enough formula (5.2) holds, then  $\Delta_\kappa^{m+1}u = 0$  in  $B$ .*

REFERENCES

[1] E.F. Beckenbach, M. Reade, *Mean values and harmonic polynomials*, Trans. Amer. Math. Soc. **51** (1945), 240–245.  
 [2] C.F. Dunkl, *Reflection groups and orthogonal polynomials on the sphere*, Math. Z. **197** (1988), 33–60.  
 [3] C.F. Dunkl, *Differential-difference operators associated to reflection groups*, Trans. Amer. Math. Soc. **311** (1989), 167–183.  
 [4] C.F. Dunkl, *Operators commuting with Coxeter group action on polynomials*, [in:] Invariant Theory and Tableaux, IMA Vol. Math. Appl. **19**, Springer-Verlag, 1990, 107–117.

- [5] C.F. Dunkl, *Reflection groups in analysis and applications*, Japan J. Math. **3** (2008), 215–246.
- [6] C.F. Dunkl, Y. Xu, *Orthogonal Polynomials of Several Variables*, Cambridge Univ. Press, 2001.
- [7] K. Hassine, *Mean value property associated with the Dunkl Laplacian*, <http://arxiv.org/pdf/1401.1949.pdf>
- [8] G. Łysik, *On the mean-value property for polyharmonic functions*, Acta Math. Hungar. **133** (2011), 133–139.
- [9] M. Maslouhi, *On the generalized Poisson transform*, Integral Transforms Spec. Func. **20** (2009), 775–784.
- [10] M. Maslouhi, E.H. Youssfi, *Harmonic functions associated to Dunkl operators*, Monatsh. Math. **152** (2007), 337–345.
- [11] M. Maslouhi, R. Daher, *Weil’s lemma and converse mean value for Dunkl operators*, [in:] Operator Theory Adv. Math. **205**, Birkhäuser, 2009, 91–100.
- [12] H. Mejjaoli, K. Triméche, *On a mean value property associated with the Dunkl Laplacian operator and applications*, Integral Transforms Spec. Func. **12** (2001), 279–302.
- [13] M. Rösler, *Positivity of Dunkl’s intertwining operator*, Duke Math. J. **98** (1999), 445–463.
- [14] M. Rösler, *A positivity radial product formula for the Dunkl kernel*, Trans. Amer. Math. Soc. **355** (2003), 2413–2438.
- [15] N.B. Salem, K. Touahri, *Pizzetti series and polyharmonicity associated with the Dunkl Laplacian*, Mediterr. J. Math. **7** (2010), 455–470.
- [16] N.B. Salem, K. Touahri, *Cubature formulae associated with the Dunkl Laplacian*, Results Math. **58** (2010), 119–144.
- [17] K. Triméche, *The Dunkl intertwining operator on spaces of functions and distributions and integral representation of its dual*, Integral Transform Spec. Func. **12** (2001), 349–374.

Grzegorz Łysik  
lysik@impan.pl

Polish Academy of Sciences  
Institute of Mathematics  
Śniadeckich 8, 00-656 Warsaw, Poland

Jan Kochanowski University  
Faculty of Mathematics and Natural Sciences  
ul. Świętokrzyska 15, 25-406 Kielce, Poland

*Received: December 13, 2013.*

*Revised: July 4, 2014.*

*Accepted: July 8, 2014.*