

Jincai Wang

J-CONVEXITY CONSTANTS

Abstract. We introduce the J -convexity constants on Banach spaces and give some properties of the constants. We give the relations between the J -convexity constants and the n -th von Neumann-Jordan constants. Using the quantitative indices we estimate the value of J -convexity constants in Orlicz spaces.

Keywords: new quantitative index, J -convexity constants, n -th von Neumann-Jordan constants, Orlicz spaces.

Mathematics Subject Classification: 46B20, 46E30.

1. INTRODUCTION

Much of the significance of the concept of superreflexivity of a Banach space X is due to its numerous equivalent characterizations, see, e. g., Beauzamy [1, Part 4]. One of these characterizations is $J(n, \varepsilon)$ -convexity. We restate the definition from [2] as follows.

Definition 1.1. Given n and $0 < \varepsilon < 1$, we say that a Banach spaces X is $J(n, \varepsilon)$ -convex if for all elements $z_1, \dots, z_n \in U_X = \{x \in X : \|x\| \leq 1\}$ there is

$$\inf_{1 \leq k \leq n} \left\| \sum_{h=1}^k z_h - \sum_{h=k+1}^n z_h \right\| < n(1 - \varepsilon).$$

Definition 1.2. We define the J -convexity constants, for $n \geq 2$, by

$$J(n, X) = \sup \left\{ \inf_{1 \leq k \leq n} \left\| \sum_{n=1}^k z_h - \sum_{n=k+1}^n z_h \right\| : z_1, \dots, z_n \in U_X \right\}, \quad (1)$$

and

$$J_n(X) = \inf \left\{ \varepsilon : 0 < \varepsilon < 1, \text{ and there exists } z_1, \dots, z_n \in U_X \text{ such that} \right. \\ \left. \inf_{1 \leq k \leq n} \left\| \sum_{n=1}^k z_h - \sum_{n=k+1}^n z_h \right\| \geq n(1 - \varepsilon) \right\}. \quad (2)$$

It is known that a Banach space is superreflexive if and only if it is $J(n, \varepsilon)$ -convex for some n and $\varepsilon > 0$ ([3] and [4]). It is evident that:

- (i) $J(n, X) \leq n$ and $0 \leq J_n(X) < 1$ for $n \geq 2$.
- (ii) X is superreflexive if and only if $J_n(X) > 0$ for some n or, equivalently, $J(n, X) < n$ for some n .
- (iii) $J(n, X) = n(1 - \varepsilon)$ if and only if $J_n(X) = \varepsilon$.
- (iv) $J(n, X) = n$ if and only if $J_n(X) = 0$.
- (v) For a Banach spaces X the following conditions are equivalent:
 - 1) X is not superreflexive,
 - 2) $J_n(X) = 0$ for all $n \in \mathbb{N}$,
 - 3) $J(n, X) = n$ for all $n \in \mathbb{N}$.

Let

$$\Phi(u) = \int_0^{|u|} \phi(t)dt, \quad \Psi(v) = \int_0^{|v|} \psi(s)ds$$

be a pair of complementary N -functions, where the right derivative ϕ of Φ is right-continuous and nondecreasing, $\phi(t) > 0$ whenever $t > 0$, $\phi(0) = 0$ and $\lim_{t \rightarrow \infty} \phi(t) = \infty$; the right derivative ψ of Ψ satisfies the same conditions as ϕ . Assume that (G, Σ, m) is a (Lebesgue) measure space and $L^0(G, \Sigma, m)$ is the space of Σ -measurable functions defined on G . The Orlicz space is defined as

$$L^\Phi(G) = \{x \in L^0 : x \text{ is measurable in } G, \rho_\Phi(\lambda x)dt < \infty \text{ for some } \lambda > 0\},$$

where $\rho_\Phi(x) = \int_G \Phi(x(t))dt$. The Luxemburg norm (gauge norm) and the Orlicz norm in $L^\Phi(G)$ are defined, respectively, by

$$\|x\|_{(\Phi)} = \inf\{c > 0 : \rho_\Phi\left(\frac{x}{c}\right) \leq 1\}$$

and

$$\|x\|_\Phi = \inf_{k>0} \frac{1}{k} [1 + \rho_\Phi(kx)].$$

As usual, we denote $L^{(\Phi)} = (L^\Phi, \|\cdot\|_{(\Phi)})$, $L^\Phi = (L^\Phi, \|\cdot\|_\Phi)$ for short.

An N -function $\Phi(u)$ is said to satisfy the Δ_2 -condition for small u (for all u or for large u), which is written as $\Phi \in \Delta_2(0)$ ($\Phi \in \Delta_2$ or $\Phi \in \Delta_2(\infty)$), if there exist $u_0 > 0$ and $c > 0$ such that $\Phi(2u) \leq c\Phi(u)$ for $0 \leq u \leq u_0$ (for all $u \geq 0$ or for $u \geq u_0$). An N -function $\Phi(u)$ satisfies the ∇_2 -condition for small u (for all $u \geq 0$ or for large u), which is written as $\Phi \in \nabla_2(0)$ ($\Phi \in \nabla_2$ or $\Phi \in \nabla_2(\infty)$), if its complementary N -function (see [6] or [8]) $\Psi \in \Delta_2(0)$ ($\Psi \in \Delta_2$ or $\Psi \in \Delta_2(\infty)$). The basic facts on Orlicz spaces can be found in [8].

New quantitative indices for an N -function Φ are defined by

$$\begin{aligned} \alpha_\Phi(n) &= \liminf_{u \rightarrow \infty} \frac{\Phi^{-1}(u)}{\Phi^{-1}(nu)}, & \beta_\Phi(n) &= \limsup_{u \rightarrow \infty} \frac{\Phi^{-1}(u)}{\Phi^{-1}(nu)}, \\ \alpha_\Phi^0(n) &= \liminf_{u \rightarrow 0} \frac{\Phi^{-1}(u)}{\Phi^{-1}(nu)}, & \beta_\Phi^0(n) &= \limsup_{u \rightarrow 0} \frac{\Phi^{-1}(u)}{\Phi^{-1}(nu)}, \\ \bar{\alpha}_\Phi(n) &= \inf_{u > 0} \frac{\Phi^{-1}(u)}{\Phi^{-1}(nu)}, & \bar{\beta}_\Phi(n) &= \sup_{u > 0} \frac{\Phi^{-1}(u)}{\Phi^{-1}(nu)}. \end{aligned}$$

For $n = 2$, we denote these constants by $\alpha_\Phi, \beta_\Phi, \alpha_\Phi^0, \beta_\Phi^0, \bar{\alpha}_\Phi$ and $\bar{\beta}_\Phi$ (see [8]). Clearly, $\frac{1}{n} \leq \bar{\alpha}_\Phi(n) \leq \min\{\alpha_\Phi(n), \alpha_\Phi^0(n)\}$, $\max\{\beta_\Phi(n), \beta_\Phi^0(n)\} \leq \bar{\beta}_\Phi(n) \leq 1$.

Proposition 1.1. ([7]) *Let Φ be an N -function. Then:*

- (i) $\Phi \notin \Delta_2(\infty) \iff \beta_\Phi(n) = 1$; $\Phi \notin \nabla_2(\infty) \iff \alpha_\Phi(n) = \frac{1}{n}$.
- (ii) $\Phi \notin \Delta_2(0) \iff \beta_\Phi^0(n) = 1$; $\Phi \notin \nabla_2(0) \iff \alpha_\Phi^0(n) = \frac{1}{n}$.
- (iii) $\Phi \notin \Delta_2 \iff \bar{\beta}_\Phi(n) = 1$; $\Phi \notin \nabla_2 \iff \bar{\alpha}_\Phi(n) = \frac{1}{n}$.

The following results concern these new indices.

Proposition 1.2. ([7]) *Let Φ and Ψ be a pair of complementary N -functions and $n \geq 2$. Then:*

$$n\alpha_\Phi(n)\beta_\Psi(n) = 1 = n\alpha_\Psi(n)\beta_\Phi(n), \tag{3}$$

$$n\alpha_\Phi^0(n)\beta_\Psi^0(n) = 1 = n\alpha_\Psi^0(n)\beta_\Phi^0(n), \tag{4}$$

$$n\bar{\alpha}_\Phi(n)\bar{\beta}_\Psi(n) = 1 = n\bar{\alpha}_\Psi(n)\bar{\beta}_\Phi(n). \tag{5}$$

2. THE RELATIONS BETWEEN J -CONVEXITY CONSTANTS AND VON NEUMANN-JORDAN CONSTANTS

In order to discuss the J -convexity constants, we need the n -th von Neumann-Jordan constants defined as follows.

Definition 2.1. *We define the n -th von Neumann-Jordan constants, for $n \geq 2$, by*

$$C_{NJ}^{(n)}(X) = \sup \left\{ \frac{\sum_{k=1}^n \left\| \sum_{h=1}^k z_h - \sum_{h=k+1}^n z_h \right\|}{n \sum_{i=1}^n \|z_i\|^2} : z_i \in X, \sum_{i=1}^n \|z_i\|^2 \neq 0 \right\}.$$

When $n = 2$, $C_{NJ}^{(2)}(X)$ is the von Neumann-Jordan constants of a Banach space X (see [8]).

Theorem 2.1.

(i) For any Banach space X , there holds

$$J(n, X) \leq \sqrt{nC_{NJ}^{(n)}(X)}. \quad (6)$$

(ii) $J(n, X) < n$ if and only if $C_{NJ}^{(n)}(X) < n$.

Proof. (i) Let $x_1, x_2, \dots, x_n \in U(X)$. Then

$$\begin{aligned} n \min_{1 \leq i \leq n} \left\| \sum_{i=1}^k x_i - \sum_{i=k+1}^n x_i \right\|^2 &\leq \sum_{k=1}^n \left\| \sum_{i=1}^k x_i - \sum_{i=k+1}^n x_i \right\|^2 \leq \\ &\leq C_{NJ}^{(n)}(X) n \sum_{i=1}^n \|x_i\|^2 \leq \\ &\leq n^2 C_{NJ}^{(n)}(X). \end{aligned}$$

(ii) By (i), the sufficiency is clear. Now we prove the necessity.

$$\begin{aligned} C_{NJ}^{(n)}(X) &= \sup \left\{ \frac{\sum_{1 \leq k \leq n} \left\| \sum_{i=1}^k x_i - \sum_{i=k+1}^n x_i \right\|}{n \sum_{i=1}^n \|x_i\|^2} : \{x_i\}_1^n \subset X \text{ and } \sum_{i=1}^n \|x_i\|^2 \neq 0 \right\} = \\ &= \sup \left\{ \frac{\sum_{1 \leq k \leq n} \left\| \sum_{i=1}^k x_i - \sum_{i=k+1}^n x_i \right\|}{n \left(\sum_{i=1}^{n-1} \|x_i\|^2 + 1 \right)} : \{x_i\}_1^n \subset X, \|x_i\| \leq \|x_n\| = 1 \right\}. \end{aligned}$$

Since $J(n, X) < n$, there exists $0 < \delta < 1$ such that

$$\sup \left\{ \inf_{1 \leq k \leq n} \left\| \sum_{i=1}^k x_i - \sum_{i=k+1}^n x_i \right\| : \{x_i\}_1^n \subset U_X \right\} < n - \delta. \quad (7)$$

Suppose $1 = \|x_n\| \geq \|x_i\| > 1 - \frac{\delta}{2(n-1)}$ ($i = 1, 2, \dots, n-1$). By (7), there is

$$\inf_{1 \leq k \leq n} \left\| \sum_{i=1}^k x_i - \sum_{i=k+1}^n x_i \right\| < n - \delta.$$

Without loss of generality, we assume that

$$\|x_1 - x_2 - \dots - x_n\| = \inf_{1 \leq k \leq n} \left\| \sum_{i=1}^k x_i - \sum_{i=k+1}^n x_i \right\| < n - \delta.$$

Hence

$$\begin{aligned} \frac{\sum_{1 \leq k \leq n} \left\| \sum_{i=1}^k x_i - \sum_{i=k+1}^n x_i \right\|^2}{n \left(\sum_{i=1}^{n-1} \|x_i\|^2 + 1 \right)} &= \frac{\|x_1 - x_2 - \dots - x_n\|^2 + \sum_{2 \leq k \leq n} \left\| \sum_{i=1}^k x_i - \sum_{i=k+1}^n x_i \right\|^2}{n \left(\sum_{i=1}^{n-1} \|x_i\|^2 + 1 \right)} \leq \\ &\leq \frac{(n - \delta)^2}{n[(n - 1) \cdot (1 - \frac{\delta}{2(n-1)})^2 + 1]} + \frac{\sum_{2 \leq k \leq n} (\sum_{i=1}^{n-1} \|x_i\| + 1)^2}{n \left(\sum_{i=1}^{n-1} \|x_i\|^2 + 1 \right)} \leq \\ &\leq \frac{(n - \delta)^2}{n[(n - 1) - \delta + \frac{\delta^2}{4(n-1)} + 1]} + \frac{\sum_{2 \leq k \leq n} n \left(\sum_{i=1}^{n-1} \|x_i\|^2 + 1 \right)}{n \left(\sum_{i=1}^{n-1} \|x_i\|^2 + 1 \right)} = \\ &= \frac{(n - \delta)^2}{n[n - \delta + \frac{\delta^2}{4(n-1)}]} + n - 1 < \frac{n - \delta}{n} + n - 1 = n - \frac{\delta}{n}. \end{aligned}$$

If there exists $1 \leq i \leq n - 1$ such that $\|x_i\| \leq 1 - \frac{\delta}{2(n-1)}$, we may, without loss of generality, assume that $\|x_1\| \leq 1 - \frac{\delta}{2(n-1)}$. Then

$$\begin{aligned} \frac{\sum_{1 \leq k \leq n} \left\| \sum_{i=1}^k x_i - \sum_{i=k+1}^n x_i \right\|^2}{n \left(\sum_{i=1}^{n-1} \|x_i\|^2 + 1 \right)} &\leq \frac{n(\|x_1\| + \|x_2\| + \dots + \|x_{n-1}\| + 1)^2}{n \left(\sum_{i=1}^{n-1} \|x_i\|^2 + 1 \right)} = \\ &= n - \frac{n \left(\sum_{i=1}^{n-1} \|x_i\|^2 + 1 \right) - (\|x_1\| + \|x_2\| + \dots + \|x_{n-1}\| + 1)^2}{\left(\sum_{i=1}^{n-1} \|x_i\|^2 + 1 \right)} = \\ &= n - \left\{ n \left(\sum_{i=1}^{n-1} \|x_i\|^2 + 1 \right) - [\|x_1\|^2 + 2\|x_1\|(\|x_2\| + \dots + \|x_{n-1}\| + 1) + \right. \\ &\quad \left. + (\|x_2\| + \dots + \|x_{n-1}\| + 1)^2] \right\} / \left(\sum_{i=1}^{n-1} \|x_i\|^2 + 1 \right) = \\ &= n - \left\{ (1 - \|x_1\|)^2 + (\|x_1\| - \|x_2\|)^2 + \dots + (\|x_1\| - \|x_{n-1}\|)^2 + \right. \\ &\quad \left. + (n - 1) \left(\sum_{i=2}^{n-1} \|x_i\|^2 + 1 \right) - (\|x_2\| + \dots + \|x_{n-1}\| + 1)^2 \right\} / \left(\sum_{i=1}^{n-1} \|x_i\|^2 + 1 \right) \leq \\ &\leq n - \frac{(1 - \|x_1\|)^2}{\sum_{i=1}^{n-1} \|x_i\|^2 + 1} \leq n - \frac{[1 - (1 - \frac{\delta}{2(n-1)})]^2}{n} = n - \frac{\delta^2}{4n(n-1)^2}. \end{aligned}$$

Therefore,

$$C_{NJ}^{(n)}(X) \leq \max \left\{ n - \frac{\delta}{n}, n - \frac{\delta^2}{4n(n-1)^2} \right\} < n.$$

□

3. LOWER BOUNDS FOR J -CONVEXITY CONSTANTS IN ORLICZ SPACES

In this section, we will give lower bounds of J -convexity for three Orlicz spaces.

Theorem 3.1. *Let Φ be an N -function and $n \geq 2$. Then:*

$$\max \left[\frac{1}{\alpha_\Phi(n)}, n\beta_\Phi(n) \right] \leq J(n, L^{(\Phi)}[0, 1]), \tag{8}$$

$$\max \left[\frac{1}{\alpha_\Phi^0(n)}, n\beta_\Phi^0(n) \right] \leq J(n, l^{(\Phi)}), \tag{9}$$

$$\max \left[\frac{1}{\bar{\alpha}_\Phi(n)}, n\bar{\beta}_\Phi(n) \right] \leq J(n, L^{(\Phi)}[0, \infty)). \tag{10}$$

Proof. We only prove (8). The proofs of inequalities (9) and (10) are similar. By the definition of $\alpha_\Phi(n)$, there exists $0 < u_k \nearrow \infty$ such that

$$\lim_{k \rightarrow \infty} \frac{\Phi^{-1}(u_k)}{\Phi^{-1}(nu_k)} = \alpha_\Phi(n).$$

So for $\varepsilon > 0$, there exists $k_0 \geq 1$ such that for any $k \geq k_0$, there is

$$\frac{\Phi^{-1}(u_k)}{\Phi^{-1}(nu_k)} < \alpha_\Phi(n) + \varepsilon. \tag{11}$$

Put $u_0 = u_{k_0} > 1$. Let $G_i, 1 \leq i \leq n$ be non-overlapping subsets of $[0, 1]$, and $m(G_i) = \frac{1}{nu_0}, 1 \leq i \leq n$. Define $x_i(t) = \Phi^{-1}(nu_0)\chi_{G_i}(t), 1 \leq i \leq n$. Then $\|x_i\|_{(\Phi)} = 1$, and for any $1 \leq k \leq n$, there holds

$$\begin{aligned} \left\| \sum_{i=1}^k x_i - \sum_{k+1}^n x_i \right\|_{(\Phi)} &= \Phi^{-1}(nu_0) \|\chi_{G_1 \cup G_2 \cup \dots \cup G_n}\|_{(\Phi)} = \\ &= \frac{\Phi^{-1}(nu_0)}{\Phi^{-1}(u_0)} > \frac{1}{\alpha_\Phi(n) + \varepsilon}. \end{aligned}$$

Therefore

$$J(n, L^{(\Phi)}[0, 1]) > \frac{1}{\alpha_\Phi(n) + \varepsilon},$$

which proves (8), because ε is arbitrary.

Now we prove that $n\beta_\Phi(n) \leq J(n, L^{(\Phi)}[0, 1])$. By the definition of $\beta_\Phi(n)$, for any given $\varepsilon > 0$ we choose a $v_0 > 1$ such that $\frac{\Phi^{-1}(v_0)}{\Phi^{-1}(nv_0)} > \beta_\Phi(n) - \frac{\varepsilon}{n}$. We divide $[0, \frac{1}{v_0}]$ into n non-overlapping intervals A_1, A_2, \dots, A_n of the same length. Define:

$$\begin{aligned} x_1(t) &= \Phi^{-1}(v_0) (\chi_{A_1} + \chi_{A_2} + \chi_{A_3} + \dots + \chi_{A_n}), \\ x_2(t) &= \Phi^{-1}(v_0) (-\chi_{A_1} + \chi_{A_2} + \chi_{A_3} + \dots + \chi_{A_n}), \\ x_3(t) &= \Phi^{-1}(v_0) ((-\chi_{A_1} - \chi_{A_2} + \chi_{A_3} + \dots + \chi_{A_n})), \\ &\dots\dots \\ x_n(t) &= \Phi^{-1}(v_0) (-\chi_{A_1} - \chi_{A_2} - \chi_{A_3} - \dots - \chi_{A_{n-1}} + \chi_{A_n}). \end{aligned}$$

Obviously, $\|x_i\|_{(\Phi)} = 1 (i = 1, 2, \dots, n)$. For any $1 \leq k \leq n$,

$$\left\| \sum_{i=1}^k x_i - \sum_{i=k+1}^n x_i \right\|_{(\Phi)} \geq \|\Phi^{-1}(v_0)n\chi_{A_k}\|_{(\Phi)} = \frac{n\Phi^{-1}(v_0)}{\Phi^{-1}(nv_0)} \geq n\beta_\Phi(n) - \varepsilon.$$

The latter implies that $J(n, L^{(\Phi)}[0, 1]) \geq n\beta_\Phi(2^{n-1}) - \varepsilon$. This proves that $J_n(L^{(\Phi)}[0, 1]) \geq n\beta_\Phi(n)$, because ε is arbitrary. \square

Corollary 3.1.

- (i) If $\Phi \notin \Delta_2(\infty) \cap \nabla_2(\infty)$, then $J(n, L^{(\Phi)}[0, 1]) = n$.
- (ii) If $\Phi \notin \Delta_2(0) \cap \nabla_2(0)$, then $J(n, l^{(\Phi)}) = n$.
- (iii) If $\Phi \notin \Delta_2 \cap \nabla_2$, then $J(n, L^{(\Phi)}[0, \infty)) = n$.

Proof. We only prove (i). If $\Phi \notin \Delta_2(\infty)$, then $\beta_\Phi(n) = 1$ by Proposition 1.1. By (8), there is $J(n, L^{(\Phi)}[0, 1]) = n$. If $\Phi \notin \nabla_2(\infty)$, then $\alpha_\Phi(n) = \frac{1}{n}$. By (8), the same equality holds true. \square

Corollary 3.2. Let $1 < p < \infty, L^p \in \{L^p[0, 1], L^p[0, \infty), l^p\}$. Then for $n \geq 2$,

$$\begin{aligned} \max \left\{ n^{\frac{1}{p}}, n^{1-\frac{1}{p}} \right\} &\leq J(n, L^p), \\ \max \left\{ n^{\frac{2}{p}-1}, n^{1-\frac{2}{p}} \right\} &\leq C_{NJ}^{(n)}(L^p). \end{aligned}$$

Proof. We put $\Phi_p(u) = |u|^p$. Then $L^{(\Phi)} = L^p$ and $\|\cdot\|_{(\Phi)} = \|\cdot\|_p$. The result is easy to verify. \square

Similarly, for the Orlicz spaces with the Orlicz norm, the following theorem holds.

Theorem 3.2. Let Φ be an N -function, $n \geq 2$ and Ψ be a complementary N -function for Φ . Then:

$$\max \left[n\beta_\Psi(n), \frac{1}{\alpha_\Psi(n)} \right] \leq J(n, L^\Phi[0, 1]), \tag{12}$$

$$\max \left[n\beta_\Psi^0(n), \frac{1}{\alpha_\Psi^0(n)} \right] \leq J(n, l^\Phi), \tag{13}$$

$$\max \left[n\bar{\beta}_\Psi(n), \frac{1}{\bar{\alpha}_\Psi(n)} \right] \leq J(n, L^\Phi[0, \infty)). \tag{14}$$

Remark. (i) By Proposition 1.2, there is

$$\max \left\{ \frac{1}{\alpha_{\Phi}(n)}, n\beta_{\Phi}(n) \right\} = \max \left\{ n\beta_{\Psi}(n), \frac{1}{\alpha_{\Psi}(n)} \right\}.$$

That is to say that $J(n, L^{(\Phi)}[0, 1])$ and $J(n, L^{(\Psi)}[0, 1])$ have the same lower bounds etc.

(ii) If X_{Φ} denotes one of the Orlicz spaces in Theorem 3.1 and Theorem 3.2, then

$$\sqrt{n} \leq J(n, X_{\Phi}).$$

In fact, assume that $X_{\Phi} = L^{(\Phi)}$, then by Theorem 3.1, there holds

$$\sqrt{n} \leq \max \left\{ \frac{1}{\alpha_{\Phi}(n)}, n\beta_{\Phi}(n) \right\} \leq J(n, X_{\Phi}).$$

REFERENCES

- [1] B. Beauzamy, *Introduction to Banach spaces and their geometry*, Volume 68 of North-Holland Mathematics Studies, North-Holland, 2nd ed., 1985.
- [2] J. Wenzel, *Superreflexivity and J -convexity of Banach spaces*, Acta Math. Univ. Comenianae, Vol. LXVI, 1 (1997), 135–147.
- [3] R.C. James, *Some self dual properites of normed linear spaces*, Symposium on infinite dimensional topology, Baton Rouge, 1967, Volume 69 of Annals of Mathematics studies, 1972, 159–175.
- [4] J.J. Schäfler, K. Sundaresan, *Reflexivity and girth of spheres*, Math. Ann. 184 (1970) 3, 163–168.
- [5] Z.D. Ren, *Non-square constants of Orlicz space*, Stochastic processes and Function Analysis, edited by J.A. Goldstein, N. E. Gretskey and J. J. Uhl, Jr., Marcel Dekker, Lecture Notes in Pure and Applied Math. 186 (1997), 79–197.
- [6] S.T. Chen, *Geometry of Orlicz Spaces*, Dissertationes Math. 356 (1996), 1–204.
- [7] J. Wang, *On non- $l_n^{(1)}$ constants and n -th Von Neumann-Jordan constants for Orlicz spaces*, Commentationes Mathematicae, to appear.
- [8] M.M. Rao, Z.D. Ren, *Theory of Orlicz spaces*, Marcel Dekker, New York, 1991.

Jincai Wang
www.wangj@163.com

Suzhou University
Department of Mathematics
P.R. China, 215006

Received: December 29, 2005.