

Natalia Florencka*

The Content of Heavy Metals in the Alluvial Formations of Mountain Torrent**

1. Introduction

Base deposit gathered in the bottom of rivers and water basins as a result of mineral and organic material sedimentation retains different harmful substances containing heavy metals and durable organic compounds. Some compounds could be return ad to the water in the result of chemical and biochemical processes going on in sediments (it determines the caution to the living organisms), as well as in the result of mechanical movement of earlier deposited contaminated sediments. [2, 3, 10, 16, 22, 26]. Contaminated sediments could be transported to the soil of flood terrace or down with the river during water rising or flooding. Accumulative properties of deposit cause several times higher concentration of contaminants than those in the water. Quality of bed sediments of rivers is the factor of the anthropogenic pollution, as well gives information on potential threads for the ecosystem [9, 15, 17, 21, 23, 25]

The aim of the investigations was description of heavy metal content in the alluvial deposits gathered in the sandbanks of channels of Jalowiecki torrent, with consideration of amount of deposit fraction.

2. Methodic

The object of investigations was part of the basin of Jalowiecki torrent, placed in the grounds of Zawoja district, in the west part of Carpatian Mountains, in the Beskid Makowski region. Basin is limited in the North-West direction by the Jalowieckie mountain range and in the South-East direction by the Babiogórskie

* AGH University of Science and Technology, Faculty of Mining Surveying and Environmental Engineering, Krakow, Poland

** The work was carried out under financial support of Statutory Research AGH-UST no. 11.11.150.008

mountain range. Investigated stream is the mountain torrent with surface area of the basin of 19.33 km². Lower part of the torrent basin is built with magura sandstones, more often medium- and fine-grained, which together with quartz contains feldspar, slivers of dark phyllite shist, chlorite-sericite shist and muscovite. It comes from higher part of middle and upper eocene. The bottom of the valley is built with quaternary formation and gravels forming the terraces. It is covered not-bold gravels formed contemporary in the stream channel. The age of the gravel terraces is dated to the early post-glacial period, probably [18].

Intensive transport in the torrent channel of the river rubble is going on, it is gradually sorted and breaking up to smaller grains and its accumulation leads creating a visible channel formation. This formations are less durable, and in the dependence of value of the flow undergo to dynamic changes. It influences directly on torrent flow increasing its roughness and resistance. In the medium part of Jalowiecki torrent, in the longitude of 1.1 km three channel form appears. Meander sandbank, lateral "A", is formed in the higher part of measuring length on the right bank. Next formation is lateral sandbank "B" characterized by the significant size, its length is approx. 30 m. Last formation is lateral sandbank "C" of 12 m length, which state the bank protection of the torrent right side and despite visible erosion of bank slope is particularly stable.

Investigated material is taken from layer down to 20 cm, in the several points of each sandbank. Points are localized along sandbanks proportionally to its length. Fractions of >6 mm, 6–2 mm, 2–1 mm, 1–0.5 mm, 0.5–0.25 mm, 0.25–0.1 mm, 0.1–0.05 mm, 0.05–0.025, <0.025 mm were derived from the fetched material after drying. In the grains of size under 2 mm content of cadmium, copper, nickel, lead, chromium and zinc were determined by the atomic absorption spectrophotometer method.

3. Results

Sediment contained in river-bed sandbanks is characterized by changeable graining and different content of analyzed heavy metals. Sandbank A is very strong and highly skeletal built, sandbank B is highly skeletal, and sandbank C – medium skeletal [20]. Detailed presentation of grain size distribution of investigated samples is presented on figure 1.

Results of the chemical analysis of sediment samples are presented in the table 1. Content of cadmium oscillate in the range 0.01–0.95 mg·kg⁻¹, but the lowest values are stated in the fraction 1–0.5 mm. Lead contains in the range of 8.2–98.50 mg·kg⁻¹ and maximal values are denoted in the finest fractions <0.025 mm. Amount of chromium contains in the range from 21.10 mg·kg⁻¹ to

144.80 mg·kg⁻¹. Concentration of the chromium decreases gradually reaching minimal values in the fractions 0.5–0.25 mm and consecutively increases reaching the highest value for diameter <0.025 mm. Copper content contains in the range of 12.3–62.5 mg·kg⁻¹. The lowest amount of this metal was stated in the fraction 0.5–0.25 mm, and highest one for the grains diameter <0.025 mm. Manganese concentration oscillates in the wide range from 142 mg·kg⁻¹ to 1233 mg·kg⁻¹, the highest one are denoted in the fraction 0.05–0.025 mm. The amount of nickel is in the range 14.3–222.4 mg·kg⁻¹. This metal was highly cumulated in the fraction <0.025 mm. Zinc level oscillates from 33 mg·kg⁻¹ to 149.6 mg·kg⁻¹, the lowest amounts are denoted in the fraction 0.5–0.025 mm, and highest one in the grains <0.025 mm.

Investigated sediments contains trace elements with the different amount. The highest concentration of copper, chromium, nickel, lead, cadmium and zinc are denoted in the finest fraction (Fig. 2). Similar tendencies presented in the publications of different authors [1, 11, 13, 15, 24, 25].

Deposit sieved by the sieve of mesh diameter 2 mm should be taken into account under the analysis of taken material in the quality point of view [12]. Exceeded content of nickel in the 50% of samples are stated. Over normative amount appears mostly in the fraction <0.1 mm and sporadically in some samples of higher mesh diameter. Remain elements concentration is significantly under official regulations.




Fig. 1. Percentage contribution of derived fractions in the investigated river-bed sandbanks

Table 1. The content of chosen heavy metals in investigated fractions of sediments

Element		Fraction [mm]						
		2.00–1.00	1.00–0.50	0.50–0.25	0.25–0.01	0.10–0.05	0.05–0.025	<0.025
Chromium [mg·kg ⁻¹]	Average*	32.22	29.68	25.75	49.52	58.52	53.95	103.21
	Stand.Dev.	5.36	8.12	3.76	15.02	12.20	16.94	29.20
	Quartile 25%	29.93	25.00	23.75	38.65	50.35	41.40	76.55
	Quartile 75%	34.50	33.10	27.00	56.60	62.50	66.55	132.50
Manganese [mg·kg ⁻¹]	Average	559.6	635.0	570.8	648.1	818.0	948.6	698.5
	Stand.Dev.	283.4	196.9	85.9	59.1	47.6	69.9	293.0
	Quartile 25%	451.5	527.0	521.5	614.5	795.5	928.5	491.0
	Quartile 75%	612.5	718.5	598.0	682.5	827.0	974.0	971.0
Nickel [mg·kg ⁻¹]	Average	70.00	59.90	60.68	60.17	78.60	101.10	130.20
	Stand.Dev.	24.87	19.01	9.38	7.54	15.52	13.16	67.69
	Quartile 25%	67.15	53.00	59.50	54.65	65.20	90.30	69.43
	Quartile 75%	82.85	74.40	63.70	66.00	91.50	112.45	199.50
Copper [mg·kg ⁻¹]	Average	24.71	23.88	20.18	21.14	29.15	33.61	43.30
	Stand.Dev.	5.44	4.63	1.96	2.51	2.96	8.30	10.83
	Quartile 25%	22.38	23.25	19.45	19.45	26.70	32.40	38.70
	Quartile 75%	28.60	26.45	21.95	22.95	31.65	37.05	51.00
Zinc [mg·kg ⁻¹]	Average	65.63	67.11	55.44	62.14	90.14	108.8	119.72
	Stand.Dev.	8.34	15.16	5.32	6.34	6.96	7.88	15.92
	Quartile 25%	65.78	59.75	52.25	58.65	84.95	103.40	108.50
	Quartile 75%	70.65	77.10	60.05	64.25	93.15	111.55	130.60
Cadmium [mg·kg ⁻¹]	Average	0.32	0.27	0.28	0.36	0.41	0.44	0.46
	Stand.Dev.	0.19	0.13	0.26	0.30	0.29	0.24	0.19
	Quartile 25%	0.15	0.23	0.15	0.20	0.23	0.25	0.30
	Quartile 75%	0.44	0.35	0.35	0.45	0.68	0.58	0.60
Lead [mg·kg ⁻¹]	Average	20.54	18.96	23.01	16.05	20.15	30.61	52.96
	Stand.Dev.	9.74	14.05	25.42	3.45	2.58	18.53	11.08
	Quartile 25%	12.73	12.75	11.85	13.35	18.30	23.90	48.20
	Quartile 75%	22.80	18.60	17.05	18.45	21.80	26.90	56.90

* Average content of each metal was obtained on the basis of 11 samples.




Fig. 2. Average content of chosen heavy metals in the several sediment fractions

National Geological Institute in Poland apply quality classification of bottom sediments based on trace elements content in the fraction ≤ 0.2 mm [3, 5–8]. This fraction represent the content of the heavy metals in the environment in view of the adhesive properties point. Finer fractions are characterized by higher content of of heavy metal in comparison to thicker fraction of the same sample [7]. In the geochemical researches of water sediments incomplete distribution of samples is applied, because for the estimation of the pollution level of the environment the amount of an element which easily moves to the surface water is interested. Total content of an element in the deposit is less essential because remains of its content constitutes the mineral resistant to decay is not important for quality of the environment [4]

Established quality classification of bottom deposits describes threshold values taking lethal influence of accumulated pollutants on water organisms into account (I class – sediments weakly polluted, II class – sediments averagely polluted, III class – sediments polluted). Exceeding of at least one lethal component defines the quality of the deposit. Concentration of investigated metals significantly exceeds values of geochemical background. Especially amounts of nickel are on the high level (all samples of diameter < 0.025 mm), it cause the assignment of the deposit to the III class (Fig. 3).

Several sandbanks not differ each-other by the concentration of investigated metals, only approximately 10% lower content of lead and nickel is stated in the sandbank C in comparison to the others.




Fig. 3. Percentage contribution of metals in the investigated deposits according to geochemical classification

Source: [6]

4. Conclusions

1. Sediment collected in the River-bed sandbanks of Jalowiecki torrent divers in respect of graining from formation of highly skeletal to medium skeletal structure.
2. Content of analyzed heavy metals in the alluvial materials oscillates in the wide range. The highest content of chromium, copper, cadmium, nickel, lead and zinc are denoted in the finest fractions (<0.025 mm), while the lowest amount of zinc, chromium and copper contains an alluvial materials with diameter 0.5–0.25 mm.
3. Studied base materials are assigned to the III class, accordingly to the geochemical criteria, in the effect of exceeded concentration of Ni.

References

- [1] Aleksander-Kwaterczak U., Sikora W.S., Wójcik R.: *Rozkład zawartości metali ciężkich pomiędzy frakcje ziarnowe w osadach dennych rzeki Odry*. Geologia, t. 30, z. 2, 2004, pp. 165–174.
- [2] Belzile N., Chen Y., Gunn J.M., Dixit S.S.: *Sediment trace profiles in lakes of Killarney Park, Canada: from regional to continental influence*. Environmental Pollution, vol. 130, 2004, pp. 239–248.
- [3] Bojakowska I.: *Kryteria oceny zanieczyszczenia osadów wodnych*. Przegląd Geologiczny, vol. 49, nr 3, 2001, pp. 213–218.

-
- [4] Bojakowska I., Gliwicz T., Małecka K.: *Wyniki geochemicznych badań osadów wodnych Polski w latach 2003–2005*. BMŚ, Warszawa 2006.
- [5] Bojakowska I., Gliwicz T., Sokołowska G.: *Wyniki monitoringu geochemicznego osadów wodnych Polski w latach 1996 i 1997*. BMŚ, PIOŚ, Warszawa 1998.
- [6] Bojakowska I., Sokołowska G.: *Geochemiczne klasy czystości osadów wodnych*. Przegląd Geologiczny, vol. 46, nr 1, 1998, pp. 49–54.
- [7] Bojakowska I., Sokołowska G.: *Wyniki monitoringu geochemicznego osadów wodnych Polski w latach 1994–1995*. BMŚ, PIOŚ, Warszawa 1996.
- [8] Bojakowska I., Sokołowska G.: *Wyniki monitoringu geochemicznego osadów wodnych Polski w latach 1993–1993*. BMŚ, PIOŚ, Warszawa 1994.
- [9] Calmano W., Hong J., Förster U.: *Binding and mobilization of heavy metals in contaminated sediments affected by pH and redox potential*. Water Science and Technology, vol. 28, 1994, pp. 223–235.
- [10] Cieślęwicz J., Róžański S.: *Metale ciężkie w osadach dennych jezior*. Ochrona Środowiska i Zasobów Naturalnych, nr 44, 2010, pp. 72–82.
- [11] Ciszewski D., Malik I., Wardas M.: *Uwarunkowania geomorfologiczne migracji metali ciężkich w osadach fluwialnych: dolina Małej Panwi*. Przegląd Geologiczny, vol. 52, nr 2, 2004, pp. 163–174.
- [12] *Rozporządzenie Ministra Środowiska z dnia 16 kwietnia 2002 r. w sprawie rodzajów oraz stężeń substancji powodujących, że urobek jest zanieczyszczony*. Dz. U. Nr 55, poz. 498.
- [13] Frankowski M., Ziōła-Frankowska A., Kowalski A., Siepak J.: *Fractionation of heavy metals in bottom sediments using Tessier procedure*. Environmental Earth Sciences, vol. 60, no. 6, 2010, pp. 1165–1178.
- [14] Helios-Rybicka E.: *Rola minerałów ilastych w wiązaniu metali ciężkich przez osady rzeczne górnej Wisły*. ZN AGH nr 1050, Geologia, z. 32, 1986.
- [15] Helios-Rybicka E.: *Akumulacja i mobilizacja metali ciężkich w osadach środowiska wodnego: osady datowane jako wskaźnik chronologiczny*. Mat. Konf. "Geologiczne aspekty ochrony środowiska", Kraków 1991, pp. 18–23.
- [16] Hudson-Edwards K., Macklin M.G., Curtis C.D., Vaughan D.J.: *Chemical remobilization of contaminant metals within floodplain sediments in an incising river system: implications for dating and chemostratigraphy*. Earth Surface Processes and Land Forms, vol. 123, 1998, pp. 671–684.
- [17] Kabata-Pendias A., Pendias H.: *Biogeochemia pierwiastków śladowych*. Wyd. Nauk. PWN, Warszawa 1999.
- [18] Książkiewicz M.: *Geologia dynamiczna*. Wydawnictwa Geologiczne, Warszawa 1972.
- [19] Lis J., Pasieczna A.: *Atlas geochemiczny Polski w skali 1:2 500 000*. Państwowy Instytut Geologiczny, Warszawa 1995.

-
- [20] *Klasyfikacja uziarnienia gleb i utworów mineralnych – PTG 2008*. Roczniki Gleboznawcze, t. 60, z. 2, 2009, pp. 5–16.
- [21] Radecki-Pawlik A., Florencka N.: *Odkładanie wybranych pierwiastków chemicznych w łachach korytowych potoku górskiego pod wpływem antropopresji*. II Ogólnopolskie Sympozjum “Dynamika zmian środowiska geograficznego pod wpływem antropopresji”, Kraków – Sanok 25–27 września 2003.
- [22] Sobczyński T., Siepak J.: *Badanie kumulacji związków biogenicznych i specjacji metali w osadach dennych jezior Wielkopolskiego Parku Narodowego*. Ogólnopolska Konferencja Naukowa “Kompleksowe i szczegółowe problemy inżynierii środowiska”, Ustronie Morskie 2001, pp. 265–290.
- [23] Tam N.F.Y., Wong Y.S.: *Spatial variation of heavy metals in surface sediments of Hong Kong mangrove swamps*. Environmental Pollution, vol. 110, 2000, pp. 195–205.
- [24] Woszczyk M., Spychalski W.: *Czynniki czasowej zmienności zawartości wybranych metali ciężkich w osadach jeziora Sarbsko (Nizina Gardnieńsko-Łebska) na tle genezy zbiornika*. Ochrona Środowiska i Zasobów Naturalnych, nr 31, 2007, pp. 485–497.
- [25] Zioła A., Frankowski M., Sobczyński T., Siepak J.: *Wpływ uziarnienia na zawartość metali ciężkich w aluwialnych osadach rzeki Odry*. Nowoczesne metody przygotowania próbek i oznaczania śladowych ilości pierwiastków. Materiały XV Poznańskiego Konwersatorium Analitycznego, Poznań 2006.
- [26] Zoumis T., Schmidt A., Grigorova L. & Calmano W.: *Contaminants in sediments: remobilisation and demobilisation*. Science of the Total Environment, vol. 266, 2001, pp. 195–202.