

Chong Li, Józef Myjak

**POROUS SETS FOR MUTUALLY NEAREST POINTS
IN BANACH SPACES**

Abstract. Let $\mathfrak{B}(X)$ denote the family of all nonempty closed bounded subsets of a real Banach space X , endowed with the Hausdorff metric. For $E, F \in \mathfrak{B}(X)$ we set $\lambda_{EF} = \inf \{\|z - x\| : x \in E, z \in F\}$. Let \mathfrak{D} denote the closure (under the maximum distance) of the set of all $(E, F) \in \mathfrak{B}(X) \times \mathfrak{B}(X)$ such that $\lambda_{EF} > 0$. It is proved that the set of all $(E, F) \in \mathfrak{D}$ for which the minimization problem $\min_{x \in E, z \in F} \|x - z\|$ fails to be well posed in a σ -porous subset of \mathfrak{D} .

Keywords: minimization problem, well-posedness, H_ρ -topology, σ -porous set.

Mathematics Subject Classification: Primary 41A65, 54E52; Secondary 46B20.

1. INTRODUCTION

Let X be a real Banach space. By $\mathfrak{B}(X)$ we denote the family of all nonempty closed bounded subsets of X . For $E, F \in \mathfrak{B}(X)$ we set

$$\lambda_{EF} := \inf \{\|z - x\| : x \in E, z \in F\}.$$

We consider the minimization problem, denoted by $\min(E, F)$, of finding a pair (x_0, z_0) with $x_0 \in E, z_0 \in F$ such that $\|x_0 - z_0\| = \lambda_{EF}$. Such a pair is called a solution of the minimization problem $\min(E, F)$. Moreover, any sequence $\{(x_n, z_n)\}$ with $x_n \in E, z_n \in F$ such that $\lim_{n \rightarrow \infty} \|x_n - z_n\| = \lambda_{EF}$ is called a *minimizing* sequence for the problem $\min(E, F)$. A minimization problem is said to be *well-posed* if it has a unique solution and every minimizing sequence converges strongly to this solution.

Recall that the Hausdorff distance on the space $\mathfrak{B}(X)$ is defined by

$$h(A, B) = \max \left\{ \sup_{a \in A} \inf_{b \in B} \|a - b\|, \sup_{b \in B} \inf_{a \in A} \|a - b\| \right\}, \quad A, B \in \mathfrak{B}(X).$$

It is well known that $\mathfrak{B}(X)$ endowed with the Hausdorff distance is a complete metric space.

Define $\mathfrak{C}(X) = \{A \in \mathfrak{B}(X) : A \text{ is convex}\}$. For a given set G , let $\mathfrak{C}_G(X)$ stand for the closure of the set $\{A \in \mathfrak{C}(X) : \lambda_{AG} > 0\}$. It is proved in [3] that if X is a uniformly convex Banach space, then the set of all $A \in \mathfrak{C}_G(X)$ such that the minimization problem $\min(A, G)$ is well-posed is a dense G_δ -subset of $\mathfrak{C}_G(X)$. This result has been extended to the framework of strongly convex and/or strictly convex Banach spaces in [7-12]. For further related results see [5, 6, 14-16].

Let $\mathfrak{B}(X) \times \mathfrak{B}(X)$ denote the Cartesian product endowed with the distance

$$d((A, B), (E, F)) = \max\{h(A, E), h(B, F)\} \quad \text{for } A, B, E, F \in \mathfrak{B}(X).$$

Let \mathfrak{D} denote the closure of the set of all $(E, F) \in \mathfrak{B}(X) \times \mathfrak{B}(X)$ such that $\lambda_{EF} > 0$. In this note, we will show that the set of all $(E, F) \in \mathfrak{D}$ such that the minimization problem $\min(E, F)$ is well-posed is a dense G_δ -subset of \mathfrak{D} . In particular, we also show that the set of all $(E, F) \in \mathfrak{D}$ such that the minimization problem $\min(E, F)$ fails to be well-posed is a σ -porous subset of \mathfrak{D} .

2. AUXILIARY RESULTS

For a subset A of X , \bar{A} stands for the closure of A , $\text{diam } A$ for the diameter of A , $\overline{\text{co}}A$ for the closed convex hull of A , and $d(x, A)$ for the distance from x to A . We use $S(x, r)$ to denote the closed ball with center x and radius r in X , in particular, S stands for $S(0, 1)$.

Let $E, F \in \mathfrak{B}(X)$ and $\sigma > 0$. Define

$$L_{E,F}(\sigma) := \{x \in E : d(x, F) \leq \lambda_{EF} + \sigma\} \quad (2.1)$$

It is clear that $L_{E,F}(\sigma_1) \subseteq L_{E,F}(\sigma_2)$ if $\sigma_1 \leq \sigma_2$. The following propositions can be found in [3, 4].

Proposition 2.1. *Let $E, F \in \mathfrak{B}(X)$. Then the problem $\min(E, F)$ is well-posed if and only if*

$$\inf_{\sigma > 0} \text{diam} L_{E,F}(\sigma) = 0 \quad \text{and} \quad \inf_{\sigma > 0} \text{diam} L_{F,E}(\sigma) = 0.$$

Proposition 2.2. *Let $A, B, E, F \in \mathfrak{B}(X)$ and $z \in X$. Then:*

- (i) $|d(z, E) - d(z, F)| \leq h(E, F)$;
- (ii) $\lambda_{EF} \leq d(z, E) + d(z, F)$;
- (iii) $|\lambda_{AB} - \lambda_{EF}| \leq 2d((A, B), (E, F))$.

Define the function Λ on \mathfrak{D} by the formula

$$\Lambda(E, F) = \inf_{\sigma > 0} \text{diam} L_{E,F}(\sigma), \quad \text{for } (E, F) \in \mathfrak{D}.$$

Proposition 2.3. Λ is upper semi-continuous on \mathfrak{D} .

Proof. Let $(E_0, F_0) \in \mathfrak{D}$. Let $\sigma > 0$ and $\delta > 0$. We will show that

$$L_{E,F}(\sigma) \subseteq L_{E_0,F_0}(\sigma + 4\delta) + \delta S \quad (2.2)$$

holds for all $(E, F) \in \mathfrak{D}$ with $d((E, F), (E_0, F_0)) < \delta$. Indeed let $y \in L_{E,F}(\sigma)$. Since $h(E, E_0) < \delta$, there exists $x \in E_0$ such that $\|x - y\| < \delta$. Hence, by Proposition 2.2 and relation (2.1) there is

$$\begin{aligned} d(x, F_0) &\leq d(x, F) + h(F, F_0) \leq d(y, F) + \|x - y\| + h(F, F_0) \leq \\ &\leq d(y, F) + 2\delta \leq \lambda_{EF} + \sigma + 2\delta \leq \lambda_{E_0F_0} + \sigma + 4\delta, \end{aligned}$$

which shows that $x \in L_{E_0,F_0}(\sigma + 4\delta)$. Hence (2.2) holds. Let $\varepsilon > 0$. Choose $\tau > 0$ such that

$$\text{diam } L_{E_0,F_0}(\tau) < \Lambda(E_0, F_0) + \frac{\varepsilon}{2}. \quad (2.3)$$

Taking $\sigma > 0$ and $\delta > 0$ such that $\sigma + 4\delta < \tau$ and $\delta < \varepsilon/4$, by (2.2) and (2.3), we obtain

$$\Lambda(E, F) \leq \text{diam } L_{E,F}(\sigma) \leq \text{diam } L_{E_0,F_0}(\sigma + 4\delta) + 2\delta < \Lambda(E_0, F_0) + \varepsilon$$

for all $(E, F) \in \mathfrak{D}$ with $d((E, F), (E_0, F_0)) < \delta$. This shows that Λ is upper semi-continuous at (E_0, F_0) . \square

The following lemma (see [4]) is essential in our proofs.

Lemma 2.1. Let $\varepsilon > 0$, $\rho > 0$ and let $E \in \mathfrak{C}(X)$. Let $\delta_0 = (\rho/2) \min\{1, \varepsilon\}$. Then for each $u \in X$ with $d(u, E) \geq \rho$ and each $0 < \delta \leq \delta_0$, there is

$$\text{diam } C_{E,u}(\delta) < (\text{diam } E + \delta)\varepsilon,$$

where

$$C_{E,u}(\delta) = \overline{\text{co}}(E \cup \{u\}) \setminus (E + (d(u, E) - \delta)S).$$

3. A GENERIC RESULT FOR MUTUALLY NEAREST POINTS

Let \mathfrak{D}_0 denote the set of all $(E, F) \in \mathfrak{D}$ such that the minimization problem $\min(E, F)$ is well-posed. By virtue of Proposition 2.1,

$$\mathfrak{D}_0 = \bigcap_{k \in \mathbb{N}} \mathfrak{D}_k, \quad (3.1)$$

where

$$\mathfrak{D}_k := \left\{ (E, F) \in \mathfrak{D} : \Lambda(E, F) < \frac{1}{k}, \Lambda(F, E) < \frac{1}{k} \right\}.$$

Theorem 3.1. \mathfrak{D}_0 is a dense G_δ subset of \mathfrak{D} .

Proof. By (3.1), it suffices to verify that each \mathfrak{D}_k ($k \in \mathbb{N}$) is open and dense in \mathcal{D} . The openness of \mathfrak{D}_k is a direct consequence of Proposition 2.3. It remains to show that for every $k \in \mathbb{N}$ the set \mathfrak{D}_k is dense in \mathfrak{D} . To this end, let $(E, F) \in \mathfrak{D}$. Without loss of generality, we may assume that $\lambda_{EF} > 0$. Let $k \in \mathbb{N}$ and $0 < r < \lambda_{EF}/4$. By Lemma 2.1, there exists $0 < \delta < r/2$ such that, for all $u \in X$ with $d(u, E) \geq r/2$ and all $v \in X$ with $d(v, F) \geq r/2$, there holds

$$\text{diam } C_{E,u}(\delta) < \frac{1}{k} \quad \text{and} \quad \text{diam } C_{F,v}(\delta) < \frac{1}{k}.$$

Pick $\hat{x} \in E$ and $\hat{y} \in F$ such that

$$\|\hat{x} - \hat{y}\| < \lambda_{EF} + \delta/2.$$

Note that $\|\hat{x} - \hat{y}\| \geq \lambda_{EF} \geq 4r$. Choose such two points u and v in the interval $[\hat{x}, \hat{y}]$ that $\|\hat{x} - u\| = \|\hat{y} - v\| = r$ and define

$$\tilde{E} = \overline{\text{co}}(E \cup \{u\}), \quad \tilde{F} = \overline{\text{co}}(F \cup \{v\}).$$

Obviously $h(\tilde{E}, E) \leq r$, $h(\tilde{F}, F) \leq r$ and $\lambda_{\tilde{E}\tilde{F}} \geq \lambda_{EF} - 2r > 0$. Hence $(\tilde{E}, \tilde{F}) \in \mathfrak{D}$. To complete the proof it suffices to show that $(\tilde{E}, \tilde{F}) \in \mathfrak{D}_k$ for every $k \in \mathbb{N}$. Note that

$$\|u - \hat{y}\| = \|\hat{x} - \hat{y}\| - \|u - \hat{x}\| \leq \lambda_{EF} + \frac{\delta}{2} - r$$

and

$$d(u, F) \leq \|u - \hat{y}\| \leq \lambda_{EF} + \frac{\delta}{2} - r.$$

From Proposition 2.2, the last inequality and the choice of δ , we conclude

$$d(u, E) \geq \lambda_{EF} - d(u, F) \geq r - \frac{\delta}{2} \geq \frac{3r}{4}. \quad (3.2)$$

On the other hand, since $u \in \tilde{E}$ and $v \in \tilde{F}$, then

$$\lambda_{\tilde{E}\tilde{F}} = \|u - v\| \leq \|\hat{x} - \hat{y}\| - 2r \leq \lambda_{EF} + \frac{\delta}{2} - 2r. \quad (3.3)$$

We claim that

$$L_{\tilde{E}\tilde{F}}(\delta/2) \subseteq C_{E,u}(\delta). \quad (3.4)$$

Indeed, let $y \in L_{\tilde{E}\tilde{F}}(\delta/2) = \overline{\text{co}}(E \cup \{u\})$. By (2.1) and (3.3), there holds

$$d(y, \tilde{F}) \leq \lambda_{\tilde{E}\tilde{F}} + \frac{\delta}{2} \leq \lambda_{EF} + \delta - 2r. \quad (3.5)$$

Then, by Proposition 2.2, relation (3.5), and the inequality $d(u, E) \leq \|u - \hat{x}\| = r$, there follows

$$\begin{aligned} d(y, E) &\geq \lambda_{E\tilde{F}} - d(y, \tilde{F}) \geq \lambda_{E\tilde{F}} - (\lambda_{EF} + \delta - 2r) \geq \\ &\geq \lambda_{EF} - r - (\lambda_{EF} + \delta - 2r) = r - \delta \geq d(u, E) - \delta, \end{aligned}$$

This means that (3.4) holds. From (3.4) and Lemma 2.1 it follows that

$$\Lambda(\tilde{E}, \tilde{F}) \leq \text{diam } C_{E,u}(\delta) < \frac{1}{k}.$$

Similarly, one can show that

$$\Lambda(\tilde{F}, \tilde{E}) \leq \text{diam } C_{F,v}(\delta) < \frac{1}{k}.$$

This means that $(\tilde{E}, \tilde{F}) \in \mathfrak{D}_k$, which completes the proof. \square

4. A POROSITY RESULT

Definition 4.1. A subset Y in a metric space (X, d) is said to be porous in X if there are $0 < t \leq 1$ and $r_0 > 0$ such that for every $x \in X$ and $r \in (0, r_0]$ there is a point $y \in X$ such that $S(y, tr) \subseteq S(x, r) \cap (X \setminus Y)$. A subset Y is said to be σ -porous in X if it is a countable union of sets which are porous in X .

Note that an equivalent definition of a porous set can be obtained by replacing “for every $x \in X$ ” with “for every $x \in Y$ ” (see [1, 3]).

For $(E, F) \in \mathfrak{D}_0$, let $(u_E, u_F) \in E \times F$ denote the unique solution of the minimization problem $\min(E, F)$. Let

$$u_{\alpha, E} = (1 - \alpha)u_E + \alpha u_F, \quad \text{and} \quad E_\alpha = \overline{\text{co}}(E \cup \{u_{\alpha, E}\}), \quad \alpha \in [0, 1].$$

Furthermore, for $r > 0$, set

$$\mathcal{O}(F, r) = \{E \in \mathcal{B}(X) : h(E, F) < r\}.$$

Define

$$\tilde{\mathfrak{D}} = \bigcap_{k \in \mathbb{N}} \bigcup_{(E, F) \in \mathfrak{D}_0} \bigcup_{0 \leq \alpha \leq 1/4} \left(\mathcal{O}(E_\alpha, \gamma_{E_\alpha}(1/k)) \times \mathcal{O}(F_\alpha, \gamma_{F_\alpha}(1/k)) \right),$$

where

$$\gamma_{E_\alpha}(\varepsilon) = \min \{d(u_{\alpha, E}, E), 1\} \varepsilon, \quad \gamma_{F_\alpha}(\varepsilon) = \min \{d(u_{\alpha, F}, F), 1\} \varepsilon.$$

Lemma 4.1. $\tilde{\mathfrak{D}} \subseteq \mathfrak{D}_0$.

Proof. Let $(E, F) \in \tilde{\mathfrak{D}}$. By Proposition 2.2, we only need to show that

$$\Lambda(E, F) = \lim_{\delta \rightarrow 0^+} \text{diam } L_{E, F}(\delta) = 0 \quad \text{and} \quad \Lambda(F, E) = \lim_{\delta \rightarrow 0^+} \text{diam } L_{F, E}(\delta) = 0. \quad (4.1)$$

By the definition of $\tilde{\mathfrak{D}}$, for each $k \in \mathbb{N}$, there exist $(E^k, F^k) \in \mathfrak{D}_0$ and $0 \leq \alpha_k \leq 1/4$ such that

$$h(E, E_{\alpha_k}^k) \leq \gamma_{E_{\alpha_k}^k}(1/k) \quad \text{and} \quad h(F, F_{\alpha_k}^k) \leq \gamma_{F_{\alpha_k}^k}(1/k). \quad (4.2)$$

Without loss of generality, we may assume that $\lambda_{E^k F^k} > 0$ and $\alpha_k > 0$ for each $k \in \mathbb{N}$. For convenience, we write

$$r_k = \lambda_{E^k F^k} \quad \text{and} \quad \delta_k = \gamma_{E_{\alpha_k}^k}(1/k) = \gamma_{F_{\alpha_k}^k}(1/k).$$

Then, it is easy to see that, for each $k \in \mathbb{N}$,

$$\delta_k \leq \alpha_k r_k / k,$$

$$\lambda_{E_{\alpha_k}^k F_{\alpha_k}^k} = (1 - 2\alpha_k)r_k, \quad (4.3)$$

$$\lambda_{E^k F_{\alpha_k}^k} = \lambda_{F^k E_{\alpha_k}^k} = (1 - \alpha_k)r_k, \quad (4.4)$$

$$d(u_{\alpha_k, E^k}, E^k) = d(u_{\alpha_k, F^k}, F^k) = \alpha_k r_k. \quad (4.5)$$

We claim that, for each $\delta > 0$,

$$L_{E_{\alpha_k}^k, F_{\alpha_k}^k}(\delta/2) \subseteq C_{E^k, u_{\alpha_k, E^k}}(\delta) \quad \text{for } k \in \mathbb{N}. \quad (4.6)$$

To see this, let $k \in \mathbb{N}$, $\delta > 0$ and $y \in L_{E_{\alpha_k}^k, F_{\alpha_k}^k}(\delta/2)$. Obviously, $y \in \overline{co}(E^k \cup \{u_{\alpha_k, E^k}\})$. By (2.1) and (4.3), there is

$$d(y, F_{\alpha_k}^k) \leq \lambda_{E_{\alpha_k}^k F_{\alpha_k}^k} + \frac{\delta}{2} = (1 - 2\alpha_k)r_k + \frac{\delta}{2}. \quad (4.7)$$

Consequently, by Proposition 2.2, relation (4.4), (4.7) and (4.5) we obtain

$$\begin{aligned} d(y, E^k) &\geq \lambda_{E^k F_{\alpha_k}^k} - d(y, F_{\alpha_k}^k) \geq (1 - \alpha_k)r_k - (1 - 2\alpha_k)r_k - \delta/2 = \\ &= \alpha_k r_k - \delta/2 = d(u_{\alpha_k, E^k}, E^k) - \delta/2 > d(u_{\alpha_k, E^k}, E^k) - \delta. \end{aligned}$$

Hence $y \in C_{E^k, u_{\alpha_k, E^k}}(\delta)$. Since

$$d((E, F), (E_{\alpha_k}^k, F_{\alpha_k}^k)) < \delta_k,$$

from (2.2) it follows that

$$L_{E, F}(\delta_k) \subseteq L_{E_{\alpha_k}^k, F_{\alpha_k}^k}(5\delta_k) + \delta_k S.$$

By the last inclusion and (4.6) we obtain

$$\begin{aligned} \Lambda(E, F) &\leq \text{diam } L_{E, F}(\delta_k) \leq \text{diam } L_{E_{\alpha_k}^k, F_{\alpha_k}^k}(5\delta_k) + 2\delta_k \leq \\ &\leq \text{diam } C_{E^k, u_{\alpha_k}}(10\delta_k) + 2\delta_k \end{aligned} \quad (4.8)$$

for each $k \in \mathbb{N}$. Recall that

$$d(u_{\alpha_k, E^k}, E^k) = \alpha_k r_k \quad \text{and} \quad \delta_k \leq \frac{\alpha_k r_k}{k} \quad \text{for each } k > 1.$$

Then using Lemma 2.1 we conclude that

$$\text{diam } C_{E^k, u_{\alpha_k}}(10\delta_k) \leq \frac{2}{k}(\text{diam } E^k + 10\alpha_k r_k),$$

and hence, by (4.8),

$$\Lambda(E, F) \leq \frac{2}{k}(\text{diam } E^k + 10\alpha_k r_k) + 2\delta_k \leq \frac{2}{k}(\text{diam } E^k + 11\alpha_k r_k). \quad (4.9)$$

Note that

$$h(E, E^k) \leq h(E, E_{\alpha_k}^k) \leq \gamma_{E_{\alpha_k}^k}(1/k) \leq 1.$$

Analogously $h(F, F^k) \leq 1$. Thus $h(E^k, F^k) \leq h(E, F) + 2$. It follows that sequences $\{\text{diam } E^k\}$ and $\{r_k\}$ are bounded. Hence (4.9) implies that $\Lambda(E, F) = 0$. Similarly, we can verify that $\Lambda(F, E) = 0$. Hence (4.1) holds and the proof of Lemma 4.1 is complete. \square

Theorem 4.1. *The set $\mathfrak{D} \setminus \mathfrak{D}_0$ is σ -porous in \mathfrak{D} .*

Proof. For $k, l \in \mathbb{N}$, define

$$\tilde{\mathfrak{D}}_k = \mathfrak{D} \setminus \bigcup_{(E, F) \in \mathfrak{D}_0} \bigcup_{0 \leq \alpha \leq 1/4} \left(\mathcal{O}(E_\alpha, \gamma_{E_\alpha}(1/k)) \times \mathcal{O}(F_\alpha, \gamma_{F_\alpha}(1/k)) \right)$$

and

$$\tilde{\mathcal{D}}_k^l = \left\{ (E, F) \in \tilde{\mathfrak{D}}_k : \frac{1}{l} < \lambda_{EF} < l \right\}.$$

Observe that

$$\mathfrak{D} \setminus \mathfrak{D}_0 \subseteq \mathfrak{D} \setminus \tilde{\mathfrak{D}} = \bigcup_{k \in \mathbb{N}} \bigcup_{l \in \mathbb{N}} \tilde{\mathcal{D}}_k^l.$$

It suffices to verify that, $\tilde{\mathfrak{D}}_k^l$ is porous in \mathfrak{D} for each $k, l \in \mathbb{N}$. To this end, let $k, l \in \mathbb{N}$ be arbitrary. Define $r_0 = 1/(2l)$ and $\alpha = 1/(4k)$. Let $(E, F) \in \tilde{\mathfrak{D}}_k^l$ and $0 < r \leq r_0$. Then, by Theorem 2.1, there exists $(\bar{E}, \bar{F}) \in \mathfrak{D}_0$ such that

$$h(E, \bar{E}) < \frac{r}{4}, \quad h(F, \bar{F}) < \frac{r}{4}$$

and

$$\frac{1}{l} < \lambda_{\bar{E}\bar{F}} < l.$$

Set $\bar{u}_{1/2} = (u_{\bar{E}} + u_{\bar{F}})/2$. Then

$$\begin{aligned} h(\bar{E}_{1/2}, E) &\geq h(\bar{E}_{1/2}, \bar{E}) - h(\bar{E}, E) \geq \\ &\geq \sup_{y \in \bar{E}_{1/2}} d(y, \bar{E}) - r/4 \geq \\ &\geq d(\bar{u}_{1/2}, \bar{E}) - r/4 = \\ &= (1/2)\lambda_{\bar{E}\bar{F}} - r/4 \geq 3r/4. \end{aligned}$$

Similarly, one can prove that

$$h(\bar{F}_{1/2}, F) \geq 3r/4.$$

From the previous two inequalities it follows that there exist $0 < t_1, t_2 \leq 1/2$ such that $h(\bar{E}_{t_1}, E) = 3r/4$ and $h(\bar{F}_{t_2}, F) = 3r/4$, where $\bar{E}_{t_1} = \bar{c}\bar{o}(\bar{E} \cup u_{t_1, \bar{E}})$ and $\bar{F}_{t_2} = \bar{c}\bar{o}(\bar{F} \cup u_{t_2, \bar{F}})$. Observe that

$$\mathcal{O}(\bar{E}_{t_1}, \alpha r) \subseteq \mathcal{O}(E, r) \quad \text{and} \quad \mathcal{O}(\bar{F}_{t_2}, \alpha r) \subseteq \mathcal{O}(F, r). \quad (4.10)$$

Indeed, for each $A \in \mathcal{O}(\bar{E}_{t_1}, \alpha r)$

$$h(A, E) \leq h(A, \bar{E}_{t_1}) + h(\bar{E}_{t_1}, E) \leq \alpha r + 3r/4 \leq r.$$

Hence the first inequality of (4.10) is proved. The second one can be proved analogously.

Now we claim that

$$\alpha r \leq \gamma_{\bar{E}_{t_1}}(1/k) \quad \text{and} \quad \alpha r \leq \gamma_{\bar{F}_{t_2}}(1/k). \quad (4.11)$$

Indeed, note that

$$h(\bar{E}_{t_1}, \bar{E}) \geq h(\bar{E}_{t_1}, E) - h(E, \bar{E}) \geq r/2.$$

Therefore,

$$\alpha r \leq 2\alpha h(\bar{E}_{t_1}, \bar{E}) \leq h(\bar{E}_{t_1}, \bar{E})/k = d(u_{t_1, \bar{E}}, \bar{E})/k.$$

Since obviously $\alpha r \leq 1/k$, the first inequality of (4.11) is proved. The second one can be proved analogously.

From (4.11) it follows that

$$\mathcal{O}(\bar{E}_{t_1}, \alpha r) \times \mathcal{O}(\bar{F}_{t_2}, \alpha r) \subseteq \mathcal{O}(\bar{E}_{t_1}, \gamma_{\bar{E}_{t_1}}(1/k)) \times \mathcal{O}(\bar{F}_{t_2}, \gamma_{\bar{F}_{t_2}}(1/k)).$$

This implies that

$$\mathcal{O}(\bar{E}_{t_1}, \alpha r) \times \mathcal{O}(\bar{F}_{t_2}, \alpha r) \subseteq \mathfrak{D} \setminus \tilde{\mathfrak{D}}_k^l.$$

From this last inclusion and relation (4.10) it immediately follows that the set $\tilde{\mathfrak{D}}_k^l$ is porous in \mathfrak{D} . \square

Acknowledgements

Supported in part by the National Natural Science Foundations of China (Grant No. 10271025).

REFERENCES

- [1] F.S. De Blasi, J. Myjak, *Sur la porosité des ensemble des contractions sans point fixe*, C.R. Acad. Sci. Paris **308** (1989), 51–54.

- [2] F.S. De Blasi, J. Myjak, P.L. Papini, *Starshaped sets and best approximation*, Arch. Math. **56** (1991), 41–48.
- [3] F.S. De Blasi, J. Myjak, P.L. Papini, *Porous sets in best approximation theory*, J. London Math. Soc. **44** (1991), 135–142.
- [4] F.S. De Blasi, J. Myjak, P.L. Papini, *On mutually nearest and mutually furthest points of sets in Banach spaces*, J. Approx. Theory **70** (1992), 142–155.
- [5] J.M. Borwein, S. Fitzpatrick, *Existence of nearest points in Banach spaces*, Can. J. Math. **41** (1989), 702–720.
- [6] S.V. Konjagin, *On approximation properties of closed sets in Banach spaces and characterization of strongly convex spaces*, Soviet Math. Dokl. **21** (1980), 418–422.
- [7] C. Li, *On mutually nearest and mutually furthest points in reflexive Banach spaces*, J. Approx. Theory, **103** (2000), 1–17.
- [8] C. Li, *On well-posedness of best simultaneous approximation problems in Banach spaces*, Science in China (Ser A), **44** (2001), 1558–1570.
- [9] C. Li, J. Myjak, *On mutually nearest points of unbounded sets in Banach spaces*, J. Nonlinear Convex Anal. **8** (2007), 165–177.
- [10] C. Li, R.X. Ni, *On well-posed mutually nearest and mutually furthest point problems in Banach spaces*, Acta Math. Sinica, New Ser. **20** (2004), 147–156.
- [11] C. Li, H.K. Xu, *On almost well-posed mutually nearest and mutually furthest points problems*, Numer. Funct. Anal. Optim. **23** (2002), 323–331.
- [12] C. Li, H.K. Xu, *Porosity of mutually nearest and mutually furthest points in Banach spaces*, J Approx. Theory **125** (2003), 10–25.
- [13] S. Reich, A.J. Zaslavski, *Well-posedness and porosity in best approximation problems*, Topol. Methods Nonlinear Anal. **18** (2001), 395–408.
- [14] S. Reich, A.J. Zaslavski, *A porosity results in best approximation theory*, J. Nonlinear Convex Anal. **4** (2003), 165–173.
- [15] S. Reich, A.J. Zaslavski, *Well-posedness of generalized best approximation problems*, Nonlinear Funct. Anal. Appl. **7** (2002), 115–128.
- [16] S. Reich, A.J. Zaslavski, *Porous sets and generalized best approximation problems*, Nonlinear Anal. Forum **9** (2004), 135–152.

Chong Li
cli@zju.edu.cn

Zhejiang University
Department of Mathematics
Hangzhou 310027, P. R. China

Józef Myjak
myjak@univag.it

AGH University of Science and Technology
Faculty of Applied Mathematics
al. Mickiewicza 30, 30-059 Cracow

Received: April 2, 2007.

Revised: June 9, 2007.

Accepted: June 12, 2007.