

**RIFT-RELATED ENVIRONMENTAL CHANGE  
IN THE NORTHERN TETHYS  
BASED ON ND AND SR ISOTOPES IN THE MIDDLE  
AND UPPER JURASSIC CARBONATES  
OF THE PIENINY KLIPPEN BELT**

**Magdalena SIDORCZUK<sup>1</sup>,  
Marek LEWANDOWSKI<sup>2, 3</sup> & Zdzisław BEŁKA<sup>4</sup>**

<sup>1</sup>*University of Warsaw, Faculty of Geology;  
ul. Żwirki i Wigury 93, 02-089 Warsaw; e-mail: sima@uw.edu.pl*

<sup>2</sup>*Polish Academy of Sciences, Institute of Geophysics;  
ul. ks. Janusza 64, 01-452 Warsaw; e-mail: lemar@igf.edu.pl*

<sup>3</sup>*Polish Academy of Sciences, Institute of Geological Sciences;  
ul. Twarda 51/55, 00-818 Warsaw*

<sup>4</sup>*Adam Mickiewicz University, Isotope Laboratory;  
ul. Maków Polnych 16, 61-606 Poznań; e-mail: zbelka@amu.edu.pl*

The Pieniny Klippen Belt was formed during the Late Cretaceous-Cenozoic tectonic movements due to the closure of the Pieniny Klippen Basin (PKB), a former part of northern sector of the Tethys Ocean. Data collected from the Middle and Upper Jurassic rocks from the Veliky Kamenets section, Western Ukraine (see Lewandowski *et al.*, 2005, 3Palaeo, 216, 53–72) to the Western Slovakia (Lewandowski *et al.*, in prep.) speak in favor of significant paleogeographic rearrangement of the crustal units, which formed the substratum of PKB. A direct implication of observed shallowing of paleolatitude, amounting 10° between the Early Callovian and the Middle Oxfordian (*ca.* 10 Ma), was a rapid and a long-range drift of PKB basement, rifted off the northern European Craton toward the South. Sedimentary record of this event is lacking, a hiatus embracing the Middle Callovian/Early Oxfordian time span. Our goal was to test the rifting scenario using geochemical signatures of rocks bracketing the hiatus.

In our study, we have collected carbonate rocks underlying and overlying the hiatus, following the same outcrops that were sampled for paleomagnetic studies. Some samples from the manganese crusts were also collected. We performed geochemical studies including REE analysis using ICP-OES and ICP-MS techniques, which demonstrated higher concentrations of REE and U, Th, K, Nb, Zr and Hf in samples collected from limestones underlying the hiatus, as well as a significantly different concentrations of La/Yb-Sc/Ni and Zr/Th between the Callovian and the Oxfordian limestones. We tentatively interpreted this

record as an effect of change in a source of elemental alimentation for the PKB that took place between Batonian and Oxfordian time, the latter including more mafic components (Sidorczuk *et al.* 2009, IAS Abstract book).

In a current presentation, we demonstrate change of the oceanic environment of the Pieniny Klippen Basin using Nd and Sr isotopic composition of the carbonate and silicate fractions. Analyses were performed using Finnigan MAT 261 MC-MS at the Isotope Lab., UAM, Poznań (Poland).

Neodymium isotopic composition of the carbonate fraction analyzed from the strata bracketing the hiatus is regarded as a direct record of the contemporaneous composition of the seawater. It shows no evidence for localized input of old continental Nd from the adjacent land areas. Our analyses revealed that both the Early Callovian and the Middle/Late Oxfordian seawaters of eastern part of the Pieniny Klippen Basin were isotopically homogeneous in each case. The main result is, however, that a significant exchange of the seawater happened between Early Callovian and Middle Oxfordian time. Samples collected from strata below the hiatus yielded  $\epsilon_{\text{Nd}}$  values ranging from  $-6.6$  to  $-7.0$ . These isotopic signatures are identical to those known from the Alpine part of the Tethys. Above the hiatus, however, the  $\epsilon_{\text{Nd}}$  values are constantly higher, between  $-5.3$  and  $-5.8$ , marking opposite trend in  $\epsilon_{\text{Nd}}$  evolution than postulated for the Western Tethys in the same time span (*cf.* Stille *et al.* 1996, EPSL, 144, 9-19). Our data, therefore, record enter of more radiogenic oceanic waters into the basin. Because the seawater in the western segment (Alpine-Penninic) of the Tethys was predominantly less radiogenic during the Middle Oxfordian, we speculate that oceanic waters could be introduced into the Pieniny Basin from the Pacific Ocean. This event could be causally linked to a rapid rifting process and dramatic widening of the Pieniny Basin, in line with the scenario of the substantial paleogeographic change indicated by paleomagnetic results.

*Acknowledgment Grant no. N30704332/195 MNiSW.*