

**STRONG CONVERGENCE THEOREM
OF A HYBRID PROJECTION ALGORITHM
FOR A FAMILY OF QUASI- ϕ -ASYMPTOTICALLY
NONEXPANSIVE MAPPINGS**

J.F. Tang, S.S. Chang, M. Liu, J.A. Liu

Abstract. The main purpose of this paper is by using a new hybrid projection iterative algorithm to prove some strong convergence theorems for a family of quasi- ϕ -asymptotically nonexpansive mappings. The results presented in the paper improve and extend the corresponding results announced by some authors.

Keywords: quasi- ϕ -asymptotically nonexpansive mapping, asymptotically regular mapping, hybrid projection iterative algorithm, strong convergence theorem.

Mathematics Subject Classification: 47H09, 47H10.

1. INTRODUCTION

Throughout this paper, we assume that E is a real Banach space, E^* is the dual space of E , C is a nonempty closed convex subset of E , and $\langle \cdot, \cdot \rangle$ is the pairing between E and E^* . Recall that a mapping $T : C \rightarrow C$ is said to be asymptotically nonexpansive [1] if there exists a sequence $\{k_n\} \subset [1, \infty)$ with $k_n \rightarrow 1$ such that

$$\|T^n x - T^n y\| \leq k_n \|x - y\| \quad \forall x, y \in C \quad \text{and} \quad \forall n \geq 1. \quad (1.1)$$

In recent years, nonexpansive mappings and asymptotically nonexpansive mappings have been studied extensively by many authors. In 2003, Nakajo and Takahashi [2] proposed the following modification of the Mann iteration method for a nonexpansive mapping T in a Hilbert space H :

$$\begin{cases} x_0 \in C, \\ y_n = \alpha_n x_n + (1 - \alpha_n) T x_n, \\ C_n = \{z \in C : \|y_n - z\| \leq \|x_n - z\|\}, \\ Q_n = \{z \in C : \langle x_n - z, x_0 - x_n \rangle \geq 0\}, \\ x_{n+1} = P_{C_n \cap Q_n}(x_0). \end{cases} \quad (1.2)$$

where C is a closed convex subset of H , and P_K is the metric projection from H onto a closed convex subset K of H . They proved that if the sequence $\{\alpha_n\}$ is bounded above from one, then the sequence $\{x_n\}$ generated by (1.2) converges strongly to $P_{F(T)}(x_0)$.

In 2006, Kim and Xu [4] proposed the following modification of the Mann iteration method for a asymptotically nonexpansive mapping T in a Hilbert space H :

$$\begin{cases} x_0 \in C, \\ y_n = \alpha_n x_n + (1 - \alpha_n) T^n x_n, \\ C_n = \{z \in C : \|y_n - z\|^2 \leq \|x_n - z\|^2 + \theta_n\}, \\ Q_n = \{z \in C : \langle x_n - z, x_0 - x_n \rangle \geq 0\}, \\ x_{n+1} = P_{C_n \cap Q_n}(x_0), \end{cases} \quad (1.3)$$

where C is a bounded closed convex subset and $\theta_n = (1 - \alpha_n)(k_n^2 - 1)(\text{diam}C)^2 \rightarrow 0$ ($n \rightarrow \infty$). They proved that if the sequence $\{\alpha_n\}$ is bounded above from one, then the sequence $\{x_n\}$ generated by (1.3) converges strongly to $P_{F(T)}(x_0)$.

In 2005, Matsushita and Takahashi [3] proposed the following hybrid iteration method with generalized projection for a relatively nonexpansive mapping T in a Banach space E :

$$\begin{cases} x_0 \in C, \\ y_n = J^{-1}(\alpha_n Jx_n + (1 - \alpha_n)JT x_n), \\ C_n = \{z \in C : \phi(z, y_n) \leq \phi(z, x_n)\}, \\ Q_n = \{z \in C : \langle x_n - z, Jx_0 - Jx_n \rangle \geq 0\}, \\ x_{n+1} = \Pi_{C_n \cap Q_n}(x_0). \end{cases} \quad (1.4)$$

Under suitable conditions they proved that the sequence $\{x_n\}$ generated by (1.4) converges strongly to $\Pi_{F(T)}(x_0)$.

In 2009, Zhou and Gao [5] proposed the following modified hybrid iteration method with generalized projection for a family of closed and quasi- ϕ -asymptotically nonexpansive mappings which are asymptotically regular in a Banach space E :

$$\begin{cases} x_0 \in C, \\ y_{n,i} = J^{-1}(\alpha_n Jx_n + (1 - \alpha_n)JT_i^n x_n), \\ C_{n,i} = \{z \in C : \phi(z, y_{n,i}) \leq \phi(z, x_n) + \xi_{n,i}\}, \\ C_n = \bigcap_{i \in I} C_{n,i}, \\ Q_0 = C, \\ Q_n = \{z \in Q_{n-1} : \langle x_n - z, Jx_0 - Jx_n \rangle \geq 0\}, \\ x_{n+1} = \Pi_{C_n \cap Q_n}(x_0). \end{cases} \quad (1.5)$$

Under suitable conditions they proved that the sequence $\{x_n\}$ generated by (1.5) converges strongly to $\Pi_{\bigcap_{i \in I} F(T_i)}(x_0)$.

Motivated and inspired by the research going on in this direction, the purpose of this paper is to introduce a hybrid projection iterative algorithm and prove strong

some convergence theorems for a family of quasi- ϕ -asymptotically nonexpansive mappings in the setting of Banach spaces. The results presented in the paper improve and extend the corresponding results in [2–5].

2. PRELIMINARIES

Let E be a Banach space with a dual E^* and C be a nonempty closed convex subsets of E . We denote by $J : E \rightarrow 2^{E^*}$ the normalized duality mapping defined by

$$Jx = \{f \in E^* : \langle x, f \rangle = \|x\|^2 = \|f\|^2\}, \quad x \in E.$$

It is well known that if E is uniformly convex and uniformly smooth, then J and J^{-1} both are uniformly continuous on bounded subsets of E and E^* , respectively.

In the sequel, we always denote by $\phi : E \times E \rightarrow R^+$ the Lyapunov functional defined by

$$\phi(x, y) = \|x\|^2 - 2\langle x, Jy \rangle + \|y\|^2, \quad \forall x, y \in E. \quad (2.1)$$

From the definition of ϕ , it is obvious that

$$(\|x\| - \|y\|)^2 \leq \phi(x, y) \leq (\|x\| + \|y\|)^2, \quad \forall x, y \in E. \quad (2.2)$$

The *generalized projection* $\Pi_C : E \rightarrow C$ is defined by

$$\Pi_C(x) = \inf_{y \in C} \phi(y, x), \quad \forall x \in E. \quad (2.3)$$

Lemma 2.1 ([6]). *Let E be a smooth, strict convex and reflexive Banach space and C be a nonempty closed convex subset of E . Then, the following conclusions hold:*

- (i) $\phi(x, \Pi_C y) + \phi(\Pi_C y, y) \leq \phi(x, y), \quad \forall x \in C, y \in E.$
- (ii) *Let $x \in E$ and $z \in C$, then*

$$z = \Pi_C x \iff \langle z - y, Jx - Jz \rangle \geq 0, \quad \forall y \in C. \quad (2.4)$$

Let C be a closed convex subset of E , and T a mapping from C into itself. T is said to be *ϕ -asymptotically nonexpansive*, if there exists a sequence $\{k_n\} \subset [1, \infty)$ with $k_n \rightarrow 1$ such that $\phi(T^n x, T^n y) \leq k_n \phi(x, y)$ for all $n \geq 1$ and $x, y \in C$. T is said to be *quasi- ϕ -asymptotically nonexpansive*, if $F(T) \neq \emptyset$ and there exists a sequence $\{k_n\} \subset [1, \infty)$ with $k_n \rightarrow 1$ such that $\phi(p, T^n x) \leq k_n \phi(p, x)$ for all $n \geq 1$, $x \in C$ and $p \in F(T)$. T is said to be *closed*, if for any $\{x_n\}$ with $x_n \rightarrow x$ and $Tx_n \rightarrow y$, then we have $Tx = y$.

T is said to be *asymptotically regular* on C if, for any bounded subset D of C , the following equality holds:

$$\limsup_{n \rightarrow \infty} \{\|T^{n+1}x - T^n x\| : x \in D\} = 0.$$

The following lemmas will play an important role in the proof of the main results in this paper.

Lemma 2.2 ([6]). *Let E be a uniformly convex and smooth Banach space and $\{x_n\}, \{y_n\}$ be sequences of E . If $\phi(x_n, y_n) \rightarrow 0$ (as $n \rightarrow \infty$) and either $\{x_n\}$ or $\{y_n\}$ is bounded, then $x_n - y_n \rightarrow 0$ (as $n \rightarrow \infty$).*

Lemma 2.3 ([5]). *Let E be a uniformly convex and smooth Banach space, C be a closed convex subset of E , and T be a closed and quasi- ϕ -asymptotically nonexpansive mapping from C into itself. Then $F(T)$ is a closed convex subset of C .*

Lemma 2.4 ([7]). *Let E be a uniformly convex Banach space, $r > 0$ be a positive number, and $B_r(0)$ be a closed ball of E . For any given points $\{x_1, x_2, \dots, x_n, \dots\} \subset B_r(0)$ and for any given positive numbers $\{\lambda_1, \lambda_2, \dots, \lambda_n, \dots\}$ with $\sum_{n=1}^{\infty} \lambda_n = 1$, there exists a continuous, strictly increasing and convex function $g : [0, 2r) \rightarrow [0, \infty)$ with $g(0) = 0$ such that for any $i, j \in \{1, 2, \dots\}, i < j$,*

$$\left\| \sum_{n=1}^{\infty} \lambda_n x_n \right\|^2 \leq \sum_{n=1}^{\infty} \lambda_n \|x_n\|^2 - \lambda_i \lambda_j g(\|x_i - x_j\|). \quad (2.5)$$

3. MAIN RESULTS

Theorem 3.1. *Let C be a nonempty bounded closed convex subset of a uniformly convex and uniformly smooth Banach space E . For each $i = 1, 2, \dots$, let $T_i : C \rightarrow C$ be a closed and quasi- ϕ -asymptotically nonexpansive mapping with a sequence $\{k_{n,i}\} \subset [1, \infty)$ such that $k_n := \sup_{i \geq 1} k_{n,i} \rightarrow 1$ ($n \rightarrow \infty$) and $F = \bigcap_{i=1}^{\infty} F(T_i) \neq \emptyset$. Suppose further that for each $i = 1, 2, \dots$, T_i is asymptotically regular on C . Let $\{x_n\}$ be the sequence in C defined by:*

$$\begin{cases} x_0 \in C, & C_0 = C, \\ y_n = J^{-1}(\alpha_{n0} Jx_n + \sum_{i=1}^{\infty} \alpha_{ni} J T_i^n x_n), \\ C_{n+1} = \{z \in C_n : \phi(z, y_n) \leq \phi(z, x_n) + \xi_n\}, \\ x_{n+1} = \Pi_{C_{n+1}}(x_0), \end{cases} \quad (3.1)$$

where $J : E \rightarrow E^*$ is the normalized duality mapping, $M = \sup_{z \in F, n \geq 1} \phi(z, x_n)$, $\xi_n = \sum_{i=1}^{\infty} \alpha_{ni} (k_n - 1) M$, and $\{\alpha_{ni}\}$ is the sequence in $[0, 1]$ satisfying the following conditions:

- (a) $\sum_{i=0}^{\infty} \alpha_{ni} = 1, \forall n \geq 0$;
- (b) $\liminf_{n \rightarrow \infty} \alpha_{n0} \alpha_{ni} > 0, i = 1, 2, \dots$

Then $\{x_n\}$ converges strongly to $\Pi_F x_0$, where Π_F is the generalized projection of E onto F .

Proof. (I) Because $\phi(z, y_n) \leq \phi(z, x_n) + \xi_n$ is equivalent to $2\langle z, Jx_n - Jy_n \rangle \leq \|x_n\|^2 - \|y_n\|^2 + \xi_n$. This implies that C_n is a closed and convex subset of C for all $n \geq 0$.

(II) Next, we prove that $F := \bigcap_{i=1}^{\infty} F(T_i) \subset C_n, \forall n \geq 0$.

Indeed, it is obvious that $F \subset C_0$. Suppose that $F \subset C_n$ for some $n \in \mathbb{N}$. Noting that $\|\cdot\|^2$ is convex and using (2.1), for any $z \in F \subset C_n$ and for any $\forall m, j \in \{0, 1, 2, \dots\}, m < j$, we have that

$$\begin{aligned}
 \phi(z, y_n) &= \phi(z, J^{-1}(\alpha_{n0}Jx_n + \sum_{i=1}^{\infty} \alpha_{ni}JT_i^n x_n)) = \\
 &= \|z\|^2 - 2\langle z, \sum_{i=0}^{\infty} \alpha_{ni}JT_i^n x_n \rangle + \|\sum_{i=0}^{\infty} \alpha_{ni}JT_i^n x_n\|^2 \text{ (where } T_0 = I) \leq \\
 &\leq \|z\|^2 - \sum_{i=0}^{\infty} \alpha_{ni}2\langle z, JT_i^n x_n \rangle + \sum_{i=0}^{\infty} \alpha_{ni}\|T_i^n x_n\|^2 - \\
 &\quad - \alpha_{nm}\alpha_{nj}g(\|JT_m^n x_n - JT_j^n x_n\|) = \\
 &= \sum_{i=0}^{\infty} \alpha_{ni}(\|z\|^2 - 2\langle z, JT_i^n x_n \rangle + \|T_i^n x_n\|^2) - \\
 &\quad - \alpha_{nm}\alpha_{nj}g(\|JT_m^n x_n - JT_j^n x_n\|) = \\
 &= \alpha_{n0}\phi(z, x_n) + \sum_{i=1}^{\infty} \alpha_{ni}\phi(z, T_i^n x_n) - \alpha_{nm}\alpha_{nj}g(\|JT_m^n x_n - JT_j^n x_n\|) \leq \\
 &\leq \alpha_{n0}\phi(z, x_n) + \sum_{i=1}^{\infty} \alpha_{ni}((k_{n,i} - 1) + 1)\phi(z, x_n) - \\
 &\quad - \alpha_{nm}\alpha_{nj}g(\|JT_m^n x_n - JT_j^n x_n\|) \leq \\
 &\leq \phi(z, x_n) + \xi_n - \alpha_{nm}\alpha_{nj}g(\|JT_m^n x_n - JT_j^n x_n\|) \leq \phi(z, x_n) + \xi_n.
 \end{aligned} \tag{3.2}$$

This implies that $z \in C_n$. Thereby, $F \subset C_n, \forall n \geq 0$.

(III) Now, we prove that $\{x_n\}$ is a Cauchy sequence.

Indeed, since $x_{n+1} = \Pi_{C_{n+1}}x_0$ and $x_n = \Pi_{C_n}x_0, x_{n+1} \in C_{n+1} \subset C_n$, from the definition of Π_{C_n} we have

$$\phi(x_n, x_0) \leq \phi(x_{n+1}, x_0), \quad \forall n \geq 0. \tag{3.3}$$

Therefore $\{\phi(x_n, x_0)\}$ is nondecreasing. By the assumption that C is bounded, hence from (2.2) we know that $\{\phi(x_n, x_0)\}$ is bounded. This together with (3.3) ensures that the limit $\{\phi(x_n, x_0)\}$ exists. Write

$$\lim_{n \rightarrow \infty} \phi(x_n, x_0) = d. \tag{3.4}$$

From Lemma 2.1, we have, for any positive integer $m \geq n$, that

$$\begin{aligned}
 \phi(x_m, x_n) &= \phi(x_m, \Pi_{C_n}x_0) \leq \phi(x_m, x_0) - \phi(\Pi_{C_n}x_0, x_0) = \\
 &= \phi(x_m, x_0) - \phi(x_n, x_0)
 \end{aligned} \tag{3.5}$$

This implies that

$$\lim_{n, m \rightarrow \infty} \phi(x_m, x_n) = 0. \tag{3.6}$$

By Lemma 2.2, we know that $x_m - x_n \rightarrow 0$ ($n, m \rightarrow \infty$), hence, $\{x_n\}$ is a Cauchy sequence. Without loss of generality, we can assume that $x_n \rightarrow p \in C$ ($n \rightarrow \infty$).

(IV) Now, we prove $\|x_n - T_i^n x_n\| \rightarrow 0$ for each $i = 1, 2, \dots$

In fact, taking $m = n + 1$ in (3.5) we have that

$$\phi(x_{n+1}, x_n) \leq \phi(x_{n+1}, x_0) - \phi(x_n, x_0) \rightarrow 0 \quad (n \rightarrow \infty), \quad (3.7)$$

and hence $x_{n+1} - x_n \rightarrow 0$ as $n \rightarrow \infty$ by Lemma 2.2. Since $x_{n+1} = \Pi_{C_{n+1}} x_0 \in C_{n+1}$, and by the assumption that $\xi_n \rightarrow 0$ (as $n \rightarrow \infty$), hence from the definition of C_{n+1} , we have

$$\phi(x_{n+1}, y_n) \leq \phi(x_{n+1}, x_n) + \xi_n \rightarrow 0 \quad (n \rightarrow \infty), \quad (3.8)$$

and so $x_{n+1} - y_n \rightarrow 0$ ($n \rightarrow \infty$) by Lemma 2.2 Thus we have

$$\|y_n - x_n\| \leq \|y_n - x_{n+1}\| + \|x_{n+1} - x_n\| \rightarrow 0 \quad (n \rightarrow \infty). \quad (3.9)$$

Since J is uniformly continuous on any bounded sets of E , we conclude that

$$\lim_{n \rightarrow \infty} \|Jx_n - Jy_n\| = 0. \quad (3.10)$$

On the other hand, taking $m = 0$ and $j = 1, 2, \dots$ in (3.2), for any $z \in F$, we have

$$\phi(z, y_n) \leq \phi(z, x_n) + \xi_n - \alpha_{n0} \alpha_{nj} g(\|Jx_n - JT_j^n x_n\|),$$

i.e.,

$$\alpha_{n0} \alpha_{nj} g(\|Jx_n - JT_j^n x_n\|) \leq \phi(z, x_n) - \phi(z, y_n) + \xi_n. \quad (3.11)$$

Since

$$\begin{aligned} \phi(z, x_n) - \phi(z, y_n) + \xi_n &= \|x_n\|^2 - \|y_n\|^2 - 2\langle z, Jx_n - Jy_n \rangle + \xi_n \leq \\ &\leq \|x_n\|^2 - \|y_n\|^2 + 2\|z\| \|Jx_n - Jy_n\| + \xi_n \leq \\ &\leq \|x_n - y_n\| (\|x_n + y_n\|) + 2\|z\| \|Jx_n - Jy_n\| + \xi_n \end{aligned} \quad (3.12)$$

from (3.9) and (3.10), it follows that $\phi(z, x_n) - \phi(z, y_n) + \xi_n \rightarrow 0$ ($n \rightarrow \infty$). Hence, from (3.11) and condition (b) in Theorem 3.1, we have that

$$g(\|Jx_n - JT_j^n x_n\|) \rightarrow 0 \quad (n \rightarrow \infty), \quad \forall j = 1, 2, \dots \quad (3.13)$$

Since g is continuous and strictly increasing with $g(0) = 0$, it follows from (3.13) that

$$\|Jx_n - JT_j^n x_n\| \rightarrow 0 \quad (as \ n \rightarrow \infty) \quad \text{and for each } j = 1, 2, \dots$$

Again by the assumption that E is uniformly convex and so E^* is uniformly smooth, hence J^{-1} is uniformly continuous on any bounded subset of E^* . Therefore we have

$$\|x_n - T_j^n x_n\| \rightarrow 0 \quad (n \rightarrow \infty), \quad \text{for each } j = 1, 2, \dots \quad (3.14)$$

(V) Now, we prove $p \in F$.

From $x_n \rightarrow p$ ($n \rightarrow \infty$) and (3.14), we have

$$T_j^n x_n \rightarrow p \quad (n \rightarrow \infty) \quad \text{for each } j = 1, 2, \dots \tag{3.15}$$

Noting that

$$\|T_i^{n+1} x_n - p\| \leq \|T_i^{n+1} x_n - T_i^n x_n\| + \|T_i^n x_n - p\|, \tag{3.16}$$

using (3.15) and the asymptotic regularity of T_i , from (3.16) we have

$$T_i^{n+1} x_n \rightarrow p \quad (n \rightarrow \infty), \quad \text{i.e., } T_i T_i^n x_n \rightarrow p \quad (n \rightarrow \infty). \tag{3.17}$$

By virtue of the closeness of T_i , it follows from (3.15) and (3.17) that p is a fixed point of T_i , $\forall i \geq 1$, i.e., $p \in F$.

(VI) Now, we prove $x_n \rightarrow p = \Pi_F x_0$ ($n \rightarrow \infty$).

Let $w = \Pi_F x_0$. From $w \in F \subset C_{n+1}$ and $x_{n+1} = \Pi_{C_{n+1}} x_0$, we have $\phi(x_{n+1}, x_0) \leq \phi(w, x_0)$, $\forall n \geq 0$. This implies that

$$\phi(p, x_0) = \lim_{n \rightarrow \infty} \phi(x_n, x_0) \leq \phi(w, x_0). \tag{3.18}$$

By the definition of $\Pi_F x_0$ and (3.18), we have $p = w$. Therefore, $x_n \rightarrow p = \Pi_F x_0$ ($n \rightarrow \infty$).

This completes the proof of theorem 3.1. □

Remark 3.2. *The asymptotic regularity assumption on T_i in Theorem 3.1 can be weakened to the assumption that $T_i^{n+1} x_n - T_i^n x_n \rightarrow 0$ as $n \rightarrow \infty$. The assumption that $T_i^{n+1} x_n - T_i^n x_n \rightarrow 0$ as $n \rightarrow \infty$ can be replaced by the uniform Lipschitz continuous of T_i .*

Therefore, we have the following convergence result.

Corollary 3.3. *Let C be a nonempty bounded closed convex subset of a uniformly convex and uniformly smooth Banach space E , and $\{T_i\}_{i=1}^\infty : C \rightarrow C$ be a family of closed and uniformly Lipschitz continuous and quasi- ϕ -asymptotically nonexpansive mappings with sequence $\{k_{n,i}\}_1^\infty \subset [1, \infty)$ such that $k_n := \sup_{i \geq 1} k_{n,i} \rightarrow 1$ ($n \rightarrow \infty$) and $F = \bigcap_{i=1}^\infty F(T_i) \neq \emptyset$. Let $\{x_n\}$ and $\{\alpha_n\}$ be the same sequences as given in Theorem 3.1 Then $\{x_n\} \rightarrow$ converges strongly to $\Pi_F x_0$.*

Because each quasi- ϕ nonexpansive mapping is a quasi- ϕ -asymptotically nonexpansive mapping with sequence $\{k_{n,i} = 1\}$, therefore we have the following

Corollary 3.4. *Let C be a nonempty bounded closed convex subset of a uniformly convex and uniformly smooth Banach space E and $\{T_i\}_{i=1}^\infty : C \rightarrow C$ be a family of closed and quasi- ϕ nonexpansive mappings such that $F = \bigcap_{i=1}^\infty F(T_i) \neq \emptyset$. Let $\{x_n\}$ be the sequence defined by:*

$$\begin{cases} x_0 \in C, & C_0 = C, \\ y_n = J^{-1}(\alpha_{n0} Jx_n + \sum_{i=1}^\infty \alpha_{ni} J T_i x_n), \\ C_{n+1} = \{z \in C_n : \phi(z, y_n) \leq \phi(z, x_n)\}, \\ x_{n+1} = \Pi_{C_{n+1}}(x_0), \end{cases} \tag{3.19}$$

where $\{\alpha_{ni}\} \subset [0, 1]$ is the sequence satisfying conditions (a), (b) in Theorem 3.1. Then $\{x_n\}$ converges strongly to $\Pi_F x_0$.

REFERENCES

- [1] K. Goebel, W.A. Kirk, *A fixed point theorem for asymptotically nonexpansive mappings*, Proc. Amer. Math. Soc. **35** (1972), 171–174.
- [2] K. Nakajo, W. Takahashi, *Strong convergence theorems for nonexpansive mappings and nonexpansive semigroups*, J. Math. Anal. Appl. **279** (2003), 372–379.
- [3] S.Y. Matsushita, W. Takahashi, *A strong convergence theorem for relatively nonexpansive mappings in a Banach space*, J. Approx. Theory **134** (2005), 257–266.
- [4] T.H. Kim, H.K. Xu, *Strong convergence of modified Mann iterations for asymptotically mappings and semigroups*, Nonlinear Anal. **64** (2006), 1140–1152.
- [5] H.Y. Zhou, G.L. Gao, *Convergence theorems of a modified hybrid algorithm for a family of quasi- ϕ -asymptotically nonexpansive mappings*, J. Appl. Math. Comput. DOI:10.1007/s12190-009-0263-4.
- [6] S. Kamimura, W. Takahashi, *Strong convergence of a proximal-type algorithm in a Banach space*, SIAM J. Optim. **13** (2002), 938–945.
- [7] Zhang Shi-sheng, *The generalized mixed equilibrium problem in Banach space*, Appl. Math. Mech. **30** (2009), 1105–1112.

J.F. Tang

Yibin University
Department of Mathematics
Yibin, Sichuan 644007, China

S.S. Chang
changss@yahoo.cn

Yibin University
Department of Mathematics
Yibin, Sichuan 644007, China

M. Liu

Yibin University
Department of Mathematics
Yibin, Sichuan 644007, China

J.A. Liu

Beijing Institute of Petro-Chemical Industry
Department of Mathematics and Physics
Beijing, 102617, China

Received: March 15, 2010.

Accepted: March 18, 2010.