

Discrete approximation of nonconvex hyperbolic optimal control problems with state constraints

by

I. Chrysosoverghi, A. Bacopoulos, J. Coletsos and B. Kokkinis

Department of Mathematics,
National Technical University, Athens, Greece

Abstract: We consider an optimal control problem for systems defined by nonlinear hyperbolic partial differential equations with state constraints. Since no convexity assumptions are made on the data, we also consider the control problem in relaxed form. We discretize both the classical and the relaxed problems by using a finite element method in space and a finite difference scheme in time, the controls being approximated by piecewise constant ones. We develop the existence theory and the necessary conditions for optimality, for the continuous and the discrete problems. Finally, we study the behaviour in the limit of discrete optimality, admissibility and extremality properties.

Keywords: Optimal control, nonlinear hyperbolic systems, discretization, existence theory, minimum principle, relaxed controls.

1. Introduction

Optimal control problems without any convexity assumptions on the data have no classical solutions in general. In order to prove the existence of optimal controls, the convexity of some extended velocity set is usually assumed, which is clearly unrealistic when nonlinear systems are involved. To overcome this difficulty, one has to relax, or convexify, the problem in some manner, and then work on the relaxed problem. As a result, relaxation theory has been extensively used, not only to prove existence theorems and derive necessary conditions for optimality, see Warga (1972), Ekeland (1972), Chrysosoverghi (1986), Fattorini (1994), Fattorini (1997), Roubiřek (1997), but also to develop approximation schemes, see Roubiřek (1991), Chrysosoverghi et al. (1993), and optimization methods, see Warga (1977), Teo et al. (1984), Chrysosoverghi et al. (1997).

Here, we consider an optimal distributed control problem for systems defined by nonlinear hyperbolic partial differential equations with several equality and inequality constraints (for hyperbolic systems, see also Bittner, 1975, Sloss

et al., 1995, Sadek et al., 1996). Under reasonable assumptions, the existence of optimal controls and the necessary conditions for optimality are established for the relaxed problem. We then discretize the classical (resp. relaxed) problem by using the Galerkin finite element method in space and the semi-implicit finite difference scheme in time for approximating the state equations, while the controls are approximated by piecewise constant classical (resp. relaxed) ones with respect to an independent partition of the space-time domain (see Cullum, 1971, and Casas, 1996, for discretizations of classical problems). This independent control discretization corresponds to the use of controls of simple and flexible form for numerical and/or engineering reasons. The discretization of both the classical and the relaxed continuous problems is motivated by the fact that in practice classical (resp. relaxed) optimization methods are usually applied to the classical (resp. relaxed) problem after some discretization. We then prove the existence of optimal controls for both the discrete classical and the discrete relaxed problems, and derive a piecewise minimum principle of optimality for the discrete relaxed problem. Finally, we study the behaviour in the limit of the above approximations. More precisely, we prove that, under appropriate assumptions, accumulation points of sequences of optimal discrete classical controls are optimal for the continuous relaxed problem, and that accumulation points of sequences of optimal (resp. admissible extremal) discrete relaxed controls are optimal (resp. admissible extremal) for the continuous relaxed problem.

2. The continuous optimal control problems

Let Ω be a bounded domain in \mathbf{R}^d with Lipschitz boundary Γ , and let $I := (0, T)$, $0 < T < \infty$. Consider the following nonlinear hyperbolic state equations

$$\partial^2 y / \partial t^2 + A(t)y = f(x, t, y(x, t), w(x, t)), \quad \text{in } Q := \Omega \times I, \quad (1)$$

$$y(x, t) = 0, \quad \text{in } \Sigma := \Gamma \times I, \quad (2)$$

$$y(x, 0) = y^0(x), \quad \text{in } \Omega, \quad (3)$$

$$(\partial y / \partial t)(x, 0) = y^1(x), \quad \text{in } \Omega, \quad (4)$$

where $A(t)$ is the second order differential operator

$$A(t)y := - \sum_{i,j=1}^d (\partial / \partial x_i)[a_{ij}(x, t)(\partial y / \partial x_j)].$$

The constraints on the control w are

$$w(x, t) \in U, \quad \text{in } Q,$$

where U is a compact subset of $\mathbf{R}^{d'}$. The constraints on the state and the control variables y, w are

$$J_m(w) := \int_Q g_m(x, t, y(x, t), w(x, t)) dx dt = 0, \quad 1 \leq m \leq p,$$

$$J_m(w) := \int_Q g_m(x, t, y(x, t), w(x, t)) dx dt \leq 0, \quad p < m \leq q,$$

and the cost functional is

$$J_0(w) := \int_Q g_0(x, t, y(x, t), w(x, t)) dx dt,$$

where $y := y_w$ is the solution of (1-4) for the control w . The optimal control problem is to minimize $J_0(w)$ subject to the above constraints.

Since such problems have no classical solutions in general, without additional convexity assumptions on the data, it is standard (see Warga, 1972) to work on the so-called relaxed form of the problem, which we define below.

Define the set of classical controls

$$W := \{w : (x, t) \longrightarrow w(x, t) \mid w \text{ measurable from } \bar{Q} \text{ to } U\},$$

and the set of relaxed controls

$$R := \{r : (x, t) \longrightarrow r(x, t) \mid r \text{ weakly measurable from } \bar{Q} \text{ to } M_1(U)\},$$

where the set $M_1(U)$ of probability measures on U is a subset of the space $M(U) \approx C(U)^*$ of Radon measures on U , and has here the relative weak star topology. We have

$$R \subset L_w^\infty(\bar{Q}, M(U)) \approx L^1(\bar{Q}, C(U))^* \approx B(\bar{Q}, U)^*,$$

where $L_w^\infty(\bar{Q}, M(U))$ is the set of (equivalence classes of) functions from \bar{Q} to $M(U)$ which are measurable w.r.t. a weak norm topology on $M(U)$ (which coincides on $M_1(U)$ with the relative weak star topology) and essentially bounded w.r.t. the strong dual norm on $M(U)$, and $B(\bar{Q}, U)$ is the space of Caratheodory functions on $\bar{Q} \times U$ in the sense of Warga (1972). The subset R is endowed with relative weak star topology. The sets $M_1(U)$ and R are convex and, with their respective topology, metrizable and compact. For $\phi \in B(\bar{Q}, U)$, and $r \in \text{span}(R)$, we use the notation

$$\phi(x, t, r(x, t)) := \int_U \phi(x, t, u) r(x, t)(du).$$

Note that this expression is linear in r . A sequence $\{r_k\}$ converges to r in R if

$$\lim_{k \rightarrow \infty} \int_Q \phi(x, t, r_k(x, t)) dx dt = \int_Q \phi(x, t, r(x, t)) dx dt,$$

for every $\phi \in B(\overline{Q}, U)$, or equivalently for every $\phi \in C(\overline{Q} \times U)$. In addition, we identify every classical control $w(\cdot, \cdot)$ with its associated Dirac relaxed control $\delta_{w(\cdot, \cdot)}$. Thus, we have $W \subset R$, and it is proved in Warga (1972) that W is dense in R .

We denote by (\cdot, \cdot) and $|\cdot|$ the inner product and norm in $L^2(\Omega)$, by $((\cdot, \cdot))$ and $\|\cdot\|$ the usual inner product and the norm in the Sobolev space $V := H_0^1(\Omega)$ and by $\langle \cdot, \cdot \rangle$ the duality bracket between V and its dual V^* . Define the family of bilinear forms on V

$$a(t, v, w) := \sum_{i,j=1}^d \int_{\Omega} a_{ij}(x, t) (\partial v / \partial x_j) (\partial w / \partial x_i) dx.$$

In order to define the relaxed as well as the classical solutions of our problem, we shall first interpret the state equation in the following weak and relaxed form:

$$\langle y'', v \rangle + a(t, y, v) = \int_{\Omega} f(x, t, y(x, t), r(x, t)) v(x) dx, \quad (5)$$

for every $v \in V$, a.e. in I ,

$$y(x, 0) = y^0(x), \quad \text{in } \Omega, \quad (6)$$

$$y'(x, 0) = y^1(x), \quad \text{in } \Omega, \quad (7)$$

where the derivatives are taken in the sense of distributions (cf. Lions, 1969, p. 115). Note that, accordingly to our notation, the relaxed control r appears here in mean-value form. Defining the functionals

$$J_m(r) := \int_Q g_m(x, t, y(x, t), r(x, t)) dx dt, \quad 0 \leq m \leq q, \quad (8)$$

the continuous relaxed optimal control problem (CRP) is to minimize $J_0(r)$ subject to the constraints $r \in R$, $J_m(r) = 0$, $1 \leq m \leq p$, and $J_m(r) \leq 0$, $p < m \leq q$, where $y = y_r$ is the (unique) solution of (2,5,6,7). The continuous classical problem (CCP) is the problem CRP with additional constraint $r \in W$.

We suppose that the operators $A(t)$ satisfy the following conditions

$$\sum_{i,j=1}^d a_{ij}(x, t) z_i z_j \geq \alpha \sum_{i=1}^d z_i^2, \quad (x, t) \in \overline{Q}, \quad z_i \in \mathbf{R}, \quad 1 \leq i \leq d, \quad \text{with } \alpha > 0,$$

$$a_{ij} \in C^1(\overline{I}, L^\infty(\Omega)), \quad i, j = 1, \dots, d,$$

$$a_{ij} = a_{ji}, \quad i, j = 1, \dots, d,$$

which imply that

$$|a(t, v, w)| \leq \alpha_1 \|v\| \|w\|, \quad t \in \overline{I}, \quad v, w \in V,$$

$$a(t, v, v) \geq \alpha_2 \|v\|^2, \quad t \in \bar{I}, \quad v \in V,$$

for some $\alpha_1 \geq 0$, $\alpha_2 > 0$, and that $a(t, v, w)$ is symmetric.

We suppose also that the function f is defined on $\bar{Q} \times \mathbf{R} \times U$, measurable for fixed y, u , continuous for fixed x, t , and satisfies

$$|f(x, t, y, u)| \leq F(x, t) + \beta|y|, \quad \text{for every } x, t, y, u,$$

$$\text{with } F \in L^2(Q), \quad \beta \geq 0,$$

$$|f(x, t, y_1, u) - f(x, t, y_2, u)| \leq L|y_1 - y_2|, \quad \text{for every } x, t, y_1, y_2, u.$$

Then, for every $r \in R$, $y^0 \in V$ and $y^1 \in L^2(\Omega)$, it can be proved that equations (2,5,6,7) have a unique solution $y = y_r$, such that $y \in L^\infty(I, V)$, $y' \in L^\infty(I, L^2(\Omega))$ and $y'' \in L^2(I, V^*)$ (see Lions, 1972, Chap. 4, for the uniqueness in the linear case, and proof of Lemma 2.1 below, with fixed control). It follows that y is essentially equal to a function in $C(\bar{I}, L^2(\Omega))$, that y' is essentially equal to a function in $C(\bar{I}, V^*)$, and thus the initial conditions (6,7) make sense.

LEMMA 2.1 *The mapping $r \rightarrow y_r$, from R to $L^2(Q)$, is continuous.*

Proof. Let $r \in R$ be a fixed relaxed control. Let $\{r_k\}$ be any sequence converging to r in R , and set $y_k := y_{r_k}$. Note that, since the bilinear form $a(t, \cdot, \cdot)$ is defined on V only, we cannot directly replace v by $y'_k(t) \in L^2(\Omega)$ in (5). To overcome this difficulty, we shall use the following approximation. Since V is separable, there exists a sequence $\{v_i\}_{i=1}^\infty$ such that the elements v_1, \dots, v_n are linearly independent for every n , and the set $\text{span}(\{v_i\}_{i=1}^\infty)$ is dense in V . For every k and n , define the approximate solution y_{nk}

$$y_{nk}(x, t) := \sum_{j=1}^n \zeta_{jnk}(t) v_j(x),$$

which satisfies

$$(y_{nk}''(t), v_i) + a(t, y_{nk}(t), v_i) = (f(t, y_{nk}, r_k), v_i), \quad i = 1, \dots, n, \quad \text{a.e. in } I, \quad (9)$$

$$y_{nk}(0) = y_{0n} = \sum_{j=1}^n \xi_{jn}^0 v_j, \quad (10)$$

$$y_{nk}'(0) = y_{1n} = \sum_{j=1}^n \xi_{jn}^1 v_j, \quad (11)$$

where $y_{0n} \rightarrow y^0$ (resp. $y_{1n} \rightarrow y^1$) in V (resp. $L^2(\Omega)$) strongly. For example, we can choose y_{0n} (resp. y_{1n}) to be the projection of y^0 (resp. y^1) onto the

space $\text{span}(\{v_1, \dots, v_n\})$ in V (resp. $L^2(\Omega)$). By our assumptions, for each n, k , equations (9,10,11) reduce to a system of linear ordinary differential equations in the ζ_{jnk} , which has a unique solution, with ζ_{jnk} and ζ'_{jnk} absolutely continuous on \bar{I} (see Warga, 1972, Ch. II). Hence, in particular, $y_{nk}, y'_{nk} \in C(\bar{I}, V)$. From equation (9), with v_i replaced by y'_{nk} , we obtain

$$\begin{aligned} & (d/dt)[|y'_{nk}(t)|^2 + a(t, y_{nk}(t), y_{nk}(t))] - a_t(t, y_{nk}(t), y_{nk}(t)) \\ &= 2(f(t, y_{nk}(t), r_k(t)), y'_{nk}(t)), \end{aligned}$$

where $a_t(t, v, w)$ is defined by replacing the coefficients a_{ij} by $\partial a_{ij}/\partial t$ in $a(t, v, w)$. Integrating on $[0, t]$, we get

$$\begin{aligned} & |y'_{nk}(t)|^2 + a(t, y_{nk}(t), y_{nk}(t)) = |y'_{1n}|^2 + a(0, y_{0n}, y_{0n}) \\ & + \int_0^t a_t(s, y_{nk}(s), y_{nk}(s)) ds + 2 \int_0^t (f(s, y_{nk}(s), r_k(s)), y'_{nk}(s)) ds. \end{aligned}$$

By our assumptions, it follows easily that

$$\begin{aligned} & |y'_{nk}(t)|^2 + \|y_{nk}(t)\|^2 \leq c \left[|y'_{1n}|^2 + \|y_{0n}\|^2 + \int_0^T |F(s)|^2 ds \right] \\ & + c' \int_0^t \left[|y'_{nk}(s)|^2 + \|y_{nk}(s)\|^2 \right] ds. \end{aligned}$$

Hence, by Gronwall's inequality

$$|y'_{nk}(t)|^2 + \|y_{nk}(t)\|^2 \leq c, \quad t \in \bar{I},$$

which shows, in particular, that the double sequence $\{y_{nk}\}$ (resp. $\{y'_{nk}\}$) is bounded in $L^2(I, V)$ (resp. $L^2(Q)$). Now, let $\{y_{n_\mu k_\mu}\}_{\mu=1}^\infty$ be any subsequence such that $n_\mu \rightarrow \infty$ and $k_\mu \rightarrow \infty$. By the Alaoglu-Bourbaki theorem, there exists a subsequence $\{y_{n_\mu k_\mu}\}$ (same notation) such that

$$y_{n_\mu k_\mu} \rightarrow z \text{ in } L^2(I, V) \text{ weakly,}$$

$$y'_{n_\mu k_\mu} \rightarrow \bar{z} \text{ in } L^2(Q) \text{ weakly.}$$

We have, for every $v \in V$ and $\phi \in C_c^\infty(I)$

$$\int_0^T (y'_{n_\mu k_\mu}, v) \phi(t) dt = - \int_0^T (y_{n_\mu k_\mu}, v) \phi'(t) dt,$$

and passing to the limit we see that $\bar{z} = z'$ (Lemma 1.1 in Temam, 1977, p. 250). By the Aubin compactness theorem (see Temam, 1977, p. 271), we can suppose also that

$$y_{n_\mu k_\mu} \rightarrow z \text{ in } L^2(Q) \text{ strongly.}$$

We can then pass to the limit (in μ) in the approximate equations (9,10,11) in weak integrated form (see Lions, 1972), using here Proposition 2.1, from Chrysosoverghi (1986), for the term containing f , to show that $z = y_r$. Since the limit y_r is unique, it follows by contradiction that $y_{nk} \rightarrow y_r$, in $L^2(Q)$ strongly, as $n, k \rightarrow \infty$. Note that the above proof shows, in particular, that for each k , i.e. keeping the control r_k fixed, we have $y_{nk} \rightarrow y_k := y_{r_k}$ in $L^2(Q)$ strongly, as $n \rightarrow \infty$. These convergences imply that, for given ε , there exist N_k , for each k , such that

$$\|y_k - y_{nk}\|_{L^2(Q)} \leq \varepsilon/2, \text{ for every } n \geq N_k,$$

and N such that

$$\|y_{nk} - y_r\|_{L^2(Q)} \leq \varepsilon/2, \text{ for every } n \geq N \text{ and } k \geq N.$$

Hence, for each $k \geq N$, by choosing some $n \geq \max(N, N_k)$, we get

$$\|y_k - y_r\|_{L^2(Q)} \leq \varepsilon.$$

Therefore $y_k \rightarrow y_r$ in $L^2(Q)$ strongly, which proves the lemma, since R is metrizable. ■

In order to prove the existence of an optimal relaxed control, we suppose in addition that the functions g_m , $0 \leq m \leq q$, are measurable for fixed y, u , continuous for fixed x, t , and satisfy

$$|g_m(x, t, y, u)| \leq G_m(x, t) + \gamma_m |y|^2, \text{ for every } x, t, y, u,$$

$$\text{with } G_m \in L^1(Q), \gamma_m \geq 0, 0 \leq m \leq q.$$

LEMMA 2.2 *The functionals J_m , $0 \leq m \leq q$, are continuous on R .*

Proof. Follows from Lemma 2.1 and Proposition 2.1. of Chrysosoverghi (1986). ■

THEOREM 2.1 *If there exists an admissible control, i.e. a control satisfying the constraints, then there exists an optimal control for the CRP.*

Proof. By Lemma 2.2, the set $R_A \subset R$ of admissible controls is closed, hence compact, and the functional J_0 is continuous. The theorem follows. ■

In order to derive necessary conditions for optimality, we suppose in addition that f'_y and g'_{my} , $0 \leq m \leq q$, exist, are measurable for fixed y, u , continuous for fixed x, t , and satisfy

$$|g'_{my}(x, t, y, u)| \leq G_{1m}(x, t) + \gamma_{1m} |y|, \text{ for every } x, t, y, u,$$

$$\text{with } G_{1m} \in L^2(Q), \gamma_{1m} \geq 0, 0 \leq m \leq q.$$

Since f is Lipschitzian, we have also

$$|f'_y(x, t, y, u)| \leq L, \text{ for every } x, t, y, u.$$

LEMMA 2.3 For $\mu \geq 0$, $r \in R$, $\tilde{r}_i \in R$, $i = 0, \dots, \mu$, $\theta = (\theta_0, \dots, \theta_\mu)$, with $\theta_i > 0$, $i = 0, \dots, \mu$, and $\sum_{i=0}^{\mu} \theta_i \leq 1$, set $r_\theta := r + \sum_{i=0}^{\mu} \theta_i(\tilde{r}_i - r)$, $y := y_r$, $y_\theta := y_{r_\theta}$, $\Delta y_\theta := y_\theta - y$. Then

$$\|\Delta y_\theta\|_{L^2(Q)} \leq c\|\theta\|_\infty,$$

where $\|\theta\|_\infty = \max_i |\theta_i|$.

Proof. Let $y_{n\theta}$ (resp. y_n) be the solution of equations (9,10,11), where y_{nk} is replaced by $y_{n\theta}$ (resp. y_n) and r_k by r_θ (resp. r). Setting $\Delta y_{n\theta} := y_{n\theta} - y_n$, we have

$$\begin{aligned} & (\Delta y_{n\theta}''(t), \Delta y_{n\theta}'(t)) + a(t, \Delta y_{n\theta}(t), \Delta y_{n\theta}'(t)) \\ &= (f(t, y_{n\theta}, r_\theta) - f(t, y_n, r), \Delta y_{n\theta}'(t)) \\ &= (f(t, y_{n\theta}, r_\theta) - f(t, y_n, r_\theta), \Delta y_{n\theta}(t)) \\ &+ \sum_{i=0}^{\mu} \theta_i (f(t, y_n, \tilde{r}_i - r_i), \Delta y_{n\theta}'(t)). \end{aligned}$$

Since f is Lipschitzian, we deduce, similarly to the proof of Lemma 2.1, c denoting various constants, that

$$\begin{aligned} & |\Delta y_{n\theta}'(t)|^2 + \|\Delta y_{n\theta}(t)\|^2 \\ & \leq c \int_0^t [|\Delta y_{n\theta}'(s)|^2 + \|\Delta y_{n\theta}(s)\|^2] ds + c\|\theta\|_\infty^2 \sum_{i=0}^{\mu} \int_0^T |f(y_n, \tilde{r}_i - r)|^2 ds, \end{aligned}$$

hence, using Gronwall's inequality

$$\begin{aligned} & \|\Delta y_{n\theta}\|_{L^2(Q)}^2 \leq c\|\Delta y_{n\theta}\|_{L^\infty(I, L^2(\Omega))}^2 \leq c\|\Delta y_{n\theta}\|_{L^\infty(I, V)}^2 \\ & \leq c\|\theta\|_\infty^2 \sum_{i=0}^{\mu} \int_0^T |f(y_n, \tilde{r}_i - r)|^2 ds \\ & \leq c\|\theta\|_\infty^2 (\mu + 1) [\|F\|_{L^2(Q)} + \beta\|y_n\|_{L^2(Q)}]^2. \end{aligned}$$

By Lemma 2.1, with fixed control, $y_n \rightarrow y$, $y_{n\theta} \rightarrow y_\theta$ and $\Delta y_{n\theta} \rightarrow \Delta y_\theta$, in $L^2(Q)$. Therefore

$$\|\Delta y_\theta\|_{L^2(Q)} \leq c\|\theta\|_\infty.$$

LEMMA 2.4 Dropping the index m , with the notation of Lemma 2.3, we have

$$\begin{aligned} & J[r + \sum_{i=0}^{\mu} \theta_i(\tilde{r}_i - r)] - J(r) \\ &= \sum_{i=0}^{\mu} \theta_i \int_Q H(x, t, y(x, t), \phi(x, t), \tilde{r}_i(x, t) - r(x, t)) dx dt + o(\|\theta\|_\infty), \end{aligned}$$

where, for each function g , the general Hamiltonian H is defined by

$$H(x, t, y, \phi, u) := \phi f(x, t, y, u) + g(x, t, y, u), \quad (12)$$

and the general adjoint state $\phi := \phi_r$ satisfies

$$\phi'' + A(t)\phi = f'_y(y, r)\phi + g'_y(y, r), \quad \text{in } Q, \quad (13)$$

$$\phi(x, t) = 0, \quad \text{in } \Sigma, \quad (14)$$

$$\phi(x, T) = \phi'(x, T) = 0, \quad \text{in } \Omega, \quad (15)$$

where $y := y_r$. In particular, for $\mu = 0$, $r, \tilde{r} \in R$, the directional derivative of J is

$$\begin{aligned} DJ(r, \tilde{r} - r) &:= \lim_{\theta \rightarrow 0^+} [J(r + \theta(\tilde{r} - r)) - J(r)]/\theta \\ &= \int_Q H(x, t, y(x, t), \phi(x, t), \tilde{r}(x, t) - r(x, t)) dx dt. \end{aligned}$$

Proof. Under our assumptions on g and g'_y , for given $r \in R$, it can be proved (similarly to Warga (1972), II.6.7, II.6.8 and VIII.2.2, using essentially the mean-value theorem and the Lebesgue dominated convergence theorem) that the functional Ψ , with

$$\Psi(y) := \int_Q g(x, t, y, r) dx dt,$$

defined on $L^2(Q)$, has Fréchet derivative Ψ' , with

$$\Psi'(y)\Delta y = \int_Q g'_y(x, t, y, r)\Delta y dx dt.$$

By Lemma 2.3, we have

$$\begin{aligned} J(r_\theta) - J(r) &= \int_Q [g(y_\theta, r_\theta) - g(y, r_\theta) + g(y, r_\theta) - g(y, r)] dx dt \\ &= \int_Q g'_y(y, r)\Delta y_\theta dx dt + \sum_{i=0}^{\mu} \theta_i \int_Q g'_y(y, \tilde{r}_i - r)\Delta y_\theta dx dt \\ &\quad + o(\|\theta\|_\infty) + \sum_{i=0}^{\mu} \theta_i \int_Q g(y, \tilde{r}_i - r) dx dt \\ &= \int_Q g'_y(y, r)\Delta y_\theta dx dt + \sum_{i=0}^{\mu} \theta_i \int_Q g(y, \tilde{r}_i - r) dx dt + o(\|\theta\|_\infty). \end{aligned}$$

Since $\Delta y_\theta(0) = \Delta y'_\theta(0) = \phi(T) = \phi'(T) = 0$, integrating by parts, we obtain from the state equation

$$- \int_0^T (\Delta y'_\theta, \phi') dt + \int_0^T a(t, \Delta y_\theta, \phi) dt$$

$$= \int_Q f'_y(y, r) \Delta y_\theta \phi dx dt + \sum_{i=0}^{\mu} \theta_i \int_Q f(y, \tilde{r}_i - r) \phi dx dt + o(\|\theta\|_\infty),$$

and from the adjoint equation

$$\begin{aligned} & - \int_0^T (\phi', \Delta y'_\theta) dt + \int_0^T a(t, \phi, \Delta y_\theta) dt \\ & = \int_Q f'_y(y, r) \phi \Delta y_\theta dx dt + \int_Q g'_y(y, r) \Delta y_\theta dx dt, \end{aligned}$$

and the lemma follows. ■

The following theorem states necessary conditions for optimality.

THEOREM 2.2 *If $r \in R$ is optimal for either the CRP or the CCP, then r is extremal, i.e. there exist multipliers $\lambda_m \in \mathbf{R}$, $0 \leq m \leq q$, with $\lambda_0 \geq 0$, $\lambda_m \geq 0$, $p < m \leq q$, and $\sum_{m=0}^q |\lambda_m| \neq 0$, such that*

$$\begin{aligned} & H(x, t, y(x, t), \phi(x, t), r(x, t)) \\ & = \min_{u \in U} H(x, t, y(x, t), \phi(x, t), u), \quad \text{a.e. in } Q, \end{aligned} \tag{16}$$

and

$$\lambda_m J_m(r) = 0, \quad p < m \leq q, \tag{17}$$

where H and ϕ are defined by (12) and (13,14,15), with g replaced by $\sum_{m=0}^q \lambda_m g_m$.

Proof. Let

$$S := \{\theta \in \mathbf{R}_+^{p+1} \mid \sum_{m=0}^p \theta_m \leq 1\}.$$

By Lemma 2.4, the functions

$$\theta \longrightarrow J_m(r_\theta), \quad 0 \leq m \leq q,$$

from S to \mathbf{R} , have a derivative at 0 (see Warga, 1972), and by Lemma 2.2, the functionals J_m are continuous on R . If r is optimal for the CRP or the CCP, then, by Theorem V.3.2 in Warga (1972), p. 310, there exist multipliers as above such that

$$\sum_{m=0}^q \lambda_m D J_m(r, \tilde{r} - r) \geq 0, \quad \text{for every } \tilde{r} \in R,$$

which is equivalent to (16) (see Warga, 1972, p. 360), and such that (17) holds. ■

We have also the following continuity result, whose proof is similar to that of Lemma 2.1.

LEMMA 2.5 *The mapping $r \rightarrow \phi_r$, from R to $L^2(Q)$, is continuous.*

3. The discrete optimal control problems

We shall now discretize our continuous optimal control problems. For simplicity, we suppose that the domain Ω is a polyhedron.

For every integer n , let $\{S_i^n\}_{i=1}^{M(n)}$ be an admissible regular triangulation (see Temam, 1977, p. 73) of $\bar{\Omega}$ into closed d -simplices, $\{I_j^n\}_{j=0}^{N(n)-1}$ a subdivision of the interval \bar{I} into $N(n)$ intervals $I_j^n := [t_j^n, t_{j+1}^n]$, of equal length $\Delta t = T/N$, and $\{B_k^n\}_{k=1}^{P(n)}$ a partition of \bar{Q} into $P(n)$ Borel subsets with $\max_k [\text{diam}(B_k^n)] \rightarrow 0$, as $n \rightarrow \infty$. Set $Q_{ij}^n := S_i^n \times I_j^n$. For example, the B_k^n may be unions of some of the Q_{ij}^n . Let $V^n \subset V$ be the subspace of functions which are continuous on $\bar{\Omega}$ and affine on each S_i^n . Let $R^n \subset R$ be the set of piecewise constant relaxed controls

$$R^n := \{r^n \in R \mid r^n(x, t) := r_k^n \in M_1(U), \text{ on } B_k^n, k = 1, \dots, P\},$$

and $W^n := R^n \cap W$ the set of piecewise constant classical controls.

For given values $v_j^n, j = 0, \dots, N$, in a vector space, we define the following functions, a.e. on \bar{I}

$$v_-^n(t) := v_j^n, t \in I_j^n, j = 0, \dots, N-1,$$

$$v_+^n(t) := v_{j+1}^n, t \in I_j^n, j = 0, \dots, N-1,$$

$v_\wedge^n(t) :=$ the function which is affine on each I_j^n , such that

$$v_\wedge^n(t_j^n) = v_j^n, j = 0, \dots, N.$$

For a given discrete relaxed control $r^n := (r_1^n, \dots, r_k^n, \dots, r_P^n)$, the corresponding discrete state $y^n := (y_0^n, \dots, y_j^n, \dots, y_N^n)$ is given by

$$y_{j+1}^n - y_j^n = \Delta t z_{j+1}^n, j = 0, \dots, N-1, \quad (18)$$

$$(1/\Delta t)(z_{j+1}^n - z_j^n, v) + a(t_{j+1}^n, y_{j+1}^n, v) = (f_j^n, v),$$

for every $v \in V^n, j = 0, \dots, N-1,$ (19)

$$y_0^n, z_0^n \text{ given, } y_j^n, z_j^n \in V^n, j = 0, \dots, N, \quad (20)$$

$$f_j^n(x) := (1/\Delta t) \int_{I_j^n} f(x, t, y_j^n(x), r^n(x, t)) dt, j = 0, \dots, N-1. \quad (21)$$

Choosing a basis $\{v_\mu\}$ in V^n and writing

$$y_{j+1}^n = \sum_{\mu} c_\mu v_\mu, z_{j+1}^n = \sum_{\mu} \hat{c}_\mu v_\mu,$$

we reduce equations (18,19) to a linear system of the form

$$c = \Delta t c' + b, \quad (22)$$

$$(D + \Delta t^2 E)c' = b', \quad (23)$$

where D and E are the corresponding mass and stiffness matrices, respectively. This system is regular since (23) has a positive definite matrix. Defining the discrete functionals

$$J_m^n(r^n) := \int_Q g_m(x, t, y^n(x, t), r^n(x, t)) dx dt, \quad 0 \leq m \leq q,$$

the discrete constraints are $r^n \in R^n$, either of the two following constraints

$$|J_m^n(r^n)| \leq \varepsilon_m^n, \quad 1 \leq m \leq p, \quad (24)$$

$$J_m^n(r^n) = \varepsilon_m^n, \quad 1 \leq m \leq p, \quad (25)$$

and

$$J_m^n(r^n) \leq \varepsilon_m^n, \quad \varepsilon_m^n \geq 0, \quad p < m \leq q, \quad (26)$$

where the ε_m^n , $1 \leq m \leq q$, are given numbers. The discrete relaxed problem DRP^n (resp. DRP'^n) is to minimize $J_0^n(r^n)$ subject to the above constraints, case (24) (resp. (25)). The discrete classical problem DCP^n is the problem DRP^n with the additional constraint $r^n \in W^n$.

THEOREM 3.1 *If there exists an admissible control for any of the above discrete problems, then there exists an optimal control for this problem.*

Proof. We first remark that the convergence in $R^n = [M_1(U)]^P$ implies the convergence in R , hence in each set R_{ij} of restrictions $r|_{\overline{Q}_{ij}}$, for $r \in R$. The continuity of the mapping $r^n \rightarrow y^n$ is proved by induction on j , using system (22,23), and Proposition 2.1, from Chrysoverghi (1986). The same proposition shows that the functionals J_m^n are continuous. The existence of an optimal control for the DRP^n and the DRP'^n follows then from the compactness of the set R_A^n of admissible discrete relaxed controls. The existence for the DCP^n is easily proved, using here the ordinary topology of W^n . ■

LEMMA 3.1 *Dropping the index m , define the general discrete relaxed adjoint state $\phi^n = (\phi_0^n, \dots, \phi_j^n, \dots, \phi_N^n)$ by*

$$\phi_{j+1}^n - \phi_j^n = \Delta t \psi_j^n, \quad j = N-1, \dots, 0, \quad (27)$$

$$(1/\Delta t)(\psi_{j+1}^n - \psi_j^n, v) + a(t_j^n, \phi_j^n, v) = (f'_{y_j} \phi_{j+1}^n, v) + (g'_{y_j}, v),$$

$$\text{for every } v \in V^n, j = N - 1, \dots, 0, \tag{28}$$

$$\phi_N^n = \psi_N^n = 0, \phi_j^n, \psi_j^n \in V^n, j = 0, \dots, N,$$

$$f'_{y_j^n}(x) = (1/\Delta t) \int_{I_j^n} f'_y(x, t, y_j^n, r^n) dt, j = 0, \dots, N - 1, \tag{29}$$

$$g'_{y_j^n}(x) = (1/\Delta t) \int_{I_j^n} g'_y(x, t, y_j^n, r^n) dt, j = 0, \dots, N - 1. \tag{30}$$

The directional derivative of the functional J^n on R^n is given by

$$DJ^n(r^n, \tilde{r}^n - r^n) = \int_Q H(x, t, y_-^n, \phi_+^n, \tilde{r}^n - r^n) dx dt,$$

where H is defined by (12).

Proof. (Outline.) The proof parallels that of the continuous case (Lemmas 2.3 and 2.4). The discrete analogue of Lemma 2.3 is first proved similarly to Lemma 4.2 below. The analogue of Lemma 2.4 is then proved by interpreting the discrete estimates and the discrete equations in terms of y_-^n, y_+^n, y_\wedge^n , etc... and using the discrete integration by parts formula

$$\int_0^T ((v_\wedge^n)', w_+^n) dt = - \int_0^T (v_-^n, (w_\wedge^n)') dt + (v_N^n, w_N^n) - (v_0^n, w_0^n), \tag{31}$$

which is directly verified. ■

THEOREM 3.2 *If r^n is optimal for the DRP^n , then r^n is extremal, i.e. there exist multipliers $\lambda_m^n \in \mathbf{R}$, $0 \leq m \leq q$, with $\lambda_0^n \geq 0, \lambda_m^n \geq 0, p < m \leq q$, and $\sum_{m=0}^q |\lambda_m^n| = 1$, such that*

$$\begin{aligned} & \int_{B_k^n} H(x, t, y_-^n, \phi_+^n, r_k^n) dx dt = \\ & \min_{u \in U} \int_{B_k^n} H(x, t, y_-^n, \phi_+^n, u) dx dt, k = 1, \dots, P, \end{aligned} \tag{32}$$

and

$$\lambda_m^n [J_m^n(r^n) - \varepsilon_m^n] = 0, p < m \leq q, \tag{33}$$

where H, y^n, ϕ^n are given by (12), (18), (19,20,21) and (27-30) with g replaced by $\sum \lambda_m^n g_m$.

Proof. Follows from Theorem V.2.3 in Warga (1972), p. 303, noting that the inequality

$$\sum_{m=0}^q \lambda_m^n D J_m^n(r^n, \tilde{r}^n - r^n) \geq 0, \quad \text{for every } \tilde{r}^n \in R^n,$$

or

$$\sum_{k=1}^P \int_{B_k^n} H(x, t, y_-^n, \phi_+^n, \tilde{r}_k^n - r_k^n) dx dt \geq 0, \quad \text{for every } \tilde{r}^n \in R^n,$$

is equivalent to (32), since the \tilde{r}_k^n are independent. Relations (33) are the transversality conditions. ■

4. Convergence

In this section we study the behaviour of the discrete problems in the limit, as $n \rightarrow \infty$. We first state the following control approximation lemma which is proved in Chrysoverghi et al. (1993); see also Roubiřek (1991).

LEMMA 4.1 *For every $r \in R$, there exists a sequence $\{u^n \in W^n\}$ such that $u^n \rightarrow r$ in R .*

LEMMA 4.2 (Stability) *For every $r^n \in R^n$, if $\|y_0^n\|$ and $|z_0^n|$ remain bounded and Δt is sufficiently small, then*

$$\|y_j^n\| \leq c, \quad j = 0, \dots, N, \quad (34)$$

$$|z_j^n| \leq c, \quad j = 0, \dots, N, \quad (35)$$

$$\sum_{j=0}^{N-1} \|y_{j+1}^n - y_j^n\|^2 \leq c, \quad (36)$$

$$\sum_{j=0}^{N-1} |z_{j+1}^n - z_j^n|^2 \leq c. \quad (37)$$

Proof. It is easily proved (see Chrysoverghi et al., 1993) that

$$|f_j^n| \leq F_j^n + \beta |y_j^n|,$$

where

$$F_j^n := \left((1/\Delta t) \int_{\Omega \times I_j^n} F^2 dx dt \right)^{1/2}.$$

We have, since $y_{j+1}^n - y_j^n = \Delta t z_{j+1}^n$

$$\begin{aligned} & a(t_{j+1}^n, y_{j+1}^n, y_{j+1}^n) - a(t_j^n, y_j^n, y_j^n) \\ &= \Delta t a(t_{j+1}^n, y_{j+1}^n, z_{j+1}^n) + \Delta t a(t_{j+1}^n, z_{j+1}^n, y_j^n) + \\ & a(t_{j+1}^n, y_j^n, y_j^n) - a(t_j^n, y_j^n, y_j^n) \\ &= 2\Delta t a(t_{j+1}^n, y_{j+1}^n, z_{j+1}^n) - \Delta t a(t_{j+1}^n, y_{j+1}^n, z_{j+1}^n) + \Delta t a(t_{j+1}^n, z_{j+1}^n, y_j^n) \\ & \quad + a(t_{j+1}^n, y_j^n, y_j^n) - a(t_j^n, y_j^n, y_j^n) \\ &= 2\Delta t a(t_{j+1}^n, y_{j+1}^n, z_{j+1}^n) - \Delta t^2 a(t_{j+1}^n, z_{j+1}^n, z_{j+1}^n) + \\ & a(t_{j+1}^n, y_j^n, y_j^n) - a(t_j^n, y_j^n, y_j^n). \end{aligned}$$

The discrete state equation yields

$$(z_{j+1}^n - z_j^n, z_{j+1}^n) + \Delta t a(t_{j+1}^n, y_{j+1}^n, z_{j+1}^n) = \Delta t (f_j^n, z_{j+1}^n).$$

Hence, by the assumptions and the mean value theorem

$$\begin{aligned} & |z_{j+1}^n|^2 - |z_j^n|^2 + |z_{j+1}^n - z_j^n|^2 + a(t_{j+1}^n, y_{j+1}^n, y_{j+1}^n) \\ & - a(t_j^n, y_j^n, y_j^n) + \Delta t^2 a(t_{j+1}^n, z_{j+1}^n, z_{j+1}^n) \\ & \leq |a(t_{j+1}^n, y_j^n, y_j^n) - a(t_j^n, y_j^n, y_j^n)| + 2\Delta t |(f_j^n, z_{j+1}^n)| \\ & \leq c' \Delta t \|y_j^n\|^2 + c\Delta t ((F_j^n)^2 + \|y_j^n\|^2 + |z_j^n|^2 + |z_{j+1}^n - z_j^n|^2). \end{aligned}$$

Assuming that $c\Delta t \leq 1/2$ (where c does not depend on the triangulation of Ω), and summing over j , we obtain

$$\begin{aligned} & |z_{i+1}^n|^2 + (1/2) \sum_{j=0}^i |z_{j+1}^n - z_j^n|^2 + \alpha_2 \|y_{i+1}^n\|^2 + \alpha_2 \sum_{j=0}^i \|y_{j+1}^n - y_j^n\|^2 \\ & \leq |z_0^n|^2 + \alpha_1 \|y_0^n\|^2 + c \|F\|_{L^2(Q)}^2 + c'' \Delta t \sum_{j=0}^i (\|y_j^n\|^2 + |z_j^n|^2), \end{aligned}$$

and the lemma follows from the discrete Gronwall inequality (see Lees, 1960). \blacksquare

From now on, we choose (for example) y_0^n to be the projection of y^0 onto V^n in V and z_0^n the projection of y^1 onto V^n in $L^2(\Omega)$. This implies here that $y_0^n \rightarrow y^0$ in V strongly and that $z_0^n \rightarrow y^1$ in $L^2(\Omega)$ strongly.

LEMMA 4.3 (Convergence) *If $r^n \rightarrow r$ in R , then the corresponding discrete states y_+^n, y_+^n, y_-^n converge to y_r in $L^2(Q)$ strongly, as $n \rightarrow \infty$.*

Proof. Since $\|y_0^n\|$ and $|z_0^n|$ are bounded, inequalities (34), (35) show that y_-^n, y_+^n, y_-^n are bounded in $L^2(I, V)$ and that z_-^n, z_+^n, z_-^n are bounded in $L^2(Q)$. By inequalities (36), (37), we have

$$y_+^n - y_-^n \rightarrow 0, \quad \text{in } L^2(I, V) \text{ and in } L^2(Q) \text{ strongly,}$$

$$z_+^n - z_-^n \rightarrow 0, \quad \text{in } L^2(Q) \text{ strongly.}$$

It follows easily that

$$y_+^n - y_\lambda^n \rightarrow 0, \text{ in } L^2(Q) \text{ strongly,}$$

$$z_+^n - z_\lambda^n \rightarrow 0, \text{ in } L^2(Q) \text{ strongly.}$$

Therefore there exist subsequences $y_-^n, y_+^n, y_\lambda^n$ converging to some y in $L^2(I, V)$ weakly, hence in $L^2(Q)$ weakly, and subsequences $z_-^n, z_+^n, z_\lambda^n$ converging to some z in $L^2(Q)$ weakly. By the Aubin compactness theorem (Temam, 1977, p. 271), there exists a subsequence y_λ^n converging to the same y in $L^2(Q)$ strongly, hence y_-^n, y_+^n converge to y in $L^2(Q)$ strongly, for the corresponding subsequences.

In order to pass to the limit in the discrete equations, let $v \in C_c^\infty(\Omega)$, $\xi \in C^1(\bar{I})$, with $\xi(T) = 0$, $w := v\xi$, v^n (resp. ξ^n) the continuous piecewise affine interpolant of v (resp. ξ), with respect to the partition of $\bar{\Omega}$ (resp. \bar{I}), and $w^n := v^n \xi^n$. The discrete state equation (19), with $v = w_{j+1}^n$, yields by summation

$$\int_0^T ((z_\lambda^n)', w_+^n) dt + \int_0^T a(t_+^n, y_+^n, w_+^n) dt = \int_0^T (f(t, y_-^n, r^n), w_+^n) dt,$$

hence, by the discrete integration by parts formula (31)

$$\begin{aligned} & - \int_0^T (z_-^n, (w_\lambda^n)') dt + \int_0^T a(t_+^n, y_+^n, w_+^n) dt = \\ & \int_0^T (f(t, y_-^n, r^n), w_+^n) dt + (z_0^n, v^n) \xi(0). \end{aligned}$$

Moreover, since $(y_\lambda^n)' = z_+^n$, then, by integrating by parts, we get

$$- \int_0^T (y_\lambda^n, v) \xi' dt = \int_0^T (z_+^n, v) \xi dt + (y_0^n, v) \xi(0).$$

Now, we have clearly

$$\begin{aligned} & z_-^n, z_+^n \rightarrow z \text{ in } L^2(Q) \text{ weakly,} \\ & (w_\lambda^n)' \rightarrow w' \text{ in } L^\infty(Q) \text{ strongly,} \\ & t_+^n \rightarrow t \text{ in } L^\infty(I) \text{ strongly,} \\ & y_+^n \rightarrow y \text{ in } L^2(I, V) \text{ weakly,} \\ & w_+^n \rightarrow w \text{ in } L^\infty(I, V) \text{ strongly,} \\ & y_-^n, y_+^n, y_\lambda^n \rightarrow y \text{ in } L^2(Q) \text{ strongly,} \\ & r^n \rightarrow r \text{ in } R, \\ & w_+^n \rightarrow w \text{ in } L^2(Q) \text{ strongly,} \\ & z_0^n \rightarrow y^1 \text{ in } L^2(\Omega) \text{ strongly,} \\ & y_0^n \rightarrow y^0 \text{ in } V \text{ strongly,} \\ & v^n \rightarrow v \text{ in } C(\bar{\Omega}) \text{ strongly.} \end{aligned}$$

We can thus pass to the limit in the above two equations, using Proposition 2.1, from Chrysosoverghi (1986), for the term containing f , to obtain

$$-\int_0^T (z, v)\xi' dt + \int_0^T a(t, y, v)\xi dt = \int_0^T (f(t, y, r), v)\xi dt + (y^1, v)\xi(0),$$

and

$$-\int_0^T (y, v)\xi' dt = \int_0^T (z, v)\xi dt + (y^0, v)\xi(0),$$

which hold for every $v \in V$, since $C_c^\infty(\Omega)$ is dense in V . Choosing first $\xi \in C_c^\infty(I)$, these equations yield $y' = z$ and equation (5) (see Temam, 1977, p. 250). Finally, integrating by parts, we obtain the initial conditions (6), (7). Therefore, $y = y_r$, and the convergences of the lemma hold for the original sequences. ■

LEMMA 4.4 *If $r^n \rightarrow r$ in R , then $J_m^n(r^n) \rightarrow J_m(r)$, $0 \leq m \leq q$, as $n \rightarrow \infty$.*

Proof. We have

$$J_m^n(r^n) = \int_Q g_m(x, t, y_-^n, r^n) dx dt,$$

and the lemma follows from Lemma 4.3. and Proposition 2.1 from Chrysosoverghi (1986). ■

First we study the behaviour in the limit of optimal discrete controls. We assume that the CRP is feasible. Consider the DRP^n (resp. DCP^n). In the presence of state constraints, we assume that the sequences $\{\varepsilon_m^n\}$, in (24) and (26), converge to zero as $n \rightarrow \infty$ and satisfy

$$\begin{aligned} |J_m^n(\bar{r}^n)| &\leq \varepsilon_m^n, \quad 1 \leq m \leq p, \\ J_m^n(\bar{r}^n) &\leq \varepsilon_m^n, \quad \varepsilon_m^n \geq 0, \quad p < m \leq q, \end{aligned}$$

for every n , where $\{\bar{r}^n \in R^n\}$ (resp. $\{\bar{r}^n \in W^n\}$) is a sequence converging to some optimal control \bar{r} of the CRP. Such sequences exist since, by Lemma 4.1, there exists a sequence $\{\bar{r}^n \in W^n\}$ such that $\bar{r}^n \rightarrow \bar{r}$, for some optimal \bar{r} , and by Lemma 4.4

$$\begin{aligned} \lim_{n \rightarrow \infty} J_m^n(\bar{r}^n) &= J_m(\bar{r}) = 0, \quad 1 \leq m \leq p, \\ \lim_{n \rightarrow \infty} J_m^n(\bar{r}^n) &= J_m(\bar{r}) \leq 0, \quad p < m \leq q. \end{aligned}$$

In particular, the DRP^n (resp. DCP^n) is thus feasible for every n . We have the following, rather theoretical, result.

THEOREM 4.1 *For each n , let r^n be optimal for the DRP^n (resp. DCP^n). Then the sequence $\{r^n\}$ has accumulation points, and every such point is optimal for the CRP.*

Proof. Since R is compact, let $\{r^n\}$ (same notation) be a subsequence converging to some $r \in R$. Since r^n is optimal and \bar{r}^n admissible for the DRP^n (resp. DCP^n), we have

$$J_0^n(r^n) \leq J_0^n(\bar{r}^n),$$

$$|J_m^n(r^n)| \leq \varepsilon_m^n, \quad 1 \leq m \leq p,$$

$$J_m^n(r^n) \leq \varepsilon_m^n, \quad p < m \leq q,$$

Taking the limit and using Lemma 4.4, we see that r is optimal for the CRP. If there are no state constraints, by using any sequence $\{\bar{r}^n \in W^n\}$ converging to some optimal control of the CRP, we also obtain in the limit that r is optimal for the CRP. ■

In the presence of inequality state constraints only, under appropriate assumptions, we can take $\varepsilon_m^n = 0$ in the discrete state constraints, following the approach of Casas (1996). Given $\delta \in \mathbf{R}$, define the parametrized continuous problems CRP_δ and CCP_δ , with state constraints $J_m \leq \delta$, $1 \leq m \leq q$, and discrete problems DRP_δ^n and DCP_δ^n , with constraints $J_m^n \leq \delta$, $1 \leq m \leq q$, and suppose that

(i) $\bar{\delta} := \inf\{\delta \mid \text{the } CRP_\delta \text{ is feasible}\} < 0$,

(ii) the CRP_0 is **stable to the left**, i.e. $\lim_{\delta \rightarrow 0^-} (\min CRP_\delta) = \min CRP_0$.

Note that since W is dense in R , we have

$$\bar{\delta} = \min_{r \in R} \left\{ \max_{1 \leq m \leq q} J_m(r) \right\} = \inf_{w \in W} \left\{ \max_{1 \leq m \leq q} J_m(w) \right\}.$$

Condition (i) implies that the CRP_δ is feasible for every $\delta \geq \bar{\delta}$ and not feasible for every $\delta < \bar{\delta}$. It follows from condition (i) and Lemmas 4.1 and 4.4 that the DRP_0^n and the DCP_0^n are feasible for n sufficiently large. If $\bar{\delta} < 0$, condition (ii) implies that the CRP_0 is **stable**, since it is always stable to the right, if $\bar{\delta} \leq 0$.

For $\bar{\delta} \leq \delta < 0$, it can be shown that

$$\min CRP_0 \leq \inf CCP_0 \leq \min CRP_\delta \leq \inf CCP_\delta.$$

Now suppose that $\bar{\delta} < 0$ and that the CCP_0 is **stable**, i.e.

$$\lim_{\delta \rightarrow 0} (\inf CCP_\delta) = \inf CCP_0.$$

Then $\min CRP_0 = \inf CCP_0$ (Casas, 1996), hence condition (ii) holds. Note that conditions (i) and (ii) also imply that

$$\min CCP_0 = \inf CCP_0.$$

Since the function $\theta \rightarrow \min \text{CRP}_\theta$, defined on $[\bar{\delta}, +\infty)$, is non increasing, it is continuous (i.e. the CRP_θ is stable) for every $\theta \geq \bar{\delta}$, except at most a countable number of them. This assures us in some manner that the CRP_0 is almost always stable.

Under conditions (i) and (ii), Theorem 4.1, with $\varepsilon_m^n = 0$, remains valid. To see this, let $\{r^{n_k}\}$ be a subsequence of $\{r^n\}$ such that $r^{n_k} \rightarrow r \in R$, where r is feasible for the CRP_0 by Lemma 4.4. Now let $\bar{\delta} < \delta < 0$ be given, and let r_δ be optimal for the CRP_δ . There exists a sequence $\{r_\delta^n \in W^n\}$ (Lemma 4.1) such that $r_\delta^n \rightarrow r_\delta$, where r_δ^n is feasible for the DRP_0^n (resp. DCP_0^n), for $n \geq N_0$ (Lemma 4.4). We have

$$J_0^{n_k}(r^{n_k}) \leq J_0^{n_k}(r_\delta^{n_k}), \quad \text{for } n_k \geq N_0,$$

and taking the limit, as $k \rightarrow \infty$

$$J_0(r) \leq J_0(r_\delta).$$

Finally,

$$J_0(r) \leq \lim_{\delta \rightarrow 0^-} J_0(r_\delta) = \lim_{\delta \rightarrow 0^-} (\min \text{CRP}_\delta) = \min \text{CRP}_0,$$

which shows that r is optimal for the $\text{CRP}_0 = \text{CRP}$.

Next, we examine the behaviour in the limit of extremal discrete controls. This is motivated by the fact that numerical optimization methods often compute approximations of extremal controls. We shall need the following lemma whose proof is similar to that of Lemma 4.3.

LEMMA 4.5 *If $r^n \rightarrow r$ in R , then the corresponding discrete adjoint states $\phi_-^n, \phi_+^n, \phi_\lambda^n$ converge to ϕ_r in $L^2(Q)$ strongly, as $n \rightarrow \infty$.*

Now consider the DRP'^n . Sequences $\{\varepsilon_m^n\}$, in (25) and (26), converging to zero and such that the DRP'^n is feasible for every n can be constructed here as follows. Let \bar{r}^n be any solution of the unconstrained problem

$$c^n := \min_{r^n \in R^n} \left\{ \sum_{m=1}^p [J_m^n(r^n)]^2 + \sum_{m=p+1}^q [\max(0, J_m^n(r^n))]^2 \right\},$$

and set

$$\bar{\varepsilon}_m^n = J_m^n(\bar{r}^n), \quad 1 \leq m \leq p,$$

$$\bar{\varepsilon}_m^n = \max(0, J_m^n(\bar{r}^n)), \quad p < m \leq q.$$

Let \tilde{r} be an admissible control of the CRP and $\{\tilde{r}^n\}$ a sequence converging to \tilde{r} . We have

$$\lim_{n \rightarrow \infty} J_m^n(\tilde{r}^n) = J_m(\tilde{r}) = 0, \quad 1 \leq m \leq p,$$

$$\lim_{n \rightarrow \infty} J_m^n(\tilde{r}^n) = J_m(\tilde{r}) \leq 0, \quad p < m \leq q,$$

which show that $c^n \rightarrow 0$, hence $\bar{\varepsilon}_m^n \rightarrow 0$, $1 \leq m \leq q$. Now choose $\varepsilon_m^n = \bar{\varepsilon}_m^n$, $1 \leq m \leq p$, and ε_m^n such that $\varepsilon_m^n \geq \bar{\varepsilon}_m^n$ and $\varepsilon_m^n \rightarrow 0$, $p < m \leq q$. Then clearly the DRP^n is feasible for every n . We suppose that the ε_m^n , $1 \leq m \leq q$, are chosen as above.

THEOREM 4.2 *For each n , let r^n be admissible and extremal for the DRP^n . Then the sequence $\{r^n\}$ has accumulation points, and every such point is admissible and extremal for the CRP.*

Proof. Since R is compact and $\sum_m |\lambda_m^n| = 1$, we can suppose that $r^n \rightarrow r \in R$ and $\lambda_m^n \rightarrow \lambda_m$, as $n \rightarrow \infty$, with $\sum_m |\lambda_m| = 1$, $\lambda_0 \geq 0$, $\lambda_m \geq 0$, $p < m \leq q$. By Lemmas 4.1, 4.3, 4.5 and Proposition 2.1, in Chrysoverghi (1986), we have, for given $\tilde{r} \in R$ and $\tilde{r}^n \rightarrow \tilde{r}$

$$\sum_{m=0}^q \lambda_m D J_m(r, \tilde{r} - r) = \lim_{n \rightarrow \infty} \sum_{m=0}^q \lambda_m^n D J_m^n(r^n, \tilde{r}^n - r^n) \geq 0,$$

$$\lambda_m J_m(r) = \lim_{n \rightarrow \infty} \lambda_m^n [J_m^n(r^n) - \varepsilon_m^n] = 0, \quad p < m \leq q,$$

$$J_m(r) = \lim_{n \rightarrow \infty} [J_m^n(r^n) - \varepsilon_m^n] = 0, \quad 1 \leq m \leq p,$$

$$J_m(r) = \lim_{n \rightarrow \infty} [J_m^n(r^n) - \varepsilon_m^n] \leq 0, \quad p < m \leq q,$$

which show that r is extremal and admissible for the CRP. ■

5. Final comments

In order to obtain in the limit the strong relaxed minimum principle, we have used the relaxed discrete problems, since (i) the strong discrete classical minimum principle does not hold in general (see Canon et al., 1970), and (ii) weak discrete classical necessary conditions obviously cannot yield the strong conditions in the limit.

Optimization methods, when applied to the discrete relaxed problem, usually compute discrete Gamkrelidze controls. These controls, which are convex combinations of Dirac relaxed controls, can then easily be approximated by piecewise constant classical controls by a simple procedure (see Warga, 1972, and Chrysoverghi et al., 1997).

Finally, the authors wish to thank the referees for their many useful remarks.

References

- BITTNER, L. (1975) On optimal control of processes governed by abstract functional, integral, and hyperbolic differential equations. *Math. Operationsforschung u. Statistik*, **6**, 107–134.
- CANON, M.D., CULLUM, C.D., and POLAK, E. (1970) *Theory of Optimal Control and Mathematical Programming*, McGraw-Hill, New York.
- CASAS, E. (1996) The Relaxation Theory Applied to Optimal Control Problems of Semilinear Elliptic Equations. *Proceedings of the 17th Conference on System Modelling and Optimization*, Prague, July 1995, Chapman and Hall.
- CHRYSOVERGHI, I. (1985) Numerical approximation of nonconvex optimal control problems defined by parabolic equations. *Journal of Optimization Theory and Applications*, **45**, 73–88.
- CHRYSOVERGHI, I. (1986) Nonconvex optimal control of nonlinear monotone parabolic systems. *Systems and Control Letters*, **8**, 55–62.
- CHRYSOVERGHI, I. and BACOPOULOS, A. (1993) Approximation of relaxed nonlinear parabolic optimal control problems. *Journal of Optimization Theory and Applications*, **77**, 31–50.
- CHRYSOVERGHI, I., BACOPOULOS, A., KOKKINIS, B., and COLETOS, J. (1997) Mixed Frank-Wolfe penalty method with applications to nonconvex optimal control problems. *Journal of Optimization Theory and Applications*, **94**, 2, 311–334.
- CULLUM, J. (1971) An explicit procedure for discretizing continuous optimal control problems. *Journal of Optimization Theory and Applications*, **8**, 14–34.
- EKELAND, I. (1972) Sur le contrôle optimal de systèmes gouvernés par des équations elliptiques. *Journal of Functional Analysis*, **9**, 1–62.
- FATTORINI, H.O. (1994) Existence theory and the maximum principle for relaxed infinite-dimensional optimal control problems. *SIAM Journal on Control and Optimization*, **32**, 311–331.
- FATTORINI, H.O. (1997) *Infinite Dimensional Optimization and Control Theory*, Cambridge Univ. Press, Cambridge.
- LEES, M. (1960) A priori estimates for the solutions of difference approximations to parabolic differential equations. *Duke Mathematical Journal*, **27**, 287–311.
- LIONS, J.L. (1972) *Optimal Control of Systems Governed by Partial Differential Equations*, Springer, New York.
- ROUBIČEK, T. (1991) A convergent computational method for constrained optimal relaxed control problems. *Journal of Optimization Theory and Applications*, **69**, 589–603.
- ROUBIČEK, T. (1997) *Relaxation in Optimization Theory and Variational Calculus*, W. de Gruyter, Berlin.
- SADEK, I.S., SLOSS, J.M., BRUCH JR., J.C., ADALI, S. (1996) Maximum

- principle for optimal control of a hyperbolic equation in two space dimensions. *J. of Vibration and Control*, **2**, 3–15.
- SLOSS, J.M., SADEK, I.S., BRUCH JR., J.C., ADALI, S. (1995) Maximum principle for optimal control of a hyperbolic equation in one space dimension, Part I: theory. *Journal of Optimization Theory and Applications*, **87**, 33–45.
- TEMAM, R. (1977) *Navier–Stokes Equations. Theory and Numerical Analysis*, North–Holland, New York.
- TEO, K.L., CLEMENTS, D.J., WU, Z.S. and CHOO, K.G. (1984) Convergence of a strong variational algorithm for relaxed controls involving a class of hyperbolic systems. *Journal of Optimization Theory and Applications*, **42**, 467–485.
- WARGA, J. (1972) *Optimal Control of Differential and Functional Equations*, Academic Press, New York.
- WARGA, J. (1977) Steepest descent with relaxed controls. *SIAM Journal on Control and Optimization*, **15**, 674–682.