

Acritarchs from Cambrian deposits of the southern part of the Łysogóry unit in the Holy Cross Mountains, Poland

Zbigniew SZCZEPANIK



Szczepanik Z. (2001) — Acritarchs from Cambrian deposits of the southern part of the Łysogóry unit in the Holy Cross Mountains, Poland. *Geol. Quart.*, 45 (2): 117–130. Warszawa

The palynology of the Cambrian claystones and mudstones of the Góry Pieprzowe Shale Formation in the southern part of the Łysogóry region (Holy Cross Mts., Poland) are described. These deposits had been referred to the Middle Cambrian of the *Paradoxides paradoxissimus* and *Paradoxides forchhammeri* Superzones. Here, rocks from 10 shallow drillings and several exposures in the Opatów region have been examined. Scarce and poorly preserved acritarch assemblages contain several forms that are indicative for the Upper Cambrian, while the genera *Acanthodiacrodium*, *Cymatiogalea*, *Stelliferidium*, *Trunculumarium*, *Nellia*, *Impliviculus*, *Calysiella*, and *Veryhachium* excludes the possibility of a Middle Cambrian age. Rather, these forms suggest the Upper Cambrian, most probably its middle and upper part. The occurrence of the rocks of this age to both north and south of the Łysogóry quartzite outcrops indicates tectonic repetition of the geological structure in the Main Range of the Holy Cross Mountains.

The dark colours of palynoflora from the Pieprzowe Mountains in Sandomierz are identical with the colours of the Cambrian microflora from the Łysogóry region, and differ from the bright colours characterising organic matter in the Kielce region of the Holy Cross Mountains. This suggests a tectonic relation of the Pieprzowe Mountains with the Łysogóry region rather than with the Kielce region, as hitherto thought.

Zbigniew Szczepanik, Holy Cross Mts. Branch, Polish Geological Institute, Zgoda 21, PL-25-953 Kielce, Poland (received: August 26, 1999; accepted: November 3, 2000).

Key words: Holy Cross Mts., Łysogóry unit, Cambrian, acritarchs.

INTRODUCTION

The Cambrian rocks that are widely distributed in the Holy Cross Mountains have prompted much discussion (Stupnicka, 1988; Po aryski and Tomczyk, 1993; Kowalczewski, 1995; Orłowski and Mizerski, 1995; Kowalczewski and Dadlez, 1996; Znosko, 1996), mainly concerning their stratigraphy. Trilobites, although long noted, come from too few localities to resolve the stratigraphy or constrain tectonic and palaeogeographic models.

The most neglected Cambrian rock formation in the Holy Cross Mountains is the Góry Pieprzowe Shale Formation (Orłowski, 1975). Middle Cambrian trilobites have been found, in the Kamie Łukawski quarry and on P czek hill (Gürich, 1892, Orłowski, 1964, 1985), but only from part of the succession, which achieves substantial thicknesses. The rocks along the southern slopes of the Main Range (Fig. 1), included in this formation, had not yielded fossils, while lithological correlation

was uncertain. These rocks were most commonly ascribed to the Middle Cambrian — to the *P. paradoxissimus* and *P. forchhammeri* Superzones (Samsonowicz 1934; Tomczykowa, 1968; Orłowski, 1975, 1988), although the Lower Cambrian has also been suggested (Czarnocki, 1947, 1957; ak, 1962; Kowalczewski *et al.*, 1976; Kowalczewski, 1995).

Since the 1960's, acritarchs have become increasingly important in dating early Palaeozoic rocks. Acritarchs from the southern part of the Holy Cross region and from the Nida Basin (Vidal in: Po aryski *et al.*, 1981; Lenzion *et al.*, 1983; Moczyłowska in: Kowalczewski *et al.*, 1984, 1987) provided a detailed Cambrian biostratigraphy, which in turn constrained the geological structure. In this paper, acritarchs are used to date the mudstones and claystones of the Łysogóry region, which up to now were devoid of palaeontological documentation. These studies have been encouraged by preliminary work (Moczyłowska in: Kowalczewski *et al.*, 1987), which outlined the problems of age and structure in the Main Range of the Holy Cross Mountains.




Fig. 1. Location of the boreholes drilled, and Cambrian outcrops in the Holy Cross Mountains

1— Czarna Shales, 2— Ocieski Sandstones and Kamieniec Shales, 3— Słowic Sandstones, 4— Uszów Sandstones, 5— Góry Pieprzowe Shales (part of the Góry Pieprzowe Shale Formation), 6— Ublinek Beds, 7— Krajno Beds (part of the Góry Pieprzowe Shale Formation), 8— Wworków Sandstones, Łysogóry Quartzites (Wiśniówka Quarzitic Sandstone Formation), 9— Mchojce Beds, Łysogóry Beds (Klonówka Shale Formation), 10— deposits younger than Cambrian, 11— boreholes, in which acritarchs were found, 12— boreholes, in which acritarchs were not found, 13— faults, 14 Holy Cross Fault

AIM OF STUDIES

Early attempts at finding acritarchs in the Góry Pieprzowe Shale Formation from field exposures were not successful. Therefore, 10 shallow (up to 40 m) boreholes were drilled into unweathered rock, in order to recover stratigraphically useful microflora.

This sampling took place in the southern part of the Łysogóry region, adjacent to the Main Holy Cross Fault and in the Pieprzowe Mountains (Fig. 1), which as regards overall structure are included in the Kielce region of the Holy Cross Mountains.

In the west part of the area sampled, the drilling took place along the southern slopes of the Main Range, in an area limited to the south by the Main Holy Cross Fault, and to the north by the Wiśniówka Quarzitic Sandstone Formation (Orłowski, 1975). Four shallow boreholes: Kielce-Piaski PIG 1, Dąbrowa-Koszarka PIG 1, Huta Nowa PIG 1, Piórków PIG 1 (Fig. 1) were drilled here.

In the Opatów region, where most of the surface sampling was concentrated, the topography is lower and outcrop pattern is not so distinct. Acritarchs from the Kochówka, Kania and Opatówka valleys, and from the Kochówek PIG 1, Kobylanki

PIG 1 and Doły Oziębowskie PIG 1 boreholes were studied (Fig. 2).

Around Sandomierz the rocks from three boreholes were studied: Kamień Plebański PIG 1, Kamień Plebański PIG 2 and Kamień Łukawski PIG 1 (Fig. 1).

LITHOLOGY AND TECTONICS OF THE ROCKS STUDIED

The borehole material obtained was generally poor, with cores often disintegrated. The cores are dominantly of dark grey and black claystones (Fig. 3), locally accompanied by mudstones of similar colour. Scarce, thin (up to 1 cm) sandstone interlayers occur.

All rocks studied are strongly tectonized (Fig. 3), with both brittle and plastic deformation. Where bedding is visible, dips are varied, and most commonly steep. Brecciation zones are common, as fine, randomly arranged claystone clasts bound by fine-grained claystone detritus. These are generally difficult to recognise, because of fracturing during drilling, and they were obtained only in the form of loose fragments. In some drillings




Fig. 2. Geological map of the Opatów area after Z. Kowalczewski (Kowalczewski *et al.*, 1976); age interpretation changed

1 — claystone package (Upper Cambrian, most probably the middle part of the Upper Cambrian); 2 — mudstone-claystone succession (?Cambrian, age undetermined); 3 — sandstone-mudstone succession (? Cambrian, age undetermined); 4 — mudstone-claystone succession (? Cambrian, age undetermined); 5 — sandstone-quartzite-mudstone succession (Upper Cambrian, most probably lower part of the Upper Cambrian); 6 — Ordovician and Silurian; 7 — Lower Devonian; 8 — Middle Devonian; 9 — Upper Devonian; 10 — Permo-Mesozoic; 11 — Main Holy Cross Fault; 12 — faults: a — certain, b — probable; 13 — boreholes studied; 14 — localities of positive samples; 15 — localities of negative samples

(Kamie Pleba ski PIG 2, Piórków PIG 1) the fracturing was so extensive, that no material was obtained from intervals of several metres. Slickensides on bedding and fracture surfaces are common, while fracture surfaces are often inclined at about 30° bedding. Fracture patterns are often picked out by white minerals of the alum group, which also crystallized as veins and in spaces generated tectonically. Minor folds were rarely observed.

All boreholes are similarly affected, suggesting that the whole southern part of the Łysogóry region and Pieprzowe Mountains were subject to intensive tectonic deformation.

METHODOLOGY

200–400 g samples were macerated by acid dissolution and flotation. Glycerine-gelatine preparations were made of all macerates, and were studied in transmitted light. Pictures of the specimens were recorded as graphic computer files on to a database, and used as illustration in this paper.

Individual samples vary in acritarch frequency. The least common assemblages were found in the Pieprzowe Mountains, where only *ca.* 10% of the samples contain scarce palynomorphs. Acritarchs were most frequent in the

Kochówek-Ozi błów region. Along the southern slopes of the Main Range, acritarchs were found in only two out of four boreholes. In the Kielce-Piaski PIG 1 borehole, three samples yielded scarce but stratigraphically useful assemblages of poorly preserved palynomorphs. In the Piórków PIG 1 borehole an assemblage yielded numerous microspores determined by E. Turnau (*pers. inf.*) as mixed Devonian and Carboniferous, together with individual acritarchs of Lower Cambrian to lower Middle Cambrian age. The palynomorphs are relatively well preserved and their colours are bright.

The acritarchs are mostly thermally altered, with colours ranging from dark brown to black. No fractures or traces of abrasion that could indicate redeposition were found. The poor preservation state mean that most determinations are at generic level.

RESULTS OF PALYNOLOGICAL STUDIES

PIEPRZOWE MOUNTAINS

Three boreholes were drilled in the Pieprzowe Mountains. Two (Kamie Pleba ski PIG 1 and Kamie Pleba ski PIG 2) were drilled along the western limit of the Pieprzowe Moun-




Fig. 3. Profiles of selected boreholes

1 — Quaternary, 2 — Carboniferous, 3 — Devonian, 4 — Cambrian, 5 — claystones, 6 — mudstones, 7 — sandstones, 8 — breccias, 9 — loess sands, 10 — sands, 11 — samples, in which no microflora has been found, 12 — samples, in which microflora has been found

tains escarpment (Fig. 1), one where the flood dam reaches the escarpment and one in the Kamie Łukawski village (Fig. 1). These investigated the rocks on either side of the old sandstone quarry which had yielded a Middle Cambrian trilobite fauna.

All boreholes made in the Sandomierz region reached the monotonous black claystones of the Góry Pieprzowe Shale Formation (Fig. 3). These contain little or no coarser material or sedimentary structures, perhaps exacerbated by the steep bedding dip (70–90°) and therefore a small true thickness of the succession studied.

Only one of these boreholes, on the western side of the quarry (Kamie Pleba ski PIG 1), yielded acritarchs, which were scarce and badly preserved (Table 1). Most common here are *Eliasum* sp. and *Cristallinium cambriense* (Slavikova), with poorly preserved forms referable to *Micrhystridium*, including *M. lanceolatum* Vanguetaine. Most of these forms show long stratigraphic ranges, from the Middle Cambrian to the Tremadoc, or even younger, though this type of assemblage is most commonly found in the Middle Cambrian. However, the acritarchs include Upper Cambrian forms: *Calyxiella* cf. *izohoriensis* Golub et Volkova (Pl. I, Fig. 26) and *Impluviculus*. Both genera occur no earlier than the middle or upper (*Calyxiella*) part of the Upper Cambrian.

Two stratigraphic interpretations are possible:

— an atypical Upper Cambrian assemblage, dominated by long-ranging forms with only a minor contribution of typical Upper Cambrian acritarchs; in such a case, the uppermost Cambrian may be excluded, because characteristic “galeate” and Diacriodidae forms were already very common at that time;

— tectonic mixing of the claystones in the Kamie Pleba ski PIG 1 borehole; these are strongly crumpled and slickensided and contain common brecciation levels (Fig. 3); Middle Cambrian claystones could have been tectonically interleaved with Upper Cambrian deposits.

The first of these hypotheses seems more probable in explaining the presence of typical Upper Cambrian genera.

OPATÓW AREA

Bulk of the Cambrian claystone outcrops of the Opatów area, indicated on the geological maps and mentioned in the older papers, do not exist any more. In this context, the samples from the three shallow boreholes (Fig. 2) are very useful. 12 samples were studied, and acritarchs were found in 11 of them (Fig. 3, Table 1).

Several hundred acritarchs were recovered. Their poor preservation and low frequency limit assignment to the Upper Cambrian (Fig. 4), and reference to the precise biostratigraphy proposed for the East European Craton (Volkova, 1990) (Fig. 4) is not possible. In general, though, the acritarch assemblages are similar to those described from the former USSR (Volkova and Golub, 1985; Volkova, 1990).

Generally positive results have been obtained from 4 regions (Fig. 2, Table 1). These are:

— Ozi błów-Kochówek region, in the extreme south of the area studied,

- exposures near the Jurkowiec-Pipała farmstead,
- exposures in the valley escarpment at Marcinkowice,
- Karwów-Krowieniec region.

KOCHÓWEK-OZI BŁÓW REGION

The area is located along the Main Holy Cross Fault zone, between the villages of Bratków, Kobylanki and Kochówek (Fig. 2). Rocks from three boreholes were studied: Kobylanki PIG 1, Kochówek PIG 1 and Doły Ozi błowskie PIG 1 (Fig. 3), and also from exposures in Kobylanki and Kochówek. Positive results were obtained only from the borehole cores (Table 1), which were mainly of claystones with thin sandy interlayers and laminae. Sedimentological analysis was restricted by the poor core preservation, with material obtained only as clasts. This probably results both from deep weathering and from strong tectonic fracturing.

The poorest assemblage was recognised in the Kochówek PIG 1 borehole (Table 1). The acritarchs here have a wide stratigraphic range of Middle and Upper Cambrian to Lower Tremadoc. The presence of *Timofeovia* suggests a level not older than the Middle Cambrian *P. paradoxissimus* Superzone (Volkova, 1990; Lenzion and Jankauskas, 1992).

A similar, but more abundant assemblage was found in the Doły Ozi błowskie PIG 1 borehole (Table 1), mostly comprising forms ranging from the Middle to the Upper Cambrian: *Cristallinium cambriense* (Slavikova) (Pl. III, Fig. 2), *Micrhystridium* div sp. with scarcer *Timofeovia phosphoritica* Vanguetaine, *Dictyotidium* sp., *Eliasum llaniscum* Fombella, *Comasphaeridium* sp., *Polygonium* div. sp., and *Pterospermella* sp. These long-ranging (Lower/Middle Cambrian to Tremadoc; Welsch, 1986; Martin and Dean, 1988; Volkova, 1990; Moczydłowska, 1995) forms are accompanied by *Izohoria* sp., *Calyxiella* sp. (Pl. I, Fig. 27) and poorly preserved forms most probably representing *Cymatiogalea* sp. and *Stelliferidium* sp. (Pl. I, Figs. 18, 22). These are not known from deposits younger than Upper Cambrian.

The most numerous and best preserved assemblage was found in the Kobylanki PIG 1 borehole (Table 1): *Cymatiogalea* sp. (Pl. I, Fig. 17), *Stelliferidium* sp. (Pl. I, Fig. 23), *Impluviculus* sp. (Pl. II, Figs. 17, 18), *Veryhachium* sp. (Pl. II, Fig. 13), *Adara* cf. *alea* Martin (Pl. III, Fig. 13), *Eliasum llaniscum* Fombella (Pl. II, Figs. 10, 11), *Lophosphaeridium* sp., *Polygonium* sp., *Timofeovia* sp. (Pl. III, Fig. 14) and others. This assemblage suggests a level not older than WK 2 (*Olenus*) of the Upper Cambrian and probably not younger than WK 5 (*Peltura*) of the Upper Cambrian (Fig. 4).

The Upper Cambrian acritarchs are accompanied by a single specimen of *Adara* cf. *alea* Martin, which has been thought to be of lower Middle Cambrian age (Volkova, 1990). Moczydłowska (1998) suggests that this form reaches the Upper Cambrian and this find supports this interpretation.

JURKOWICE-PIPAŁA REGION

This region includes the southern Kochówka valley, south of Jurkowiec, where the claystones and mudstones, similar to those of the Kochówek-Kobylanki region are exposed. Three of the samples yielded abundant acritarchs, including a dozen or so acritarchs of the distinctive group Diacromorphitae (Table 1, Figs. 6–8, 11, 12, 24, 25), which occur abundantly not earlier than in the upper part of the Upper Cambrian. In samples 115 and 116 a few specimens of *Trunculumarium revinium* Vanguetaine (Pl. I, Figs. 2–5) were also found, indicating the

middle Upper Cambrian, i.e. the *Trunculumarium revinium-Veryhachium dumonti* Superzone (Vanguetaine and Van Looy, 1983) or WK3 *T. revinium*, *D. caudatum* Zone (Volkova, 1990) (Fig. 4) corresponding to the (3–4) *Parabolina spinulosa-Leptoplastus* Zones of the standard trilobite zonation. This stratigraphic interpretation is consistent with other observations:

— the “galeate” forms of Herkomorphitae group with a broad umbo in polar position (Pl. I, Figs. 13, 14, 19) indicate an age not older than WK 2 of the Upper Cambrian of the East European Craton (Fig. 4), and therefore corresponding at least to the *Olenus* (2) Zone in the standard trilobite zonation;

— the presence of *Veryhachium* sp. (Pl. II, Fig. 14) and *?Vogtlandia* sp. (Pl. I, Fig. 29); these are Upper Cambrian taxa, which occur not earlier than in WK 2 of the Upper Cambrian (Fig. 4), i.e. in deposits not older than the *Olenus* (2) Zone.

MARCINKOWICE REGION

Here the Cambrian rocks are well exposed, but dominated by sandstones and quartzitic sandstones, while claystones and mudstones are rare. Scarce and poorly preserved microflora assemblages have been obtained. 7 samples were studied, and positive results were obtained only from 2. The assemblages are dominated by poorly preserved *Timofeevia*, *Cristallinium* and *Retisphaeridium*. Individual specimens referable to Upper Cambrian “galeate” forms and *Impluviculus*, have, however, also been found (Pl. I, Fig. 15; Pl. II, Figs. 15, 16).

KARWÓW-KROWIENIEC REGION

Four samples have been collected from this region, two of which contained a poor microflora (Table 1). The assemblage is dominated by compact forms with a surface divided into polygonal fields including: *Cristallinium* sp., *Retisphaeridium* sp., *Timofeevia* sp., *Multiplicisphaeridium* sp. and *Micrhystridium* sp. These indicate a Middle Cambrian-Tremadoc interval.

THE SOUTHERN SLOPES OF THE MAIN RANGE

The area is a 50 km long strip of land between Piórków near Łągów (southern slopes of the Jeleniowskie Range) and the northern suburbs of Kielce (southern slopes of Krzemianka Mountain). The deposits adjoining the Main Holy Cross Fault were studied where geoelectrical studies indicated low-resistivity clay deposits. The boreholes (Figs. 1 and 3) reached unweathered Cambrian deposits at relatively shallow depths with relatively good core preservation.

The deposits are represented by claystones with scarce sandy intercalations. Bedding dips are high 30–90°, and the rocks are strongly tectonized (Fig. 3). Two out of 4 boreholes yielded acritarchs: Piórków FIG 1 and Kielce-Piaski FIG 1, while two (Huta Nowa FIG 1 and D browa-Koszarka FIG 1) were barren.

The Piórków FIG 1 borehole (Fig. 1) was drilled in the zone of the Main Holy Cross Fault. The claystones penetrated are thoroughly shattered, and core recovery was poor. The samples from this tectonic breccia yielded no acritarchs (Table 1,

Fig. 3). Instead, numerous perfectly preserved, bright spores of Devonian and Carboniferous age were found (Turnau, 1996, pers. com.). In the samples from beneath the breccias abundant Devonian and Carboniferous spores occur (more than 99% of the material), together with individual acritarchs referable to *Leiosphaeridia*, *Lophosphaeridium* and *Skiagia?*, which suggest a Lower Cambrian-lower Middle Cambrian age. Their preservation, relative to those in other boreholes in the Łysogóry region, is surprisingly good, with brightly coloured palynomorph walls. This may be due to tectonic interleaving. Alternatively, the brecciated claystones at the base may represent the Devonian and Carboniferous of the Kielce region.

The most interesting results came from the Kielce-Piaski FIG 1 borehole located in the extreme west of the area studied, on drift-covered outcrops of dark Cambrian claystones, on the southern slopes of the Krzemianka Mountain (Fig. 1). Here (Fig. 3), typically Upper Cambrian acritarchs were found: *?Stelliferidium* sp. (Pl. I, Fig. 16), *Nellia magna* Volkova (Pl. I, Fig. 28), *Trunculumarium revinium* Vanguetaine (Pl. I, Fig. 1), *?Ladogiella* sp. (Pl. II, Fig. 19), *Impluviculus*, *Acanthodiacrodium* sp. (Pl. I, Figs. 9, 10). These forms indicate at least the 3 Zone of the Upper Cambrian (Volkova, 1990) (Fig. 4).

In the Kielce-Piaski FIG 1 borehole, as at Opatów, the upper age limit is difficult to establish. The low proportion of forms with diacroidal symmetry, the numerous *Cristallinium*, and the relatively simple structure of the *Acanthodiacrodium* taxa do not suggest very high levels in the Upper Cambrian. This opinion is supported by the occurrence of a single poorly preserved specimen of *Trunculumarium*, a genus known only from the middle part of the Upper Cambrian.

STRATIGRAPHIC SIGNIFICANCE OF THE ASSEMBLAGES AND COMPARISON WITH OTHER AREAS


The acritarch assemblages recognised represent the Middle Cambrian to Upper Cambrian and Tremadoc interval. Despite more than 30 years of study of this interval, there is not yet a coherent zonation. Biostratigraphical work needs to make use of various regional zonations (Vanguetaine, 1974; Martin and Dean, 1981, 1988; Vanguetaine and Van Looy, 1983; Di Mila *et al.*, 1989; Volkova, 1990; Martin, 1992) (Fig. 4).

The stratigraphy used here is based mainly on the zones recognised by Volkova (1990) on the East European Craton (Fig. 4), with which the microflora recognised in this study show strong affinity. Use was made also of work in Newfoundland and Canada by F. Martin (Martin and Dean, 1988; Martin, 1992) (Fig. 4), where the stratigraphic scheme developed is correlated with the trilobite zonation. In Newfoundland, however, substantially fewer acritarch taxa were determined than on the East European Craton.


The oldest acritarch assemblage recognised was in the Kochówek FIG 1 borehole and in outcrops in the Karwów-Krowieniec region. The acritarch assemblage occurring here, with numerous *Cristallinium cambriense* (Slavikova), *Eliasum llaniscum* Fombella, *Micrhystridium*



1. *Trunculumarium revinium* Vanguetstaine, Kielce-Piaski FIG 1. 2–5. *Trunculumarium revinium* Vanguetstaine, Jurkowiec-Pipała. 6, 8, 11. *Dasydiacrodium* sp., Jurkowiec-Pipała. 7, 12. *Acanthodiacrodiium* sp., Jurkowiec-Pipała. 9, 10. *Acanthodiacrodiium* sp., Kielce-Piaski FIG 1. 13, 14. ?*Cymatiogalea* sp., Jurkowiec-Pipała. 15. *Acritarcha* — „galeate”, Marcinkowice. 16. ?*Stelliferidium* sp., Kielce Piaski FIG 1. 17, 21. *Cymatiogalea* sp., Kobylanki FIG 1. 18. *Stelliferidium* sp., Doły Ozi błowskie FIG 1. 19. *Cymatiogalea* sp., Jurkowiec-Pipała. 20. ?*Calyxiella* sp., Jurkowiec-Pipała. 22. *Cymatiogalea* sp., Doły Ozi błowskie FIG 1. 23. *Stelliferidium* sp., Kobylanki FIG 1. 24, 25. ?*Lophodiacrodiium* sp., Jurkowiec-Pipała. 26. *Calyxiella* cf. *izohoriensis* Golub et Volkova, Kamie Pleba ski FIG 1. 27. *Calyxiella* sp., Doły Ozi błowskie FIG 1. 28. *Nellia magna* Volkova, Kielce-Piaski FIG 1. 29. ?*Vogtlandia* sp., Jurkowiec-Pipała. 30. ?*Vulcanisphaera* sp., Jurkowiec-Pipała



1-4. *Polygonium* sp. 1, Jurkowice-Pipała. 5, 6. *Polygonium* sp. 2, Jurkowice-Pipała. 7. *Stellechinatium* sp., Kobylanki FIG 1. 8. *Leiofusa* sp., Jurkowice-Pipała. 9. *Eliasum* sp., Kobylanki FIG 1. 10, 11. *Eliasum llanicum* Fombella, Kobylanki FIG 1. 12. *Izohoria angulata* Golub et Volkova, Kobylanki FIG 1. 13. *Veryhachium* sp., Kobylanki FIG 1. 14. *Veryhachium* sp., Jurkowice-Pipała. 15, 16. *Impliviculus* sp., Marcinkowice. 17, 18. *Impliviculus* sp., Kobylanki FIG 1. 19. ?*Ladogiella* sp., Kielce-Piaski FIG 1. 20-22. *Aranidium* sp., Jurkowice-Pipała. 23, 24. *Comasphaeridium* sp., Jurkowice-Pipała



1. *Cristallinium cambiense* (Slavikova), Kobylanki FIG 1. 2. *Cristallinium cambiense* (Slavikova), Doły Ozi błowskie FIG 1. 3. *Cristallinium* sp., Kobylanki FIG 1. 4. *Cristallinium* sp., Jurkowice-Pipała. 5. *Cristallinium* sp., Marcinkowice. 6, 8. *Dictyotidium* sp., Kobylanki FIG 1. 7. *Retisphaeridium* sp., Doły Ozi błowskie FIG 1. 9. ?*Retisphaeridium* sp., Kobylanki FIG 1. 10. *Cymatiosphaera* sp., Kochówek FIG 1. 11. ?*Cymatiogalea* sp., Doły Ozi błowskie FIG 1. 12. *Retisphaeridium* sp., Kielce-Piaski FIG 1. 13. *Adara* cf. *alea* Martin, Kobylanki FIG 1. 14. *Timofeevia* sp., Kobylanki FIG 1. 15. *Timofeevia* sp., Kielce-Piaski FIG 1. 16. *Timofeevia* cf. *phosphoritica* Vanguetstaine, Karwów. 17. *Timofeevia phosphoritica* Vanguetstaine, Marcinkowice. 18. *Multiplicisphaeridium* sp., Jurkowice-Pipała. 19. *Multiplicisphaeridium* sp., Kobylanki FIG 1. 20. *Heliosphaeridium* sp., Jurkowice-Pipała. 21. *Asteridium* sp., Jurkowice-Pipała

div. sp., *Timofeevia* sp., *Retisphaeridium* sp. and others (Table 1) is difficult to interpret stratigraphically. It may represent a broad time span of the Middle Cambrian, or a depleted Upper Cambrian assemblage. Until recently *E. llaniscum* Fombella would suggest the Middle Cambrian. However, newer data (Moczyłowska, 1995, 1998) suggests that it is present also in Upper Cambrian deposits. On balance, a Middle Cambrian assemblage, not older than the *P. paradoxissimus* Superzone, is present in the Kochówek PIG 1 borehole. It resembles assemblages in SK 2 of the Middle Cambrian of the East European Craton (Volkova, 1990) and the lower part of the A2 microflora in Newfoundland (Martin and Dean, 1988). However, it most resembles Superzone *C. cambriense*, *Eliasum*, *Timofeevia* (Vanguetaine and Van Looy, 1983) (Fig. 4), correlated with the upper part of the *E. oelandicus* Superzone and lower part of the *P. forchhammeri* Superzone. Such assemblages are known from other areas, such as the Czech Republic (Vavrdová, 1982), Turkey (Erkmen and Bozdogan, 1981), Morocco (Vanguetaine and Van Looy, 1983), Belgium (Vanguetaine, 1974) and Spain (Fombella, 1977, 1979).

In the Kobylanki PIG 1, Doły Ozi błowskie PIG 1 and Kielce-Piaski PIG 1 boreholes, as well as in the Jurkowiec-Pipała outcrop, a younger assemblage has been found. This includes numerous *Cristallinium cambriense* (Slavikova) and *Polygonium* sp., and individual specimens of *P. minimum* (Timofeev), *P. martinae* (Moczyłowska), *Timofeevia lancarae* (Cramer et Diez), *Izohoria* sp., *Impluviculus* sp., *Cymatogalea* sp., *Stelliferidium* sp., *Eliasum* sp., *Trunculumarium* cf. *revinium* Vanguetaine, *Micrhystridium* sp. and *Acanthodiacrodium* sp. The genera *Stelliferidium*, *Cymatogalea*, *Impluviculus*, *Izohoria*, *Acanthodiacrodium*, *Trunculumarium* and *Nellia* are known exclusively from Upper Cambrian and younger deposits. The genera *Impluviculus*, *Izohoria*, *Acanthodiacrodium*, *Trunculumarium* and *Nellia* occur not earlier than the middle part of the Upper Cambrian, and therefore, most probably, the deposits drilled here represent the middle or the upper part of the Upper Cambrian. The assemblages found in the Kielce-Piaski PIG 1 borehole and in the Jurkowiec outcrop most compare with the WK 3–WK 4A Zones of the Upper Cambrian of the East European Craton (Fig. 4), while the Kobylanki PIG 1 and Doły Ozi błowskie PIG 1 boreholes and the outcrops at Marcinkowiec may represent the WK 2–WK 4B Zones of this division (Fig. 4).

The scarce Upper Cambrian forms found in the Kamie Pleba ski PIG 1 borehole are difficult to locate biostratigraphically. *Calyxiella* suggests the upper part of *Peltura* (5) Zone, which approximately corresponds with the WK 4B Zone of the Cambrian of the East European Craton (Volkova, 1990) (Fig. 4). Earlier works by this author (Volkova and Golub, 1985), though, noted this taxon also in the *Leptoplastus* Zone of the standard trilobite zonation. Outside of the former USSR this form has been recognised on Öland (Sweden) in the *Peltura* (5) Zone (Di Mila *et al.*, 1989) and also in of the Ublinek PIG 1 bis borehole (Szczepanik, 1997a), where it co-occurs with specimens suggesting the Cambrian/Tremadoc boundary.

The Upper Cambrian assemblages are quantitatively dominated by taxa with long stratigraphic ranges [e.g. *Cristallinium cambriense* (Slavikova) and *Micrhystridium* div. sp.]. *Polygonium* is also common, represented by small forms with relatively short outgrowths.

The most important difference vis-a-vis the platform is the longer stratigraphic range of *Eliasum llaniscum* Fombella, supporting the work of Moczyłowska (1995, 1998), and the smaller morphological diversity of *Cristallinium* and the earlier occurrence *Polygonium*.

No abundance peaks of *Timofeevia* were found, that could indicate the upper Middle Cambrian or lowermost Upper Cambrian. Such abundance peaks were found in claystones interbedded with quartzites in the Wi niówka quarries (Moczyłowska in: Kowalczewski *et al.*, 1987). Deposits of this age were probably not encountered in the boreholes.

Nowhere in the Holy Cross Mountains have assemblages with common *Raphesphaera* been found; such assemblages are characteristic of the lower part of the Upper Cambrian, and are common on the East European Craton and in Newfoundland, and also near to the platform zone (Narol PIG 2 borehole).

The assemblages recognised closely resemble to the microfloras from Newfoundland, Belgian and French Ardennes, Scandinavian Caledonides and other areas (Vanguetaine, 1974; Martin and Dean, 1981, 1988; Vanguetaine and Van Looy, 1983; Welsch, 1986; Di Mila *et al.*, 1989; Martin, 1992).

ACRITARCH COLOUR AS A PALAEOTEMPERATURE INDICATOR — TECTONIC IMPLICATIONS

Acritarchs found are dark brown to black, suggesting burial temperatures of the order of 100–200°C (AMOCO 5+–6). Acritarchs from Wi niówka (outside the study area) show even greater thermal alteration (AMOCO 6+–7) (Szczepanik, 1997b). These observations are consistent with those observed by M. Moczyłowska (in: Kowalczewski *et al.*, 1986), although showing a narrower range.

The acritarchs from the Pieprzowe Mountains thus reveal palaeotemperatures typical of the Łysogóry region (over 100°C), though their tectonic setting has in the past been linked with the Kielce region. The lithology, tectonic style and palaeothermal observations rather, however, suggest connection with the southern peripheries of the Łysogóry region. With the geology of the eastern part of the Holy Cross Mountains in mind, faults perpendicular to the Main Holy Cross Fault may be inferred along which the eastern part of the Łysogóry region has been moved southwards.

The extent to which rock heating may have been influenced by local brittle tectonics (near-fault heating) needs consideration. However, bright Devonian spores occur directly along the Holy Cross Fault (Piórków PIG 1 borehole). The factor responsible for thermal alteration was therefore a regional temperature increase caused by overburden thickness and heat production during orogeny.

CAMBRIAN		ORDOVICIAN		Volkova (1990)	Welsch (1986)	Martin and Dean (1981, 1988), Martin (1992)	Vanguetaine and Van Looy (1983)						
LOWER	MIDDLE	Paradoxides	forchhammeri	SK 2	A II	A. alea	C. cambrien. Eliasum Timofeevia						
				SK 1		R. terranov.							
				???	A I	AD-1	A. umbonul. S. compres.						
				Eccapara-doxides oleandicus									
		Protolenus	Lower Cambrian assemblages										
		UPPER	Acerocare	Peltura	Leptoplastus	Parabolina	Olenus	A. pisiformis	A. revinium V. dumonti				
										WK 2	A III	A 3	T. pentagon. V. turbata
										WK 1	A 2		
										WK 3	A 4	A 4	T. revinium V. dumonti
										???			
										WK 4A	A 5	A 5	T. revinium V. dumonti
										WK 4B			
WK 5	A IV	???	A. rommel. V. capillata										
TREMADOC	WK 5	AV	A 6	AU 3	A. destomb. V. capillata								
				C. vilco-xianum									




Fig. 4. Stratigraphic ranges of the assemblages found in particular outcrops and boreholes, together with the correlation of selected microfossil zones

CONCLUSIONS

Palynological studies in the southern part of the Łysogóry unit (boreholes: Kielce-Piaski PIG 1, Doły Ozi bławskie PIG 1, Kobylanki PIG 1 and exposures: Jurkowice-Pipała, Marcinkowice) document Upper Cambrian deposits in the southern part of the Łysogóry region, and this discovery complicates the tectonic picture of the Łysogóry region, and renders the hypothesis of a monoclinical structure in this area (Mizerski, 1992, 1994) improbable.

The acritarch assemblages of the study area resemble those from the East European Craton and Newfoundland, and also other areas, including the Peri-Gondwanan regions.

Acritarchs from the Pieprzowe Mountains show a similar preservation to those from the Łysogóry region. The degree of thermal maturity suggests heating to temperatures of 100°C and more, as in the Łysogóry region, but contrasting with that in the Cambrian rocks in the eastern part of the Kielce region, which is minimal.

The occurrence of a mixed Carboniferous-Devonian-Cambrian acritarch and spore assemblage from the Piórków PIG 1 borehole proves Variscan or post-Variscan activity along the Main Holy Cross Fault. The unaltered (bright) colours of this microflora suggests derivation of this assemblage from the Kielce region, in turn indicating an overthrust character (at least in the near-surface) for the Main Holy Cross Fault.

REFERENCES

- CZARNOCKI J. (1947) — Przewodnik XX Zjazdu Polskiego Towarzystwa Geologicznego w Górach wi tokrzyskich w 1947. Roczn. Pol. Tow. Geol., **17**: 237–295.
- CZARNOCKI J. (1957) — Tektonika Gór wi tokrzyskich, Geologia obszaru łysogórskiego. Pr. Inst. Geol., **2** (3): 11–138.
- DI MILA A., RIBECAL C. and TONGIORGI M. (1989) — Late Cambrian acritarchs from the *Peltura scarabaeoides* Trilobite Zone at Degerhamn (Öland, Sweden). Palaeontographia Italica, **76**.
- ERKMEN U. and BOZDOGAN N. (1981) — Cambrian acritarchs from the Sosnik formation in Southeast Turkey. Rev. Esp. Micropaleont., **13** (1).
- FOMBELLA M. A. (1977) — Acritarcos de edad Cambrio Medio-Inferior de la Provincia de Leon. Revista Espanola de Micropaleontologia, **9**: 115–124.
- FOMBELLA M. A. (1979) — Palinologia de Formacion Oville al Norte y Sur de la Cordillera Cantabrica, Espana Palinologia, **1**: 1–16.
- GÜRICH G. (1892) — Über eine cambrische Fauna Fauna von Sandomir in Russisch-Polen. Neues Jb. Miner., **1**. Stuttgart.
- KOWALCZEWSKI Z. (1995) — Fundamental stratigraphic problem of the Cambrian in the Holy Cross Mts. Geol. Quart., **39** (4): 449–470.
- KOWALCZEWSKI Z. and DADLEZ R. (1996) — Tectonics of the Cambrian in the Wi niówka area (Holy Cross Mts., Central Poland). Geol. Quart., **40** (1): 23–46.
- KOWALCZEWSKI Z., KULETA M., LISIK R. and MOCZYDŁOWSKA M. (1986) — New data on Cambrian and Lower Ordovician rocks in the vicinities of Wi niówka in the Góry wi tokrzyskie Mts (in Polish with English summary). Kwart. Geol., **30** (2): 201–228.
- KOWALCZEWSKI Z., KULETA M. and MOCZYDŁOWSKA M. (1984) — Charakterystyka formacyjna osadów kambru dolnego w południowo-zachodniej Małopolsce. Arch. Pa stw. Inst. Geol. Kielce.
- KOWALCZEWSKI Z., KULETA M. and MOCZYDŁOWSKA M. (1987) — Nowe dane o kambrze dolnym z okolic Katuszowa i Korytnicy w Górach wi tokrzyskich. Kwart. Geol., **31** (1): 225–226.
- KOWALCZEWSKI Z., LISIK R. and CHLEBOWSKI R. (1976) — New data on the geological structure of the Opatów area (in Polish with English summary). Biul. Inst. Geol., **296**: 167–207.
- LENDZION K. and JANKAUSKAS T. (1992) — Lower and Middle Cambrian Acritarch biozonation of the Baltic Syneclise and adjacent areas (East European Platform). Prz. Geol., **40** (9): 519–525.
- LENDZION K., MOCZYDŁOWSKA M. and AKOWA H. (1983) — Wiek utworów z otworu wiertniczego Bazów IG-1 w wietle nowych danych. Kwart. Geol., **27** (2): 451–452.
- MARTIN F. (1992) — Uppermost Cambrian and Lower Ordovician acritarchs and Lower Ordovician chitinozoas from Wilcox Pass, Alberta. Geol. Surv. Canada, Bull., **381**.
- MARTIN F. and DEAN W. T. (1981) — Middle and Upper Cambrian and Lower Ordovician acritarchs from Random Island, eastern Newfoundland. Geol. Surv. Canada Bull., **343**: 1–43.
- MARTIN F. and DEAN W. T. (1988) — Middle and Upper Cambrian acritarch and trilobite zonation at Manuels River and Random Island eastern Newfoundland. Geol. Surv. Canada, Bull., **381**: 91.
- MIZERSKI W. (1992) — Tectonics of the Cambrian sequence in the Holy Cross Mountains (in Polish with English summary). Prz. Geol., **40** (4): 142–146.
- MIZERSKI W. (1994) — Paleotectonic evolution of the Cambrian of the Holy Cross Mts (Central Poland) (in Polish with English summary). Prz. Geol., **42** (9): 721–727.
- MOCZYDŁOWSKA M. (1995) — Late Cambrian Acritarchs and their age constraints on an Ediacaran — type fauna from the Booley Bay Formation, Co. Wexford, Eire. Geol. J., **30**: 111–128.
- MOCZYDŁOWSKA M. (1998) — Cambrian acritarchs from Upper Silesia, Poland — biochronology and tectonic implications. Fossils and Strata, **46**: 1–121.
- ORŁOWSKI S. (1964) — The Middle Cambrian and its fauna in the eastern part of the Holy Cross Mts (in Polish with English summary). Stud. Geol. Pol., **16**.
- ORŁOWSKI S. (1975) — Cambrian and Upper Precambrian lithostratigraphic units in the Holy Cross Mts. (in Polish with English summary). Acta Geol. Pol., **25** (3): 431–448.
- ORŁOWSKI S. (1985) — New data on the Middle Cambrian trilobites and stratigraphy in the Holy Cross Mts. Acta Geol. Pol., **35** (3–4): 251–262.
- ORŁOWSKI S. (1988) — Stratigraphy of the Cambrian system in the Holy Cross Mts. Kwart. Geol., **32** (3/4): 525–530.
- ORŁOWSKI S. and MIZERSKI W. (1995) — Once more about geology of the Góra Wi niówka outcrops Holy Cross Mts, Central Poland (in Polish with English summary). Prz. Geol., **43** (1): 11–14.
- PO ARYSKI W. and TOMCZYK H. (1993) — Geological cross-section through SE Poland (in Polish with English summary). Prz. Geol., **41** (10): 687–695.
- PO ARYSKI W., VIDAL G. and BROCHWICZ-LEWISKI W. (1981) — New data of the Lower Cambrian at the southern margin of the Holy Cross Mts. (SE Poland). Bull. Acad. Pol. Sc. Sér. Sc. Terre., **29** (2): 167–174.
- SAMSONOWICZ J. (1934) — Objasnienie arkusza Opatów ogólnej mapy geologicznej Polski w skali 1:100 000. Pa stw. Inst. Geol. Warszawa.
- STUPNICKA E. (1988) — On the nature and origin of the wi tokrzyskie Dislocation (in Polish with English summary). Prz. Geol., **36** (1): 40–45.
- SZCZEPANIK Z. (1997a) — Palinologia ordowiku Gór wi tokrzyskich. Arch. Pa stw. Inst. Geol. Kielce.

- SZCZEPANIK Z. (1997b) — Preliminary results of thermal alternation investigations of the Cambrian acritarchs in the Holy Cross Mts. *Geol. Quart.*, **41** (3): 257–264.
- TOMCZYKOWA E. (1968) — Stratigraphy of the uppermost Cambrian deposits in the wi tokrzyskie Mountains (in Polish with English summary). *Pr. Inst. Geol.*, **54**.
- VANGUESTAINE M. (1974) — Espèces zonales d'acritarches de Cambro-Tremadocien de Belgique et de l'Ardenne Française. *Rev. Palaeobot. Palynol.*, **18**: 63–82.
- VANGUESTAINE M. and Van LOOY J. (1983) — Acritarches du Cambrien Moyen de la Vallée de Tacheddirt (Haut-Atlas, Maroc) dans le cadre d'une nouvelle zonation de Cambrien. *Ann. Soc. Geol. Belgique*, **106**: 69–85.
- VAVRDOVÁ M. (1982) — Phytoplankton communities of Cambrian and Ordovician age Central Bohemia. *Ustr. Ust. Geol.*, **57** (3): 145–155.
- VOLKOVA N. A. (1990) — The Middle and Upper Cambrian acritarchs in the East European Platform. *Nauka. Moskva*.
- VOLKOVA N. A. and GOLUB I. N. (1985) — Novye akritarchy verhnego kembrija Leningradzkoj oblasti (ladoskaja svita). *Paleont. Zur.*, **4**: 90–98.
- WELSCH M. (1986) — Die Acritarchen der Höheren Digermul-Gruppe, Mittelkambrium bis Tremadoc, Ost-Finnmark, Nord-Norwegen. *Paläontographica*, **B 201**: 1–109.
- ZNOSKO J. (1996) — Tectonic style of the Early Palaeozoic sequences in the Holy Cross Mountains. *Kwart. Geol.*, **40** (1): 1–22.
- AK C. (1962) — Preliminary study of the Middle Cambrian in the Góry Pieprzowe Mts. (In Polish with English summary). *Biul. Inst. Geol.*, **174**: 9–50.