



## The maximum ice sheet limit of the Vistulian Glaciation in northeastern Poland and neighbouring areas

Tomasz KRZYWICKI



Krzywicki T. (2002) — The maximum ice sheet limit of the Vistulian Glaciation in northeastern Poland and neighbouring areas. *Geol. Quart.*, 46 (2): 165–188. Warszawa.

This paper outlines a new approach to reconstructing Vistulian ice sheet limits in NE Poland and adjacent areas of Lithuania and Belarus. It is based on geological investigations of landforms and deposits either of a hypothetical boundary to the Vistulian ice sheet limits. The main conclusions are: 1 — the area was covered twice by the Vistulian ice sheet; the first time during the wiccie Stadial (67–50 ka BP) and the second time during the Main Stadial (24–12 ka BP); 2 — the limit of the older stadial was farther to the south a that of the younger one; 3 — due to covering of the Eemian sediments by the ice sheet of the wiccie Stadial the limit of the Vistulian Glaciation has moved about 15–20 km southwards.

Tomasz Krzywicki, Polish Geological Institute, Rakowiecka 4, 00-975 Warszawa, Poland, e-mail: tkrz@pgi.waw.pl (received: June 13, 2001; accepted: December 11, 2001).

Key words: northeastern Poland, Vistulian Glaciation, Eemian Interglacial, ice sheet limit, tills.

### INTRODUCTION

This paper deals with the limits of the Vistulian Glaciation in northeastern Poland and neighbouring territories of Lithuania and Belarus. It is a summary of the Ph. D. thesis (Krzywicki, 1999b).

The aim of the work was to investigate which of the ice sheets of the Vistulian Glaciation advanced onto the area of northeastern Poland, as well to reconstruct and delimit its maximum extent.

The Oldest (Toru ) Stadial, two thin glacial tills found by Makowska (1976) and Drozdowski (1986) in the Lower Vistula region, is not considered in this paper, which deals with the ice sheet limits of the two Vistulian Glaciations: the wiccie Stadial and the Main Stadial.

Many Polish authors have suggested that the ice sheet of the wiccie Stadial extended into northern Poland, though its presence east and south of the Mazury Great Lakes District has not been proved. Both Lithuanian and Belarussian authors suggest that there was no ice sheet in their countries at that time

(Astapova *et al.*, 1987; Karabanov, 1987; Valtchik *et al.*, 1990; Gaigalas, 1995; Kondratiene, 1996; Satkunas *et al.*, 1997).

The other ice sheet of the Vistulian Glaciation that certainly advanced onto the area of northern Poland was that of the Main Stadial (Leszno-Pomeranian). At the end of the glaciation a continuous retreat of the ice sheet occurred, interrupted by short stagnation events (of the rank of phases and subphases) marked by moraine belts. A similar idea was earlier postulated by urek (1975), Ber (1990) and Lisicki (1993). The stagnation periods were usually stable, rarely with oscillatory fluctuations, locally of a transgressive character. It seems that the only exception was the local Wigry transgression that occurred between the maximum advance of the ice sheet and the Pomeranian Phase (Lisicki, 1993).

In the north the research area embraces uplands and outwash plains of young-glacial relief. These are fragments of the Mr gowo Lakeland and Mazury Great Lakes District, Mazury Plain, part of the Elk Lakeland and Rajgród Lakeland, as well as the Augustów Plain, Dainava Plain and part of the Central Neman Lowland. In the south, the study area covers fragments of old-glacial uplands of the Kolno Upland, Białystok Upland, Sokółka Hills, Hrodna Upland and Lida

Plain, with intervening depressions of the Kurpie Plain, Biebrza Valley, High Neman Lowland. The research area covers 26 sheets of the *Detailed Geological Map of Poland*, scale 1:50 000 and occupies a total area of approximately 7700 km<sup>2</sup>. The Lithuanian and Belarussian parts cover about 3000 km<sup>2</sup>.

## PREVIOUS RESEARCH

Before 1939 Polish investigations of the last glaciation in northeastern Poland were limited to the Suwałki area (Wołosowicz, 1926, 1928; Pietkiewicz, 1928; Pawłowski, 1933). Woldstedt (1935) published a map of the Quaternary deposits of East Prussia with the ice sheet limit of the Vistulian Glaciation marked, trending along the Narew River valley near Łomża, south of Mołki and north of Sokółka (Fig. 1). The maximum extent of the Pomeranian Stadial was shown as running across the northern margin of the Mazury Lakeland. The first Polish attempt to summarise the Quaternary studies of East Prussia was the paper by Galon (1937), which also included a map of morphological landscapes of East Prussia constructed on the basis of Woldstedt's map (1935). Galon (1937) included the whole Lakeland within the extent of the ice sheet of the Pomeranian Stadial, with its limit running across the area of Mikołajki, Orzysz and Elk.

After 1939, studies of Quaternary deposits and extents of ice sheets in northeastern Poland became more systematic. Majdanowski (1947) drew the ice sheet limit of the Vistulian Glaciation south of Szczytno, north of Rozogi, between Pisz and Biała Piska, north of Grajewo, south of Rajgród, north of Sztabin and along the Wołkuszanka River valley. Halicki (1950, 1951) recognised, on the basis of palynological studies, deposits of two more glaciations above Eemian Interglacial deposits: Glaciation V (North Polesie or North Polish Glaciation) and Glaciation VI (Baltic Glaciation). The former reached the area of Supraśl, Gródek and Jałówka (Halicki, 1951). Farther east, Belarus, Halicki drew the limit of Glaciation V across Svisloch, Porozow, Lyskov, Ruzhana, Byten and south of Baranovitchi. Pachucki (1961) mapped the Neman Ice Lobe in the Suwałki area. Kondracki (1952) distinguished 9 belts of frontal moraines deposited during the Baltic Glaciation. That author drawn its maximum ice sheet limit across Nidzica, Szczytno, Niardwy Lake and Elk.

Since the beginning of the 1960s most authors assumed, after Galon and Roszkówna (1961), that the maximum limit of the Baltic Glaciation in northeastern Poland coincided with the Leszno Phase. Those authors drew the limit line between Lipowiec and Szczytno, north of Rozogi, across Jędrzejów, north of Szczuczyn, north of Grajewo, south of Rajgród, across the northern slopes of Sztabin "Island" and south of the Wołkuszanka River valley. A general outline of this limit remained basically unchanged in later papers, although several authors introduced some modifications. The limits were drawn across the area of Szczytno, Rozogi, Jędrzejów, Grajewo, Rajgród and the southern part of the Augustów Plain. In the Kolno Upland, Szafer (1955), Rühle and Sokołowska (1961), Mojski and Rühle (1965) and Mojski (1972) drew the limit of

the Leszno Phase slightly farther south, across the area of Szczuczyn and Woszczyn, approaching the macroforms described later by Musiał (1984) as the slope of a marginal contact and glaciofluvial kame rampart.

In the Kurpie Plain, Rühle and Sokołowska (1961), Słowska (1971), Bogacki (1976) and Kozłowska and Kozłowski (1995b) drew the ice sheet limit of the Baltic Glaciation across "old" frontal moraines surrounded by the youngest outwash plain deposits.

Galon and Roszkówna (1961), Kalniet and Karaszewska (1972), Różycki (1972a, b, 1978), Mojski (1972), Straszewska (1975), Wołk-Musiał (1980) and Urek (1984) drew the line across the northern border of the Biebrza Valley. Urek (1975), moreover, considered that the ice sheet entered the Biebrza marginal valley as far as the Elk and Jędrznia rivers, as shown by the Kuwasy kettle hole.

Ber (1972a, b, 1974, 1982) and Urek (1975) claimed that the ice sheet limit of the Leszno Phase in the southern part of the Augustów Plain ran across the northern slopes of the "Islands" of Sztabin, Jastrzębna, Nowy Lipsk and Lipsk. A similar opinion was expressed by Bogacki (1976), who assumed that the ice sheet reached the Biebrza Valley, covering the whole of the Augustów Plain.

Ber (1975b) reported that the ice sheet limit of the Leszno Phase which, as he assumed, coincided with the maximum extent of the North Polish Glaciation, trends across the Kolno Upland and along the northern end of the Biebrza Valley (from Cwaliny through Kowalewo, Łodygowo, Borkowo, Grajewo, Danowo, Kosyły, Drastwo Lake, Wołnawie and Tajno Lake). Recently, Ber (1999, 2000), quoting Liszkowski (1993), has considered that glacioisostasy had an impact on the maximum extent of the Vistulian Glaciation and the position, direction and extents of recession-oscillation marginal zones. Glacioisostasy operated primarily along longitudinal tectonic faults of the deeper basement. The marginal zone of the Main Stadial of the Vistulian Glaciation coincides, according to Ber (1999, 2000), with the longitudinal Sajno-Rygl fault.

Marcinkiewicz (1973) reconstructed the extent of young-glacial relief on the basis of the analysis of topographic maps of the South Mazury and the Suwałki area, at a scale of 1:25 000, understanding its limit as a boundary between an area abounding with kettle holes and an area with a regular drainage system. Marcinkiewicz (1973) suggested that this boundary corresponded to the maximum extent of the last glaciation.

Kociszewska-Musiał (1978) alone assumed that the Vistulian ice sheet advanced only onto the East Suwałki Lakeland, whereas in the western part it filled major depressions and valleys.

In recent years there were also some attempts to draw the ice sheet limit of the Vistulian Glaciation farther to the south. These attempts were based on thermoluminescence dating, and were stimulated primarily by the consciousness that there is a difference (*cf.* Halicki, 1950, 1951) between the degree of denudation in the zone typical of old-glacial areas and that in the area located to the north — although outside the commonly accepted extent of the Baltic Glaciation. These are the Mława and Przasnysz areas, the Białystok Upland and the Sokółka Hills, as well as the Biała Piska region and northern areas of the Kolno




Fig. 1. Location of research area and maximum ice sheet limits of Vistulian Glaciation (Nemunas [Neman], Poozierie) according to different authors




Fig. 2. Borehole

Upland. Research workers emphasize the freshness of the landscape of the area, the varied topography, steep slopes and a dense system of small valleys.

Fedorowicz *et al.* (1995), using TL datings, ascertained that the ice sheet front of the Vistulian Glaciation may have reached the area of Białystok. Banaszuk (1995) based his stratigraphical conclusions mainly on TL datings from the Knyszyn Forests, young-glacial areas of northeastern Poland (Banaszuk *et al.*, 1994) and the Narew River valley (Banaszuk, 1996). The results of his investigations show that the Toruń Stadial ice sheet had the greatest extent of all the Vistulian ice sheets (approximately 110 ka BP), reaching Brańsk, Wysokie Mazowieckie and the Nurzec River. The Świecie Stadial ice sheet covered only the East Suwałki Lakeland, and it was the last glaciation in that region. It is possible that the ice sheet front of the Main Stadial stopped in the extreme north of the lakelands, with ice tongues moving southwards into depressions.


After the publications by Makowska (1976, 1979, 1980) on the stratigraphy of the Vistulian Glaciation deposits in the Lower Vistula region, studies on the possible greater extent of

tills older than the Main Stadial were also initiated in the neighbouring areas.

Marks (1988) discovered that the ice sheet W3, coeval with the Świecie Stadial (60–55ka BP), reached the areas of Płock, Lidzbark Warmiński, Nidzica and Szczytno. In Warmia and the West Mazury Lakeland, the limit of the ice sheet W3 coincides with the maximum extent of the Vistulian Glaciation. In succeeding papers Marks (1990a, b, 1991, 1994, 1997a, b) upheld these ideas. Lindner (1992) was of the opinion that the Świecie Stadial ice sheet advanced again onto the Lower Vistula region and, possibly, onto Warmia and the northern part of the Mazury Lakeland. On the basis of petrographical studies and examinations of erratics Lisicki (1998) and Gałązka *et al.* (1998) moved the hypothetical line of the maximum extent of the Świecie Stadial southwards to the Mława area.

There is no controversy among geologists on the extent of the last glaciation (Poozerie Glaciation) in Belarus. The ice sheet limit was drawn by Voznyatchuk (1956) and upheld by Vaytekunas (1968) and Voznyatchuk *et al.* (1972), and is commonly accepted by Belarussian research workers. It runs across the northern slopes of the Hrodna Upland, enters the Neman River valley near Hrodna and then turns southwards to the





type sections

Aziory area (hence the name of the ice lobe). Goretskiy (1983) and Karabanov (1987) moved the line several to tens of kilometres southwards. There is a common opinion that the ice sheet did not enter the Neman River valley south of Skidel. Only Chepulite (1978, 1986) claimed that the youngest glaci- ation tills spread as far as south of the Dainava Plain, the Central Neman Lowland and Hrodna Upland.

Astapova *et al.* (1987) considered that the ice sheet of the Late Poozerie Stadial had its greatest extent during the Poozerie Glaciation (Baltijskij horizon, Orshanskij Megastadial) 17–18 ka BP.

Belarussian scientists (Karabanov, 1987; Astapova *et al.*, 1987; Valtchik *et al.*, 1990) have not found any deposits and geomorphological features of the early Poozerie Glaciation. According to Astapova *et al.* (1987), the ice sheet of the Srednie-Poozerskij Megastadial (Dvinskij subhorizon), that corresponds to the Świecie Stadial, probably reached the north- ern border of Belarus, not overstepping it.

Lithuanian geologists and geographers, among others Gaigalas (1995), Kondratiene (1996) and Satkunas *et al.* (1997), agree that ice sheet transgressions occurred only during the later phase (Upper Nemunas) of the Nemunas Glaciation. They distinguished the Gruda Stadial (including Žiogeliai

Phase) and Baltija Stadial. The stadials reflect two ice sheet ad- vances, of which the former had a greater extent.

## RESEARCH METHODS

The present report is based on the analysis of geological in- vestigations of deposits and landforms in the broadly understood zone of maximum extent of the Vistulian Glaciation, the analysis of topographic maps with particular regard to geomorphological problems, an analysis of the results of petrographical-lithological investigations (used to construct the *Detailed Geological Map of Poland*, scale 1:50 000), palynological studies of Eemian Inter- glacial sites, and thermoluminescence datings of tills, lacustrine and ice-dam lake deposits.

Topographic and geomorphological maps, at a scale of 1: 25 000, were also analysed. These gave evidence for distri- bution of frontal and dead-ice moraines, kames, eskers and gla- cial tunnel valleys, outwash fans, moraine “islands” of an older morainic plateau within outwash plain deposits, outcrops of a till basement within outwash plain and ice-dam deposits, topo- graphic lows which in the past may have formed marginal val-




Fig. 3. Research area map




Fig. 4. Geological cross-section A–B: Mrągowo Lakeland, Kurpie Plain

leys or drainage routes for the pra-Neman (Nemunas) River, channel lakes and dead-ice troughs.

62 cartographic borehole sections were analysed from the Polish part of the research area, and several tens of borehole sections from Belarus and Lithuania. 24 type sections were selected. These are characterised by the following features:

- the profiles were analysed for palynology and age determinations;

- the age of any one of the lithologic horizons is supported by 2 methods, e.g. the lithotype is determined using petrographical studies with additional TL datings;

- they possess the most complete sequences of the Vistulian Glaciation deposits;

- they show a characteristic development of deposits, most typical for the study area.

These profiles prove the age of Late Pleistocene deposits, in particular the Vistulian Glaciation and the Eemian Interglacial deposits, throughout the study area. In some cases they also il-

lustrate the typical geological structure of the sub-surface Pleistocene deposits, and form a basis for the construction of geological cross-sections and further analyses (Fig. 2).

Especially detailed analyses were made on 16 sheets of the *Detailed Geological Map of Poland*, at a scale of 1:50 000, (Maksiak, 1992; Kozłowska and Kozłowski, 1993, 1995a, b, 1996a, b, 1997, 1999; Krzywicki, 1996, 1999a, 2001a, b; Łuk, 1997, 1998a, b; Kacprzak, 2000; Lichwa, 2000; Ber, in preparation). Geological documentations were also analysed, providing much information on the lithology of end moraines, kames and outwash fans.

The aim of the work was to study the youngest glacial deposits, including tills. Their distribution, extent, thickness and lithology were examined. Supposed extents of the youngest ice-dam lakes, which came into existence during the transgression and regression of the youngest glaciation ice sheet, as well as the drainage routes of the ancient Neman River waters to the drainage systems of the Biebrza, Narew and Vistula during the




Fig. 5. Geological cross-section C–D: Augustów Plain, Biebrza Valley, Sokółka Hills

ice sheet retreat of the Vistulian Glaciation, were also investigated. A number of geological cross-sections were constructed. They show the geological structure of the Mid-Late Pleistocene deposits in the research area (Figs. 3–5).

**PALYNOLOGY**

Approximately 45 Eemian Interglacial sites have been described so far and investigated within the zone of the maximum extent of the Vistulian Glaciation in northeastern Poland, western Belarus and southwestern Lithuania. These are sites of lacustrine organic sediments both uncovered and covered by glacial deposits, most often by the Vistulian Glaciation till, its residuum or glaciofluvial deposits (Fig. 6). The climatic optimum of the Eemian Interglacial is recorded in 10 sites, and in 2 more

it is probable. The thickness of the interglacial deposits varies from 0.5 to 7.2 m (Fig. 7a, b).

In the Hrodna Upland, and High and Central Neman Lowland interglacial deposits fill abandoned river channels in the Neman River valley (Pavlovskaya, 1998). Some Belarussian and Lithuanian geologists suggest that in this area the whole of Eemian Interglacial deposits are overlain by a till or its residuum (Voznyatchuk, 1956; Chepulite, 1966; Kryger *et al.*, 1971). Other authors (Voznyatchuk, 1960; Pasyukievich *et al.*, 1972; Goretskiy, 1980) considered that some of the Eemian Interglacial sites (Pyszki, Poniemu and Druck) possess only a periglacial solifluction cover. Karabanov (1987) claims that there is no precise answer to the question whether the Eemian deposits are covered by a till or not in the peripheral zone of the Hrodna Upland near the Neman River. In western Belarus, the most widespread Eemian deposit is bog lime, often covered with peat, 1–8 m in thickness (Karabanov, 1987).






Fig. 6. Map of the Eemian Interglacial sites

Sites of the Eemian Interglacial: a — not covered with glacial sediments: in the **Mława Upland** — Nidzica (Marciniak and Kowalski, 1978), Nidzica Sewerynow (Mamakowa, 1988), Rogó ek, Rozdro e and Bartoszek (Morawski *et al.*, 1998, 1999); in the **Kolno Upland** — Niewodowo (Musiał *et al.*, 1982), Gr dy Woniecko (Banaszuk, 1980), Rakowo Nowe (Bałuk and Krupi ski, 2000); in the **Białystok Upland** — Białosuknia-Szlachta (Janczyk-Kopikowa, 1958; Ber, 1975b), Bagno-Kalinówka (Borówko-Dłu akowa and Halicki, 1957), Białousy (Janczyk-Kopikowa, 1958), Machnacz ( urek, 1990,1992; Kupryjanowicz, 1991,1994; Fedorowicz *et al.*, 1995), Czarna Wie Ko cielna (Bitner, 1956; Buczy ski, 1960); in the **Sokółka Hills** — Ludomirowo, Zacisze and Mikelewszczyzna (Czaplicka, 1952; Bitner, 1957; Ber, 1972b); in the **Hrodna Upland** — Rumłówka ( rodo , 1950; Karabanov, 1987), Grandicze (Pasyukievich *et al.*, 1972), Małoje Demidkowo, Kamienka, Czeszczewłany; in the **Central Neman Lowland** — Białe Jezioro (Karabanov, 1987), Komotowo (Voznyatchuk, 1956; Dorofieyev, 1963); in the **Lida Plain** — Psiarce and Kurki (Pasyukievich *et al.*, 1972); b — covered with glacial or fluvioglacial sediments: in the **Elk Lakeland** — Lipi skie (Janczyk-Kopikowa, 1998); in the **Suwałki Lakeland** — Szwajcaria, Błaskowizna and Smolniki (Borówko-Dłu akowa and Halicki, 1957; Andrzejeszczak, 1971; Borówko-Dłu akowa, 1971; Ber, 1973, 1974; Kociszewska-Musiał and Leciejewicz, 1982; Krzywicki, 1993); in the **Lithuania Lakeland** — Netiesos, Maksimony and Jonionys (Bremówna and Sobolewska, 1950; Borówko-Dłu akowa and Halicki, 1957; Chepulite, 1986; Kondratiene, 1996; Satkunas, 1997b; Satkunas *et al.*, 1998); in the **Dainava Plain** — Ratny ia, Druskininkai, Druskininkai-Pušinas, Liškiava, Smalnikai (Kondratiene, 1996); in the **Central Neman Lowland** — Rusota-Sołochy (Karabanov, 1987); in the **High Neman Lowland** — Pyszki (Chepulite, 1966), Druck (Halicki, 1951; Chepulite, 1986), Poniemu (Karabanov, 1987), Szczecinowo (Halicki, 1951; Chepulite, 1986), Siwkowo (Halicki, 1951; Chepulite, 1986), Kowalce (Halicki, 1951; Chepulite, 1966), ukiewiczze (Halicki, 1951), Bohatyrowicze (Halicki, 1951; Voznyatchuk and Valtchik, 1978; Chepulite, 1986)






Fig. 7. Eemian Interglacial profiles: **a** — non covered with glacial sediments, **b** — covered with glacial or fluvioglacial sediments




Fig. 8. Diagram of the coefficients O/K and K/W in the upper tills of NE Poland

The sites at Czeszczewlany, Białe Jezioro and Komotowo are undoubtedly located within the extent of the youngest ice sheet. The lack of a till horizon above Eemian Interglacial deposits results probably from later erosional and denudation processes.

A geological (palaeobotanical) criterion is one of the most important criteria permitting the delineation of the ice sheet limit, showing that the Vistulian ice sheet did not advance onto the Mazovian and Podlasie lowlands (Banaszuk, 1995, 1996, 1998; Fedorowicz *et al.*, 1995).

#### LITHOSTRATIGRAPHY OF TILLS

The major feature that allows the distinction between tills is the petrographical composition of the gravel fractions. This paper compares the petrographical composition of pebbles from the uppermost tills from the Polish part of the research area, using this data to construct a stratigraphical succession.

The methodology of petrographical studies was outlined by Lisicki (1998). Samples must contain at least 100 grains, 5–10 mm in diameter. The analysis includes the calculation of the percentage contribution of particular groups of Scandinavian rocks (crystalline rocks, limestones, dolomites, sandstones, quartzites and quartz grains) and local rocks (limestones, marls, sandstones, siltstones and claystones). Proportions between different rocks in each sample are characterised by petrographic coefficients: O/K, K/W and A/B, where: O — total of sedimentary rocks (limestones, dolomites, sandstones and shales), K — total of crystalline rocks and quartz grains, W — total of carbonates (limestones, dolomites), A — total of rocks non-resistant to weathering (limestones, dolomites, sandstones and shales), B — total of weathering-resistant rocks (crystalline rocks, quartz grains and sandstones).

Results of lithological-petrographical investigations performed for the following sheets of the *Detailed Geological Map of Poland*, at a scale of 1:50 000, were analysed: Wielbark (Jele ski, unpubl.), Lipowiec (Zawidzka, unpubl.), wi tajno and Spychowo (Fert *et al.*, unpubl.), Myszyniec (Komacka-Makowiecka, unpubl.), Ruciane-Nida, Łyse, Pisz and Kolno (Makarewicz and Zaczekiewicz, unpubl.), Biała Piska (Honczaruk, unpubl.), Rajgród, Rygol and Rudawka (Gronkowska-Krystek, unpubl.), Augustów Station (Kenig, unpubl.). Vistulian Glaciation tills are absent from the boreholes in the western part of the research area (near Ruciane-Nida, Łyse, Pisz and Kolno).

Furthermore, a special report was prepared (Gronkowska-Krystek, unpubl.) for 26 till samples collected for investigations in the southern part of the Augustów Plain and Upper Biebrza Valley (near Sztabin and Lipsk) (Krzywicki, 1997a, b).

Petrographical analyses of gravels from 62 cartographic boreholes, hand-auger holes and exposures were performed. Petrographic coefficients were calculated for each sample. Arithmetic means of these coefficients for individual till horizons were also calculated. Figure 8 shows the mutual relationships between the groups of coefficients. 5 lithotypes can be distinguish among the uppermost till horizons: B<sub>2</sub>, B<sub>1</sub>, B<sub>1bis</sub>, W<sub>2</sub> and W<sub>1</sub>. Lithotype B<sub>1</sub> occurs only east of Suwałki and Augustów (Table 1). Figure 9 shows examples of borehole sections containing tills of various lithotypes. These results were compared with the results of petrographical investigations conducted by Kenig (1998) in the Mazury and Suwałki lakelands (Fig.10).

Values of petrographical coefficients O/K, K/W and A/B (within the same lithotype) differ slightly from one study to another. The following conclusions arise from the analysis of petrographical investigations:

1. They may have a great utility in the lithostratigraphy of tills.

2. Tills B<sub>2</sub> can be correlated with the younger stadial of the Vistulian Glaciation, tills B<sub>1</sub> and B<sub>1bis</sub> correlate with the older stadial of the Vistulian Glaciation, tills W<sub>2</sub> correlate with the younger stadial of the Wartanian Glaciation, tills W<sub>1</sub> correlate with the older stadial of the Wartanian Glaciation.

3. It should be stressed that the youngest till (lithotype B<sub>2</sub>) occurs in the fresh morainic plateaux with glacial tunnel valleys. This supports Majdanowski's proposal (1947) that glacial tunnel valleys are the best indicator of the limit of the youngest glaciation, considered here as the Main Stadial of the Vistulian Glaciation, 32–13 ka BP (cf. Shackleton and Opdyke, 1973). In 4 cases (Kiejkuty, Wojnowo, Racibórz and Pikły) out of 6, tills B<sub>2</sub> are underlain by tills B<sub>1</sub>.

4. Tills B<sub>1</sub> or B<sub>1bis</sub> from the area located outside the maximum extent of the Main Stadial always form the highest horizon (except for two cases: the Krasne and Hanus boreholes in the Augustów Plain, where the upper till is weathered and its lithotype cannot be defined).

5. In the Augustów Plain, Odranian Glaciation tills are situated very high (boreholes Kolnica, Kolonia Jastrz bna, Jastrz bna II, Nowy Lipsk), and they are often overlain by tills B<sub>1bis</sub> (in the last 3 boreholes). In Krasnybór, tills W<sub>2</sub> occur near the surface and are overlain by tills B<sub>1bis</sub>.

6. Deviations from the average values of petrographic coefficients O/K, K/W and A/B are very rare. They can result from the occurrence of detached till fragments of a different age. Till from depths of 13.8–15.4 m (Nowy Lipsk borehole), with coefficients of 0.77–1.61–0.54, can serve as an example.

7. In at least 7 borehole sections, the upper tills are weathered and the results had to be rejected. Weathering processes operated over some areas during glaciations (periglacial climate) and warmer periods (moderate climate), and were probably influenced by local geomorphological and microclimatic conditions. Weathering zones often occur in the upper parts of the youngest tills.

8. In 10 borehole sections from the areas of Ruciane, Pisz, Łyse and Kolno, the first till (from top) is the Odranian Glaciation till.

## TL DATINGS

The thermoluminescence method (TL) was broadly used for absolute age determinations in the 1980s and early 1990s. It was primarily employed to examine clays and muds, but also tills and sands. This was a standard method to establish the stratigraphy during the preparation of the *Detailed Geological Map of Poland*.

The author had at his disposal results of 34 TL dates of near-surface deposits from the study area. 17 of them come from the Augustów Plain and Upper Biebrza Valley. One dates was kindly rendered accessible to me by Professor urrek. Another one comes from the paper by urrek (1990). The most important datings for the present study are those from the Augustów Plain and Upper Biebrza Valley (Fig. 11). They can be grouped into 4 time groups.

Table 1

The coefficients O/K, K/W and A/B in the upper tills of NE Poland

Litho-type	Number of samples	Mean values of petrographical coefficients O/K- K/W-A/B
B <sub>2</sub>	7	2.39–0.45–2.14
B <sub>1</sub>	21	1.92–0.55–1.71
B <sub>1bis</sub>	13	1.38–0.80–1.10
W <sub>2</sub>	22	1.69–0.62–1.49
W <sub>1</sub>	11	1.53–0.72–1.38

The first group includes only one date of 180.0 ka BP ( urrek, 1990) (profile 1 in Fig. 11). The second group contains 2 dates — 118.4±17.8 (UG-2606) and 123.0±18.4 ka BP (UG-2611) (profiles 2 and 3 in Fig. 11) (Fedorowicz, unpubl.). The third group includes 5 dates (profiles 4–8 in Fig. 11), from 47.0±7 to 63.0±9 ka BP (Lub-2088-90, UG-172) (Butrym, unpubl.; Olszak and Fedorowicz, unpubl.; Olszak, unpubl.). 4 of them were made on tills, one (the youngest) on clayey mud. The fourth group includes 4 dates from muds and muddy clays — from 23.5±3.5 to 31.8±4.8 ka BP (profiles 9–12 in Fig. 11) (UG-2607-09, 2619) (Fedorowicz, unpubl.).

The analysis of TL and petrographical investigations from the Augustów Plain and Upper Biebrza Valley were the subject of a separate paper (Krzywicki, 1997a). Its main conclusion was that tills occurring at the surface are not of the same age everywhere, and the “islands” are composed of 2 or more tills of different ages.

Dates from the Augustów Plain point to the occurrence of older and younger tills of the Wartanian Glaciation and one till of the wiccie Stadial. The younger dates from muddy deposits suggest the Grudzi dz Interstadial and Main Stadial, prior to the ice sheet advance.

TL dates can often help in stratigraphy when considered together with the results of other investigations.

If we assume the greater reliability of petrographical studies over TL datings, and tie down the above-mentioned groups of coefficients to particular lithotypes (resulting in stratigraphical conclusions), we must reject 2 of the thermoluminescence datings from the Augustów Plain.

## STRATIGRAPHY OF VISTULIAN GLACIATION DEPOSITS

Until the 1950s it was commonly accepted (Woldstedt, 1929; Kondracki, 1952; Szafer, 1953) that the last glaciation was divided into the Leszno, Pozna and Pomeranian phases (corresponding to latives of the Brandenburgian, Frankfurtian and Pomeranian phases in Germany). The Leszno Phase was considered to have the greatest extent.

Studies by Makowska (1976, 1979, 1980, 1986) resulted in the new stratigraphy of the Vistulian Glaciation in Poland. 5






Fig. 9. Examples of boreholes with different till lithotypes

tills distinguished by her were ascribed to 3 stadials (Toru , wiecie and Main stadials) or to 2 glaciations: Toru and Baltic (Vistulian) glaciations, which are separated by dated marine deposits. A warm period between the Toru and wiecie stadials was termed by Makowska the Gniew Interstadial or Krastudy Interstadial (1976, 1980), later named the Krastudy Interglacial (1986).

Results of Makowska's (1976, 1979, 1980, 1986) investigations in the Lower Vistula region were later partly upheld by Drozdowski (1986) and, in Warmia and the western part of Mazury, by Marks (1988). Marks recognised 3 tills of the Vistulian Glaciation (W<sub>2</sub>, W<sub>3</sub> and W<sub>4</sub>), corresponding to the tills BII, BIII and BIV of Makowska (1976).

A similar subdivision of the North Polish Glaciation was also accepted in the *Instruction for the Detailed Geological Map of Poland* (1991).

Currently, some Polish authors (Lindner, 1987; Marks, 1990a; Mojski 1997) accept stratigraphic schemes of the Vistulian Glaciation based on Makowska's papers with three main tills (1979, 1980). These schemes are well tied to stratigraphic schemes of the Upper Pleistocene of northern Europe (e.g. Behre, 1989), and to their correlation with <sup>18</sup>O isotope stages (Shackleton and Opdyke, 1973; Mangerud, 1992), as well as to the climatostratigraphic scheme proposed by Wysocza ski-Minkowicz (1982), and supported by age determinations.

The Belarussian authors use the stratigraphic scheme proposed by Voznyatchuk (1972, 1976). He divided the Poozerie (Valdai) Glaciation into 5 zones or subzones, equivalents to stadials and interstadials. Starting with the Middle Vistulian these are as follows: Dvinskij (Kalininskij, Srednie-Poozerskij Megastadial) subhorizon, Srednie-Poozerskij subhorizon (Krasnaya Gorka Megainterstadial) and Baltijskij horizon (Ozerie, North Poozerie, Late Poozerie, Poozerie, Poozerie Glaciation *sensu stricto*). More detailed stratigraphic schemes, based on palynological studies, have recently been worked out (Yelovicheva *et al.*, 1996, Yelovicheva, 1997; Yelovicheva and Sanko, 1999). Yelovicheva *et al.* (1996) divided the middle and upper parts of the Poozerie Glaciation into 2 parts: the Medium Level and Naroch Level — Late Glacial. The Medium Level is subdivided into 8 minor units: megastadials (cool periods) and interstadials (warm periods). Miezhinskij I Stadial corresponds to the wiecie Stadial. The Orshanskij IV Stadial corresponds to the Main Stadial. The latest paper by Yelovicheva and Sanko (1999) quotes a series of radiocarbon and TL dates from deposits of the Poozerie Glaciation in Belarus. These dates are not given in the chart of stratigraphic schemes of the Vistulian Glaciation (Fig. 12).

In Lithuania, there are two equivalent stratigraphic schemes of the Vistulian Glaciation, named the Nemunas Glaciation. The first scheme was proposed by Gaigalas and Yartsev (1992) and Gaigalas (1995). It refers to palynology sites in southern

and northern Lithuania (Jonionys, Rokai and Biržai). The Varduva Stadial (69 ka BP) is an equivalent of the wicielec Stadial. The youngest cool period is divided into 2 stadials: the Gruda Stadial (22 ka BP) with the Žiogeliai Phase, and Baltija Stadial (17 ka BP). The stadials are separated by the Pavyte Interstadial.

Guobyte and Pavlovskaya (1998) made an attempt to link the upper parts of the Lithuanian and Belarusian stratigraphic schemes of the latest glaciation (Upper Nemunas and Orshanskij Stadial of the Poozerie Glaciation, corresponding to the Main Stadial of the Vistulian Glaciation). Research around the Lithuanian/Belarusian border resulted in the conclusion that the upper stadial of the Nemunas/Poozerie Glaciation can be divided into the Gruda/Ozerie; Šventiany/Šventionys and Baltija/Braslav phases, corresponding to the Brandenburgian (Leszno), Frankfurian (Poznań) and Pomeranian phases.

Another Lithuanian stratigraphic scheme, postulated by Satkunas and Grigytė (1997), and correlated with other European stratigraphic schemes (Satkunas, 1997b), also refers to palynostratigraphy.

The Nemunas Glaciation was divided into the lower, middle and upper parts. Palynologically studied deposits of warm periods of the lower and middle Nemunas (7 units) were investigated in 2 sites at Jonionys-on-Nemunas near Merkinė and Mickunai near Vilnius. Warm periods are separated by 7 cool periods, labelled with the name of Nemunas followed by a number and a letter. A cool period corresponding to the wicielec Stadial is marked by a symbol Nemunas 2a. The upper Nemunas deposits (Nemunas 3) of the Gruda Stadial, Žiogeliai Phase and the Baltija Stadial are separated in this scheme by deposits of the Antaviliai and Ula interstadials (Satkunas, 1997a), although the stratigraphical position of these interstadials is unclear, being currently under revision.

The latest Lithuanian research (Blazhauskas *et al.*, 1998) has shown that organic deposits (peats, gyttja), exposed in high banks of the Ula River and earlier considered to have represented interstadial deposits, are younger and represent the Older Dryas, Allerød and Younger Dryas. Recent investigations of the Antaviliai section located near Vilnius (Satkunas and Hütt, 1999) also show that lacustrine deposits, correlated with the Antaviliai Interstadial, are older and represent the Drenthe-Warthe (Lublin) Interstadial. Satkunas and Hütt (1999) considered that there are no sites with interstadial or interphasial deposits of the younger part of Vistulian Glaciation in Lithuania and other Baltic countries. Therefore, there is no climatostratigraphical evidence for establishing stadials, phases and interstadials during that period.

#### VISTULIAN GLACIATION — RECONSTRUCTION OF CLIMATIC CHANGES AND ICE SHEET LIMITS

A reconstruction of geological events of the Early Vistulian Glaciation can only be approximate because no deposits of this age have been found in northeastern Poland.

During the period of 74–59 ka BP (*cf.* Mangerud, 1992) another cooling of the climate resulted in an ice-advance. In Poland this period is called the wicielec Stadial (Makowska, 1986)




Fig. 10. Minimal and maximal values of petrographic coefficients according to: a — Kenig (1998) and b — author

or Pre-Grudzi dz Stadial (Mojski, 1979). The wicielec Stadial ice sheet advanced, according to Lisicki (1996, 1997), onto the Mr gowo Lakeland from the north. Gał zka *et al.* (1998) consider that the till of the older Vistulian Glaciation contains more erratics from the Baltic-Cland and Finnish area, i.e. coming from the east, than from Sweden. A similar domination, but still more accentuated, is also observed within the younger till of the Vistulian Glaciation. It suggests that the centre of this glaciation was located to the east (Finland, Cland Islands) during both these stadials, but in the wicielec Stadial it moved to the western part of the area.

The wicielec Stadial ice sheet covered the northern part of Kurpie Plain as far as Kolno, W sosz and Klimaszewnica, released a small ice tongue into the Biebrza Valley, reached the front of Sokółka Hills in the vicinity of D browa Białostocka, and released an ice tongue into the Nurka River valley. In Belarus the ice sheet bypassed the Hrodna Upland, releasing an ice tongue into the Neman River valley as far as Hrodna. It also reached the valley east of Hrodna, covering it with a large ice lobe as far as the Svisloch River Mouths and Dubrowlany. From there, the eastern edge of the ice lobe ran towards Skidel, Ostryna and Motyle, towards the present Belarus/Lithuania border.

The extent was delimited taking into account one of the most important geomorphological criteria, namely, the occurrence of glacial troughs with lakes (Majdanowski, 1947). The absence of lakes in the old-glacial landscape (only this kind of landscape is observed within the area of the wicielec Stadial extent) is a very important indicator of landscape maturity. Only scarce relict lakes occur within that zone. It is associated with the period of filling of lake troughs with sediments (Wi ckowski, 1966). Another geomorphological criterion — the presence (in some areas of maximum extent) of outwash plains — showing a kame terraces character, morphometrically associated with marginal forms — was also considered.




Fig. 11. Borehole sections with TL dating




Fig. 12. Stratigraphical subdivision of the Vistulian Glaciation (Nemunas, Poozerie)

Among morphogenetic criteria, the occurrence of moraines, as well as the mutual relationships between end moraines and between moraine outliers on outwash plains were also taken into consideration. The most important group, however, are geological criteria. These include the extent of glacial till related to the ice advance. The wiece Stadial till was identified on the basis of petrographical investigations which comprised determining the percentage distribution of petrographic groups in gravels extracted from tills.

The thermoluminescence method (TL) enabled age determination of glacial and glaciolacustrine deposits within the area of the maximum ice sheet limit of the wiece Stadial. The distribution of sites of Eemian deposits, which are not covered by glacial deposits, allows identification of the potential maximum extent zone of the Vistulian Glaciation.

After the retreat of the wiece Stadial ice sheet, there began a warm period called the Grudzi dz Interstadial or Interpleniglacial. It lasted from 59 to 24 ka BP (cf. Mangerud, 1992). Deposits of that age are known from many sites




Fig. 13. Ice sheet limit of the Vistulian Glaciation in the Świecie Stadial and the Main Stadial






Fig. 14. Space-time schema of the Vistulian Glaciation in NE Poland and adjacent area

throughout Poland (Mojski, 1988), but the most characteristic section is a floristic succession from Konin-Maliniec I, where ancient flora is additionally <sup>14</sup>C-dated at > 42.9 ka BP (Kozarski, 1991). The Grudziądz Interstadial was characterised by a cool, subarctic climate. During its first phase, forestless tundra reigned with a willow-horsetail community. Later, stunted birch and herbs became dominant. Scattered birch-trees, typical of areas near the polar forest limit, appeared at the climatic optimum of the interstadial. There were also vast, open spaces overgrown with herbs and mugwort. At the end of the interstadial, stunted birch appeared again. Average temperatures in July for central Poland ranged from 10 to 11°C (Kozarski, 1991) during the climatic optimum.

Another cool period, called the Main or Leszno-Pomeranian Stadial, started at approximately 24.0 ka years BP. The Main Stadial ice sheet reached its maximum extent between about 18 and 20 ka BP (ice-dam lake deposits from Hołmia-on-Neman underlying the Poozerie Glaciation till, dated at 18–200 ka BP). The Pomeranian Phase is dated in the Suwałki Lakeland at approximately 15.5 ka BP. The ice sheet retreated from northeastern Poland at approximately 14.0–14.5 ka BP (Kozarski, 1991).

The ice sheet limit of the Main Stadial was reconstructed in this work largely on the basis of the occurrence of glacial tunnel valleys (Majdanowski, 1947). However, the limit was modified in many places because of use of other criteria, first of all geological and geomorphological ones.

Wi ckowski's studies (1966) on the sedimentation rate in lake troughs suggest that the relief was shaped during a period younger than the Świecie Stadial. During a period of 12 000–13 000 years, after the ice sheet retreat from the area of Mazury, Suwałki and Central Neman Lowland, most of lake troughs may have become filled with sediments only by 20–35 %.

The Main Stadial ice sheet covered the Mazury Plain as far as Szczytno, wi tajno, Spychowo, Karwica Mazurska, Turo I, Łacha, Je e and further along the high moraine belt separating the Kolno Upland from Elk Lakeland, as far as the town of Grajewo. From there, the ice sheet released a broad ice lobe into the Biebrza Valley as far south as the Białystok Upland near Goni dz. The eastern edge of that lobe ran along the Netta River valley to the villages of Bór and Gabowe Gr dy. The ice sheet covered the northern and middle parts of the Augustów Plain as far as Krasne, Ostry skie and Gruszki. North of Rudawka the ice sheet limit crossed the Poland/Belarus border. In Belarus the ice sheet reached the northern

slopes of the Hrodna Upland, released small ice tongues into the Neman and Hozhka River valleys, and reached the villages of Tabola and Lozy. Between Skidel and Ostrya the ice sheet limits of the Main and *wiecie* stadials ran along the same line.

Geomorphological, morphogenetic and geological criteria were used for delimiting the extent of the Main Stadial ice sheet. The most important were glacial tunnel valleys with lakes (Majdanowski, 1947). Assuming the lowest sedimentation rates and slow contraction of the lakes (Wi ckowski, 1966) we can suppose that the only lakes that could persist until the Holocene were those which formed after 50 ka BP, i.e. during the recession of the Main Stadial ice sheet. In the vicinity of Grajewo there is an outwash plain with a distinct outwash fan. This is the uppermost outwash plain level in that area, that is probably associated with the maximum limit of the Main Stadial ice sheet. Of morphogenetic criteria, leading to the determination of the origin and distribution of glacial forms, the occurrence of moraine belts from the area between Pisa and Grajewo were considered. The most important geological criterion was the distribution of till, identified by petrographical studies. The ice sheet limits of the Vistulian Glaciation are shown in [Figure 13](#).

Karabanov (1987) considered that the Hrodna Upland separated two ice lobes during the Brandenburgian (Leszno) Phase. To the west the Kursk Ice Lobe advanced from the Baltic Sea area (from NW), while to the east the Riga Ice Lobe advanced from the north. During the Frankfurian and Pomeranian phases, the Kursk Ice Lobe was separated into two parts in the Szeskie Heights area: the western part covered Mazury, whereas the eastern part formed an ice tongue reaching as far as the Šven ionys Upland in eastern Lithuania.

The ice sheet limit of the last glaciation was similarly drawn by Pachucki (1961). It ran from the vicinity of Wisztyniec, across Ha cza Lake, Krasnopol, Kap iamiestis, Merkine, in the vicinity of Vilnius, to Pabrade and Šven ionys. If such an eastern tongue could formed during the retreat of the Main Stadial ice sheet, it is probable that it may have also existed earlier, during the *wiecie* Stadial. The considerations are based on characteristic petrographic coefficients for the *wiecie* Stadial tills observed in the east (Augustów Plain, Upper Biebrza Valley), which are different than those calculated in Mazury. These characteristic petrographic coefficients have resulted in the separation of lithotype B<sub>1bis</sub>.

Therefore, it is highly probable that the *wiecie* Stadial ice sheet advanced onto eastern Mazury from N and NW, from western Finland, western Latvia and western Lithuania. The Suwałki and Hrodna areas were transgressed by the ice sheet advancing from the N, from Finland, Estonia, eastern Latvia nad eastern Lithuania. The area that separated both the ice tongues were the Szeskie Heights (or rather a basement elevation located west of Ha cza Lake and stretching to the south towards Suwałki), not the Hrodna Upland.

## SUMMARY AND CONCLUSIONS

This paper summarises knowledge on the near-surface geological structure (deposits of the middle and younger part of the

Vistulian Glaciation) of southeastern Mazury, the southern part of the Suwałki area and the Hrodna district. The major objective was to draw the ice sheet limits of the Vistulian Glaciation.

The main conclusions are as follows:

1. During the Vistulian Glaciation the ice sheet advanced twice onto the area of eastern Mazury and Suwałki and Hrodna districts: during the Middle Stadial (*wiecie* Stadial — approximately 55–67 ka BP) and during the Upper Stadial (Main Stadial — approximately 24–12 ka BP). The ice sheet extent was greater during the *wiecie* Stadial; it advanced from a few to about 30 km further to the south than the Main Stadial ice sheet ([Fig. 14](#)).

2. Results of petrographic composition of gravels from tills (lithotypes B<sub>2</sub>, W<sub>2</sub> and W<sub>1</sub>) tend to be highly convergent in the entire Polish part of the study area. Only the tills of the Middle Stadial of the Vistulian Glaciation have a different petrographic composition (lithotypes B<sub>1</sub> and B<sub>1bis</sub>). The results of petrographic research allowed a stratigraphic separation of the highest tills.

3. The ice sheet limit of the *wiecie* Stadial is marked by frontal moraines, outwash fans, kames and, indirectly, by sites of Eemian Interglacial sediments. In the eastern part of the research area the *wiecie* Stadial ice sheet rested against the older uplands that extended to the south.

4. The ice sheet limit of the Main Stadial is delimited by glacial tunnel valleys and lakes (except for a few small reservoirs located outside this line). Glacial sediments of the Main Stadial have often not been preserved within outwash plains at its maximum extent line. The Main Stadial ice sheet advanced onto the middle basin of the Biebrza Valley. In the Suwałki Lakeland it only reached the northern end of the Augustów Plain, and farther south it released a small ice tongue which deposited a thin till in the Augustów Plain. A similar situation might have occurred in the Mazury-Kurpie, Dainava (Lithuania) and Porechje, Aziory and Bershty (Belarus) outwash fans. Glacial sediments of the Main Stadial have commonly not been preserved in the area of its maximum limit.

5. “Till islands”, emerging from beneath the afore-mentioned outwash fans, as well as the “islands” between the Augustów Plain and Biebrza Valley, are composed of the *wiecie* Stadial tills, at least as regards their upper parts.

Lithuanian and Belarussian scientists claim that during the Middle Stadial of the Vistulian Glaciation there was no ice sheet in the territories of their countries. Nevertheless, these studies indicate that the Middle Stadial till is represented probably by the highest till observed in southwestern Lithuania and northwestern Belarus, in the area between the limits of the Middle Stadial and the Upper Stadial. The Gruda Stadial till of southwestern Lithuania, and the Ozerie Phase (of the Baltijskij horizon, Orshanskij IV Stadial) till of northwestern Belarus, probably correspond to the *wiecie* Stadial till in Poland. On the basis of lithological sections of the Eemian Interglacial and the last glaciation deposits in the Neman River valley, the ice sheet limit of the *wiecie* Stadial of the Vistulian Glaciation is postulated to run 15 to 20 km to the south of its previously drawn location.

## REFERENCES

- ANDRZEJESZCZAK B. (1971) — New locality in the Eemian Interglacial in NE Poland (in Polish with English summary). *Prz. Geogr.*, **43** (1–2): 587–596.
- ASTAPOVA S. D., SANKO A. F., ZIMENKOV O. L., LOGINOVA L. P., NAZAROV V. L., RYLOVA T. B., KHURSEVICH G. K., YAKUBOVSKAYA T. V., SHIMANOVICH S. L. and ZUBOVICH S. F. (1987) — Stratigraphic aspects of studying glacial deposits of Belarus. In: *Quaternary Glaciations in the Northern Hemisphere. Results of Investigation on the Territory of Belarus: 7–24*. Minsk.
- BAŁUK A. and KRUPI SKI K. M. (2000) — Eemian Interglacial deposits at Rakowo Nowe near Kolno (in Polish with English summary). *Prz. Geol.*, **48** (9): 789–795.
- BANASZUK H. (1980) — Geomorfologia południowej części Kotliny Biebrza skiej (in Polish with English summary). *Pr. Stud. Geogr. UW*, **2**: 7–69.
- BANASZUK H. (1995) — Geneza i rozwój rze by Puszczy Knyszy skiej w wietle analizy geomorfologicznej i bada termoluminescencyjnych. In: *Puszcza Knyszy ska. Monografia Przyrodnicza* (ed. A. Czerwi ski): 33–48.
- BANASZUK H. (1996) — Paleogeografia. Naturalne i antropogeniczne przekształcenia doliny górnej Narwi (in Polish with English summary). *Wyd. Ekonomia i rodowisko*. Białystok.
- BANASZUK H. (1998) — Ranges and course of the Wisła and Warta Glaciations in North-Eastern Poland in the latest studies. *Mat. IV Zjazdu Geomorfologów Polskich* (ed. K. P. kala): 233–240. UMCS. Lublin.
- BANASZUK H., STA SKA-PRÓSZY SKA W. and PRÓSZY SKI M. (1994) — On the paleogeography of the Wisła (Vistula) Glaciation in the NE Poland and outflow of glacial fluvial waters by Biebrza Pradolina in the light of termoluminescence investigation (in Polish with English summary). *Prz. Geogr.*, **66** (1–2): 57–69.
- BEHRE K. E. (1989) — Biostratigraphy of the last glacial period in Europe. *Quatern. Sc. Rev.*, **8**.
- BER A. (1972a) — Mapa Geologiczna Polski 1:200 000, ark. Sokółka, wyd. A (Mapa utworów powierzchniowych). *Inst. Geol. Warszawa*.
- BER A. (1972b) — Obją nienia do Mapy Geologicznej Polski 1:200 000, ark. Sejny i Sokółka. *Inst. Geol. Warszawa*.
- BER A. (1973) — Geological setting of Eemian Interglacial localities in Suwałki Lake District (in Polish with English summary). *Prz. Geol.*, **21** (7): 363–366.
- BER A. (1974) — The Quaternary of the Suwałki Lake District (in Polish with English summary). *Biul. Inst. Geol.*, **269**: 23–94.
- BER A. (1975a) — Mapa Geologiczna Polski 1:200 000, ark. Elk, wyd. A (Mapa utworów powierzchniowych). *Inst. Geol. Warszawa*.
- BER A. (1975b) — Obją nienia do Mapy Geologicznej Polski 1:200 000, ark. Elk. *Inst. Geol. Warszawa*.
- BER A. (1982) — Marginal zones and deglaciation during the North-Polish Glaciation in the Suwałki-Augustów Lakeland. *Biul. Inst. Geol.*, **343**: 71–89.
- BER A. (1990) — Obją nienia do Szczegółowej Mapy Geologicznej Polski 1:50 000, ark. Suwałki. *Pa stw. Inst. Geol. Warszawa*.
- BER A. (1999) — Glacitectonic of the Suwałki-Augustów Lakeland in connection to neotectonic movements and tectonic structures of the crystalline basement (NE Poland) (in Polish with English summary). *Prz. Geol.*, **47** (9): 831–839.
- BER A. (2000) — Pleistocene of north-eastern Poland and neighbouring areas against crystalline and sedimentary basement (in Polish with English summary). *Pr. Pa stw. Inst. Geol.*, **170**.
- BER A. (in preparation) — Szczegółowa Mapa Geologiczna Polski 1:50 000, ark. Wo na Wie . *Pa stw. Inst. Geol. Warszawa*.
- BITNER K. (1956) — Three new localities of Pleistocene flora (in Polish with English summary). *Biul. Inst. Geol.*, **100**: 247–262.
- BITNER K. (1957) — Three localities of interglacial flora in the vicinity of Sidra (Northly of Sokółka in Podlasie) (in Polish with English summary). *Biul. Inst. Geol.*, **118**: 109–154.
- BLAZHAUSKAS N., KISYELYENE D., KUCHINSKAITE V., STANCHIKAITE M., SHEIRYENE V. and SHINKUNAS P. (1998) — Late Glacial and Holocene sedimentary environment in the region of the Ula River. *Geologija*, **25**: 20–30.
- BOGACKI M. (1976) — Współczesne sandry na przedpolu Skeidar-arjökull (Islandia) i plejstoce skie sandry w Polsce północno-wschodniej. *Rozpr. UW*, **93**: 1–167.
- BORÓWKO-DEU AKOWA Z. (1971) — Fossil flora from Eemian Interglacial at Smolniki near Suwałki (in Polish with English summary). *Prz. Geogr.*, **43** (1–2): 591–600.
- BORÓWKO-DEU AKOWA Z. and HALICKI B. (1957) — Interglacial sections of the Suwałki region and of the adjacent territory (in Polish with English summary). *Acta Geol. Pol.*, **7** (4): 361–401.
- BREMÓWNA M. and SOBOLEWSKA M. (1950) — The results of botanical investigations of interglacial deposits in the Niemen basin (in Polish with English summary). *Acta Geol. Pol.*, **1** (4): 335–361.
- BU CZY SKI M. (1960) — Interglacial site at Czarna Wie (in Polish with English summary). *Prz. Geol.*, **8** (9): 483–485.
- CHEPULITE W. A. (1966) — O pyerekrytii riss-wyurmsskikh otlozheniy v okresnostyakh g. Grodno morenoy poslednego oledenyeniya. In: *Palinologiya v Geologicheskikh Issledovaniyakh Pribaltiki*. Zinatne. Riga.
- CHEPULITE W. A. (1978) — K voprosu o pyerekrytii i niepokrytii morenym suglinkom poslednego oledenyeniya miezhmorenykh i miezhlednikovykh otlozheniy yugo-vostocnoy Litwy. In: *Dostizheniya i Perspektivy Gyeologicheskogo Izucheniya Litovskoy SSR*. Vilnius.
- CHEPULITE W. A. (1986) — K diskusyi o ploshchadi rasprastranieniya morenogo suglinka poslednego oledenyeniya v yuzhnoy Pribaltike. In: *Issledovaniya Lednikovykh Obrazovaniy Pribaltiki*. Vilnius.
- CZAPLICKA J. (1952) — Opracowanie geologiczne 13 profili badawczych z utworami interglacjalnymi z okolic Sokółki na Podlasiu. *Centr. Arch. Geol. Pa stw. Inst. Geol. Warszawa*.
- DOROFIEYEV P. I. (1963) — Nowyye dannyye o pleystocenovykh florakh. In: *Materialy po Istorii Flory i Rastitelnosti SSSR*. Wyp. 4. (ed. W. N. Sukachev): 20–31. *Akad. Nauk SSSR. Moskva-Leningrad*.
- DROZDOWSKI E. (1986) — Stratigraphy and origin of Vistulian Glaciation in northern part of the Lower Vistula Region (in Polish with English summary). *Pr. Geogr., Inst. Geogr. Przestrz. Zagosp. PAN*, **146**: 1–90.
- FEDOROWICZ S., LASKOWSKI K. and LINDNER L. (1995) — O mo liwo ci dalszego zasi gu l dolodu zlodowacenia wisly w wietle datowa TL osadów lodowcowych w północnej części Wysoczyzny Białostockiej. *Prz. Geol.*, **43** (11): 941–944.
- GAIGALAS A. (1995) — Glacial history of Lithuania. In: *Glacial Deposits in North-East Europe* (eds. J. Ehlers, S. Kozarski and P. Gibbard): 127–135. *Balkema. Rotterdam*.
- GAIGALAS A. and YARTSEV W. I. (1992) — Metodicheskaya osnova izucheniya chetyvertichntch otlozheniy. *Viln. Uniw. Vilnius*.
- GALON R. (1937) — Geologia i morfologia Prus Wschodnich. In: *Słownik geograficzny Pa stwa Polskiego* **1**: 30–42. *Warszawa*.
- GALON R. and ROSZKÓWNA L. (1961) — Extents of the Scandinavian Glaciations and of their recession stages on the territory of Poland in the light of an analysis of the marginal forms on inland ice. *Prz. Geogr.*, **33** (3): 347–364.
- GAŁ ZKAD., KUSI SKI J. T. J. and MARKS L. (1998) — Próba rewizji zasi gu l dolodu zlodowacenia wisly w południowo-zachodniej części Mazur. *Mat. V Konferencji „Stratygrafia plejstocenu Polski”* (eds. S. Lisicki *et al.*): 21–23. *Iznota*.
- GORETSKIY G. I. (1980) — Osobyenosti paleopotamologii lednikovykh oblastiye. *Nauka i Tekhnika*: 1–281. *Minsk*.
- GORETSKIY G. I. (1983) — Gyeologicheskaya Karta Chetyvertichnykh Otlozhniy Bieloruskoy SSR 1:500 000. *M.G. SSSR. Moskva*.
- GUOBYTE R. and PAVLOVSKAYA I. E. (1998) — The stadial limits and maximum extent of the Weichselian Glaciation in North-Eastern Lithuania and Nord-Western Belarus. *Mat. Field Symposium on Glacial Processes and Quaternary Environment in Latvia*: 23–24. *Riga*.
- HALICKI B. (1950) — Some problems concerning the stratigraphy of the Pleistocene of the European Lowland (in Polish with English summary). *Acta Geol. Pol.*, **1** (2): 106–142.



- HALICKI B. (1951) — Principal sections of the Pleistocene in the Niemen basin (in Polish with English summary). *Acta Geol. Pol.*, **2** (1–2): 5–101.
- INSTRUKCJA W SPRAWIE OPRACOWANIA I WYDANIA SZCZEGÓŁOWEJ MAPY GEOLOGICZNEJ POLSKI W SKALI 1:50 000 (1991) — Pa. stw. Inst. Geol. Warszawa.
- JANCZYK-KOPIKOWA Z. (1958) — Wyniki analiz palinologicznych z miejscowo ci Białousy, Białosuknia, Mystki, Jezioro i Horodyszcze. *Centr. Arch. Geol. Pa. stw. Inst. Geol. Warszawa.*
- JANCZYK-KOPIKOWA Z. (1998) — Ekspertyza (orzeczenie) dotycząca próbek z otworu Lipińskie, ark. Biała Piska Szczegółowej Mapy Geologicznej Polski 1:50 000. *Centr. Arch. Geol. Pa. stw. Inst. Geol. Warszawa.*
- KACPRZAK L. (2000) — Szczegółowa Mapa Geologiczna Polski w skali 1:50 000, ark. Sztabin. *Centr. Arch. Geol. Pa. stw. Inst. Geol. Warszawa.*
- KALNIET A. and KARASZEWSKA U. (1972) — Mapa Geomorfologiczna Polski 1:500 000. *PPWK. Warszawa.*
- KARABANOV A. K. (1987) — Grodnenskaya vozvyshehnoshch. Stroyeniye, relyef i etapy formirovaniya. *Nauka i Tekhnika: 1–107. Minsk.*
- KENIG K. (1998) — Petrographic foundations of till stratigraphy of north-eastern Poland (in Polish with English summary). *Biul. Pa. stw. Inst. Geol.*, **380**.
- KOCISZEWSKA-MUSIAŁ G. (1978) — Quaternary aggregates of the Suwałki Region against its geological background (in Polish with English summary). *Pr. Muz. Ziemi*, **29**: 3–79.
- KOCISZEWSKA-MUSIAŁ G. and LECIEJEWICZ K. (1982) — Geological situation of peat from Eemian interglacial period in Blaskowizna (in Polish with English summary). *Pr. Muz. Ziemi*, **35**: 69–79.
- KONDRACKI J. (1952) — Remarks about morphological evolution of the Mazurian Lake District (in Polish with English summary). *Biul. Pa. stw. Inst. Geol.*, **65**: 513–597.
- KONDRATIENE O. (1996) — Stratigrafiya i paleogeografiya kvartera Litvy po paleobotanicheskim dannym. *Academia. Vilnius.*
- KOZARSKI S. (1977) — Ukierunkowane wytopiska na sandrach. *Stud. Geol. Pol.*, **52**: 235–248.
- KOZŁOWSKA M. and KOZŁOWSKI I. (1993) — Objasnienia do Szczegółowej Mapy Geologicznej Polski 1:50 000, ark. Pisz. *Pa. stw. Inst. Geol. Warszawa.*
- KOZŁOWSKA M. and KOZŁOWSKI I. (1995a) — Szczegółowa Mapa Geologiczna Polski 1:50 000, ark. Pisz. *Pa. stw. Inst. Geol. Warszawa.*
- KOZŁOWSKA M. and KOZŁOWSKI I. (1995b) — Objasnienia do Szczegółowej Mapy Geologicznej Polski 1:50 000, ark. Myszyniec. *Centr. Arch. Geol. Pa. stw. Inst. Geol. Warszawa.*
- KOZŁOWSKA M. and KOZŁOWSKI I. (1996a) — Objasnienia do Szczegółowej Mapy Geologicznej Polski 1:50 000, ark. Ruciane. *Pa. stw. Inst. Geol. Warszawa.*
- KOZŁOWSKA M. and KOZŁOWSKI I. (1996b) — Objasnienia do Szczegółowej Mapy Geologicznej Polski 1:50 000, ark. Łyse. *Pa. stw. Inst. Geol. Warszawa.*
- KOZŁOWSKA M. and KOZŁOWSKI I. (1997) — Objasnienia do Szczegółowej Mapy Geologicznej Polski w skali 1:50 000, ark. Lipowiec. *Centr. Arch. Geol. Pa. stw. Inst. Geol. Warszawa.*
- KOZŁOWSKA M. and KOZŁOWSKI I. (1999) — Szczegółowa Mapa Geologiczna Polski 1:50 000, ark. Biała Piska. *Centr. Arch. Geol. Pa. stw. Inst. Geol. Warszawa.*
- KRYGER M. I., KUPTSOVA I. A. and KURYERAVA L. W. (1971) — Adorvyyen mizhledavikovykh dyatamitav Paniemu –2 lya g. Grodna. In: *Antrapagen Belarusi. Minsk.*
- KRZYWICKI T. (1993) — Czwartorzęd Pojezierza Suwalskiego. *Przewodnik 64 Zjazdu Pol. Tow. Geol.*, Suwałki: 59–81. *Pa. stw. Inst. Geol. Warszawa.*
- KRZYWICKI T. (1996) — Szczegółowa Mapa Geologiczna Polski 1:50 000, ark. Stacja Augustów. *Centr. Arch. Geol. Pa. stw. Inst. Geol. Warszawa.*
- KRZYWICKI T. (1997a) — Mapa rozprzestrzenienia młodoplejstocenich glin zwałowych w południowej części Puszczy Augustowskiej jako podstawa wyznaczenia zasięgu ostatniego zlodowacenia w północno-wschodniej Polsce. *Centr. Arch. Geol. Pa. stw. Inst. Geol. Warszawa.*
- KRZYWICKI T. (1997b) — Proba pokazania maksymalnego radyusa deistviya oledneniya Wisły (Weichselian) v severo-vostocznoy Polshe. *Mat. Konferencji “Quaternary Deposits and Neotectonics in the Area of Pleistocene Glaciations”*: 31–32. *Minsk.*
- KRZYWICKI T. (1999a) — Szczegółowa Mapa Geologiczna Polski 1:50 000, ark. Rajgród. *Centr. Arch. Geol. Pa. stw. Inst. Geol. Warszawa.*
- KRZYWICKI T. (1999b) — Maksymalny zasięg lodu zlodowacenia wisły w północno-wschodniej Polsce i obszarach przyległych. *Centr. Arch. Geol. Pa. stw. Inst. Geol. Warszawa.*
- KRZYWICKI T. (2001a) — Szczegółowa Mapa Geologiczna Polski 1:50 000, ark. Rygól. *Pa. stw. Inst. Geol. Warszawa.*
- KRZYWICKI T. (2001b) — Objasnienia do Szczegółowej Mapy Geologicznej Polski 1:50 000, ark. Rygól, Rudawka. *Pa. stw. Inst. Geol. Warszawa.*
- KUPRYJANOWICZ M. (1991) — Eemian, early and late Vistulian, and Holocene vegetation in the region of Machnacz peat-bog near Białystok (NE Poland). *Acta Palaeobot.*, **31** (1–2): 215–225.
- KUPRYJANOWICZ M. (1994) — Zmiany roślinności torfowisk Machnacz w Puszczy Knyszyńskiej w okresie interglacjału eemskiego, vistulianu i holocenu. *Ph. D. thesis. Arch. Uniwersytetu w Białymstoku. Białystok.*
- LICHWA M. (2000) — Szczegółowa Mapa Geologiczna Polski 1:50 000, ark. Radziłów. *Centr. Arch. Geol. Pa. stw. Inst. Geol. Warszawa.*
- LINDNER L. (1987) — Main stratigraphic problems in the Pleistocene of Poland. *Bull. Pol. Acad. Sc. Earth Sc.*, **35** (4): 343–358.
- LINDNER L. ed. (1992) — Czwartorzęd. Osady, metody badań, stratygrafia. *PAE. Warszawa.*
- LISICKI S. (1993) — Deglacjacja Pojezierza Suwalskiego w okresie schyłku plejstocenu. *Przewodnik 64 Zjazdu Pol. Tow. Geol.*: 81–86. *Pa. stw. Inst. Geol. Warszawa.*
- LISICKI S. (1996) — Plejstocen Pojezierza Mrągowskiego. *Centr. Arch. Geol. Pa. stw. Inst. Geol. Warszawa.*
- LISICKI S. (1997) — Pleistocene of the Mrągów Lakeland. *Geol. Quart.*, **41** (3): 327–346.
- LISICKI S. (1998) — Attempt of lithostratigraphic correlation of tills in northeastern Poland and southern Lithuania. *Geol. Quart.*, **42** (2): 161–172.
- LISZKOWSKI J. (1993) — The effects of Pleistocene ice-sheet loading-unloading cycles on the bedrock structure of Poland. *Folia Quatern.*, **64**: 7–23.
- MAJDANOWSKI S. (1947) — Distribution, density and direction of lakechannels of the Polish Lowlands (in Polish with English summary). *Prz. Geogr.*, **21** (1–2): 38–69.
- MAKOWSKA A. (1976) — Stratigraphy of tills exposed along the valley of the Lower Vistula area. *Zesz. Nauk. UAM, Geografia*, **12**: 239–242.
- MAKOWSKA A. (1979) — Eemian Interglacial in the Lower Vistula Region (in Polish with English summary). *Stud. Geol. Pol.*, **63**: 1–90.
- MAKOWSKA A. (1980) — Late Eemian with preglacial and glacial part of Vistulian glaciation in the Lower Vistula Region. *Quater. Stud. Pol.*, **2**.
- MAKOWSKA A. (1986) — Morza plejstoceniczne w Polsce — osady, wiek i paleogeografia. *Pr. Inst. Geol.*, **120**.
- MAKOWSKA A. (1994) — Climatic variation in the pre-glacial part of the Toruń Glaciation in the Lower Vistula Region and the Elbląg Elevation (preliminary report). *Geol. Quart.*, **38** (1): 133–154
- MAKOWSKA A. (1999) — Quaternary and its substratum in the Elbląg Elevation as seen in the aspect of the Pleistocene Baltic Sea (in Polish with English summary). *Biul. Pa. stw. Inst. Geol.*, **386**: 59–132.
- MAKSIAK S. (1992) — Szczegółowa Mapa Geologiczna Polski 1:50 000, ark. Kolno. *Centr. Arch. Geol. Pa. stw. Inst. Geol. Warszawa.*
- MAMAKOWA K. (1988) — Pollen stratigraphy of the Eemian and adjoining glacial deposits based on continuous sequences in Poland. *Bull. Pol. Acad. Sci. Earth. Sci.*, **36** (3–4): 299–307.
- MANGERUD J. (1992) — The last interglacial/glacial cycle in Northern Europe. In: *Quaternary Landscapes* (eds. C. K. Linda *et al.*). *Univ. Minn. Press, Minneapolis. Meeting: Quaternary Landscapes, 1991. Minneapolis MN.*
- MARCINIAK B. and KOWALSKI W. W. (1978) — Dominant diatoms, pollen, chemistry and mineralogy of the Eemian lacustrine sediments from Nidzica (Northern Poland), a preliminary report. *Pol. Arch. Hydrobiol.*, **25** (1–2).

- MARCINKIEWICZ A. (1973) — Range of Young Glacial morphology in NE Poland in cartographical aspect (in Polish with English summary). *Pol. Prz. Kartogr.*, **5** (4): 151–154.
- MARKS L. (1988) — Relation of substrate to the Quaternary paleorelief and sediments, western Mazury and Warmia (Northern Poland). *Zesz. Nauk. AGH 1165. Geologia*, **14** (1): 1–76.
- MARKS L. (1990a) — Middle and Late Pleistocene glacial episodes in western Mazury and Warmia, northern Poland. *Lundqua Report*, **32**.
- MARKS L. (1990b) — Outline of geology and evolution of the northwestern Kurpie outwash, Northern Poland (in Polish with English summary). *Kwart. Geol.*, **34** (3): 489–502.
- MARKS L. (1991) — Zasięgi i dolodów zlodowacenia Wisły w rodkowej i wschodniej Polsce. *Ser. Geogr., UAM*, **50**: 531–538.
- MARKS L. (1994) — Dead-ice features at the maximum extent of the last glaciation in the northeastern Poland: 77–83. *Z. Geomorph. NF Suppl.-Bd. 95. Berlin. Stuttgart*.
- MARKS L. (1997a) — Middle and late Vistulian Glaciation in Poland. Abstract. *Mat. INQUA-SEQS Symposium “The Late Pleistocene in Eastern Europe — stratigraphy and climate”* (ed. J. Satkunas): 36. Vilnius.
- MARKS L. (1997b) — Zasięgi zlodowacenia Wisły na N i u Polskim. *Mat. IV Konferencji „Stratygrafia plejstocenu Polski”* (eds. D. Krzyszkowski and B. Przybylski). Kamieniec Zbkowicki.
- MOJSKI J. E. (1972) — Nizina Podlaska. In: *Geomorfologia Polski* (ed. R. Galon), **2**: 318–361. PWN. Warszawa.
- MOJSKI J. E. (1979) — Main stratigraphic glacial units of the Vistulian in the Polish Lowland. In: *Quaternary Glaciations in the Northern Hemisphere. Report*, **5**. Prague.
- MOJSKI J. E. (1988) — O zmianach klimatu podczas czwartorzędów. In: *Przemiany rodowiska geograficznego Polski* (ed. L. Starkel). Ossolineum. Wrocław.
- MOJSKI J. E. (1993) — Europa w plejstocenie. *Ewolucja rodowiska przyrodniczego*. Wyd. PAE. Warszawa.
- MOJSKI J. E. (1997) — Small stratigraphic units of Vistulian age in the southern Baltic Area. In: *Mat. INQUA-SEQS Symposium “The Late Pleistocene in Eastern Europe — stratigraphy and climate”* (ed. J. Satkunas): 39–40. Vilnius.
- MOJSKI J. E. and RÜHLE E. (1965) — Atlas geologiczny Polski w skali 1:3 000 000. *Zagadnienia stratygraficzno-facjalne*. Państw. Inst. Geol. Warszawa.
- MORAWSKI W., KRUPSKI K. and WINTER H. (1998) — Nowe stanowiska interglacjału eemskiego na Wysoczyźnie Mławskiej w rejonie Nidzicy. *Mat. V Konferencji „Stratygrafia plejstocenu Polski”* (eds. S. Lisicki *et al.*): 88–89. Iznota.
- MORAWSKI W., KRUPSKI K. and WINTER H. (1999) — Biogenic sediments of the Eemian Interglacial on the Nidzica Plateau, southwestern Mazury Lakeland. *Geol. Quart.*, **43** (1): 79–84.
- MUSIAŁ A. (1984) — Genesis of zone of highest elevations of the Kolno Upland (in Polish with English summary). *Pr. Stud. Geogr.* **5**: 113–153.
- MUSIAŁ A., STRASZEWSKA K. and ZIEMBSKA-TWORZYDŁO M. (1982) — Interglacial lacustrine reservoir at Niewodowo, Kolno Upland (in Polish with English summary). *Kwart. Geol.*, **26** (1): 159–176.
- PACHUCKI C. (1961) — Les moraines terminales de la dernière glaciation sur le territoire de Peribalticum (in Polish with French summary). *Rocz. Pol. Tow. Geol.*, **31** (2–4): 303–318.
- PASYUKIEVICH V. I., NELIPOVICH A. N., LEVITSKAYA R. I. and SEMENYUK A. D. (1972) — Krayevyye obrazovaniya maksimalnoy (ozyerskoy) stadii Wałdajskogo oledenieniya zapada Bielorusii. In: *Krayevyye Obrazovaniya Materikovykh Oledenieniy*: 265–268. Moskva.
- PAVLOVSKAYA I. (1998) — Muravian (Eemian) deposits as an indicator of the interglacial hydrographic network of Belarus. *Geologija*, **25**: 70–76. Akademia. Vilnius.
- PAWŁOWSKI S. (1933) — La glaciation nordique en Pologne d’après les recherches nouvelles. *Comptes Rendus du III Congrès de Geogr.* 1930. Beograd.
- PIETKIEWICZ S. (1928) — Esquisse morphologique de la partie occidentale du district de Suwałki (in Polish with French summary). *Prz. Geogr.*, **8**: 168–222.
- ROSZKO L. (1968) — Recession of last inland ice from Poland’s territory (in Polish with English summary). In: *Last Scandinavian glaciation in Poland*. *Pr. Geogr. Inst. Geogr. PAN*, **74**.
- RÓYCKI S. Z. (1972a) — Plejstocen Polski rodkowej. PWN. Warszawa.
- RÓYCKI S. Z. (1972b) — Nizina Mazowiecka. In: *Geomorfologia Polski* (ed. R. Galon), **2**: 271–317. PWN. Warszawa.
- RÓYCKI S. Z. (1978) — From Mochty to a synthesis of the Polish Prepleistocene (in Polish with English summary). *Rocz. Pol. Tow. Geol.*, **47** (3–4).
- RÜHLE E. (1965) — Czwartorzęd Polski. In: *Zarys geologii Polski*: 299–347. PWN. Warszawa.
- RÜHLE E. and SOKOŁOWSKA M. (1961) — Mapa utworów czwartorzędowych Polski 1:1 000 000. *Inst. Geol. na VI Kongres INQUA*. Warszawa.
- SATKUNAS J. (1997a) — Stadials and interstadials of the late Weichselian in Lithuania, a climatostratigraphical aspect. In: *Mat. INQUA-SEQS Symposium “The Late Pleistocene in Eastern Europe — stratigraphy, paleoenvironment and climate”* (ed. J. Satkunas): 51–52. Vilnius.
- SATKUNAS J. (1997b) — Outline of Quaternary stratigraphy of Lithuania. In: *Mat. INQUA-SEQS Symposium “The Late Pleistocene in Eastern Europe — stratigraphy, paleoenvironment and climate”* (ed. J. Satkunas): 65–68. Vilnius.
- SATKUNAS J. and GRIGYENE A. (1997) — The Jonionys site-sequence of the Eemian Interglacial and the Weichselian Interstadials. In: *Mat. INQUA-SEQS Symposium “The Late Pleistocene in Eastern Europe — stratigraphy, paleoenvironment and climate”* (ed. J. Satkunas): 77–81. Vilnius.
- SATKUNAS J., GRIGYENE A., GUOBYTE R., MARCHINKEVICHYUS V. and RACHKAUSKAS V. (1997) — Upper stratigraphy based on date from Smalvas and Dysnai localities, Northeastern Lithuania. *Geologija*, **22**: 26–35.
- SATKUNAS J., GRIGYENE A. and ROBERTSSON A.-M. (1998) — An Eemian-Middle Weichselian sequence from the Jonionys site, southern Lithuania. *Geologija*, **25**: 82–91.
- SATKUNAS J. and HÜTT G. (1999) — Stratigraphy of the section Antaviliai, eastern Lithuania, and its implication for the Upper Weichselian climatostratigraphic subdivision. *Geol. Quart.*, **43** (2): 213–218.
- SHACKLETON N. J. and OPDYKE N. D. (1973) — Oxygen isotope and paleomagnetic stratigraphy of equatorial Pacific core V28238: oxygen isotope temperatures and ice volumes on a 105 and 106 scale. *Quatern. Rec.*, **3** (1).
- SŁOWSKI W. (1971) — Objawy do Mapy Geologicznej Polski 1:200 000. *ark. Pisz. Inst. Geol. Warszawa*.
- STRASZEWSKA K. (1975) — Middle Polish (Riss) age of marginal forms in the vicinity of Łomża (north-eastern Poland). *Bull. Acad. Pol. Sc. Sér. Sc. Terre*, **23** (2): 111–118.
- SZAFER W. (1953) — Pleistocene stratigraphy of Poland from the floristical point of view (in Polish with English summary). *Rocz. Pol. Tow. Geol.*, **22** (1).
- SZAFER W. (1955) — Progress in the investigation of Quaternary floras in Poland (in Polish with English summary). *Biul. Inst. Geol.* **70**: 55–64.
- RODO A. (1950) — Rozwój roślinności pod Grodnem w czasie ostatniego interglacjału. *Acta Geol. Pol.*, **1** (4): 365–390.
- VAYTEKUNAS P. (1968) — Degradacja poslednego skandinavskogo lednikovogo pokrova na territorii Pribaltiki. *Lietuvos TSR Aukštųjų Mokyklų Mokslų Darbai. Geografija ir Geologija*, **5**: 123–135.
- VALTCHIK M. A., ZUS M. E., FEDENYA W. M. and KARABANOV A. K. (1990) — Krayevyye obrazovaniya Belorusskoy Gryady. *Nauka i Tekhnika*. Minsk.
- VOZNYATCZUK L. N. (1956) — O položenii granitsy poslednego oledenieniya w Belorusii. *Učen. Zap. Belorus. Gos. Un-ta. Ser. Geol.*, **28**.
- VOZNYATCZUK L. N. (1960) — Kratkiy ocherk stratigrafii antropogenovykh otlozheniy zapadnoy chasti Grodnenskoy oblasti. In: *Mat. konf. molodykh uczenykh AN BSSR (Jeststvennyye nauki)* Akad. Nauk. BSSR. Minsk.



- VOZNYATCHUK L. N. (1972) — Vozrost maksimalnoy stadii valdayskogo oledenieniya na siviero-zapade SSSR i osnovnyye fazy degradatsii lednika. In: *Voprosy Chetvertichnoy Geologii*, **6**. Riga.
- VOZNYATCHUK L. N. (1976) — Niekotoryye zakonomiernosti izmieniyy prirodnoy obstanovki Belorussii w Neopleystocene. In: *Voprosy Geologii Belorussii*. Minsk.
- VOZNYATCHUK L. N., KUZNIETSOV V. A. and MANDER E. R. (1972) — Nowyye dannyye o stroyeni i vierkhnicy valdayskoy terrasy Niemana. *Dok. Akad. Nauk. BSSR.*, **16** (5).
- VOZNYATCHUK L. N. and VALTCHIK M. A. (1978) — Morfologiya, stroyeniye i istoria razvitya doliny Niemana w neopleystocene i golocene. *Nauka i Tekhnika*. Minsk.
- WI CKOWSKI K. (1966) — Bottom deposits of Lake Mikołajki (in Polish with English summary). *Pr. Geogr. Inst. Geogr. PAN*, **57**.
- WOLDSTEDT P. (1929) — Das Eiszeitalter. Grundlinien einer Geologie des Diluviums: 78–91, 165–169. Stuttgart.
- WOLDSTEDT P. (1935) — Geologisch-morphologische Übersichtskarte des norddeutschen Vereisungsgebietes 1:1 500 000. Preuss. Geol. Berlin.
- WOŁK-MUSIAŁ E. (1980) — Margin of the Baltic Glaciation in the area of Kolno Plateau in the light of studies of morphologic elements (in Polish with English summary). *Pr. Stud. Geogr.*, **2**: 71–100.
- WOŁOSOWICZ S. (1926) — La moraine de forn de la transgression de Wigry et sou rôle dans la structure géologique du “pays lacustre” de Suwałki (in Polish with French summary). *Spraw. Pa stw. Inst. Geol.*, **3** (3–4): 434–467.
- WOŁOSOWICZ S. (1928) — Utwory dyluwialne i morfologia Pojezierza Suwalskiego. *Posiedz. Nauk. Pa stw. Inst. Geol.*, **21**: 50–52.
- WYSOCZA SKI-MINKOWICZ T. (1982) — Climatochronostratigraphic subdivision of the last cool period (Vistulian). *Quater. Stud.*, **3**.
- YELOVICHEVA Ya. K. (1997) — Stratigraphy, paleoenvironment and climate of the late Pleistocene in Belarus. *Mat. INQUA-SEQS Symposium “The Late Pleistocene in Eastern Europe — stratigraphy, paleoenvironment and climate”* (ed. J. Satkunas): 62. Vilnius.
- YELOVICHEVA Ya. K. and SANKO A. (1999) — Palynostratigraphy of the Poozerie Glaciation (Vistulian) in Belarus. *Geol. Quart.*, **43** (2): 203–212.
- YELOVICHEVA Ya. K., SANKO A. F. and KALECZYC E. G. (1996) — Klimatochronologiczny i archeologiczny podział pó nego plejstocenu i holocenu Białorusi. *Mat. białorusko-polskiego seminarium “Przyrodnicze procesy w plejstocene i holocene na Białorusi i w Polsce”*. Mi sk.
- UK R. (1997) — Objasnienia do Szczegółowej Mapy Geologicznej Polski 1:50 000, ark. w i tajno. *Centr. Arch. Geol. Pa stw. Inst. Geol. Warszawa*.
- UK R. (1998a) — Szczegółowa Mapa Geologiczna Polski 1:50 000, ark. Spychowo. *Centr. Arch. Geol. Pa stw. Inst. Geol. Warszawa*.
- UK R. (1998b) — Objasnienia do Szczegółowej Mapy Geologicznej Polski 1:50 000, ark. Spychowo. *Centr. Arch. Geol. Pa stw. Inst. Geol. Warszawa*.
- UREK S. (1969) — The peat deposits of Grajewo county with geomorphological conditions as background (in Polish with English summary). *Prz. Geogr.*, **41** (3): 469–484.
- UREK S. (1975) — Genesis of bog formation in the Biebrza Urstromtal (in Polish with English summary). *Pr. Geogr. Inst. Geogr. Przestrz. Zagosp. PAN*, **110**: 109.
- UREK S. (1984) — Relief, geologic structure and hydrography of the Biebrza ice marginal valley. *Pol. Ecol. Stud.*, **10** (3–4): 239–251.
- UREK S. (1990) — Zwi zek procesu zatorfienia z elementami rodowiska przyrodniczego wschodniej Polski. *Rocz. Nauk Roln. PAN, Seria D*, **220**.
- UREK S. (1992) — Wyjasnienie sytuacji chronostratygraficznej i geomorfologicznej w okolicy torfowiska Machnac. *Spraw. z Bad. Nauk. Kom. Bad. Czwart. PAN*, **9**: 39–50.