

Microorganisms in selected sulphuric springs of the Polish Carpathians

Lucyna RAJCHEL, Jacek RAJCHEL and Konrad WOŁOWSKI



Rajchel L., Rajchel J. and Wołowski K. (2002) — Microorganisms in selected sulphuric springs of the Polish Carpathians. *Geol. Quart.*, 46 (2): 189–198. Warszawa.

Microorganisms colonising sulphurous waters were found at the bottoms of the spring niches and along spring outflows. Five springs from the Carpathians and two from the Carpathian Foredeep were selected for investigations. Sulphurous flora is represented mainly by sulphuric bacteria. They occur as individual threads, spider-webs, festoons, encrustations and covers. Their colours may be white, creamy, violet, pink and purple. Altogether 31 microorganism taxa: bacteria (16), cyanoprokaryota (3) and algae (12) were identified and illustrated. The bacterial occurrences have been correlated with physico-chemical state of the waters.

Lucyna Rajchel, Department of Stratigraphy and Regional Geology, University of Mining and Metallurgy, Mickiewicza 30, 30-059 Kraków, Poland, e-mail: rajchel@geolog.geol.agh.edu.pl; Jacek Rajchel, Department of General and Mathematical Geology, University of Mining and Metallurgy, Mickiewicza 30, 30-059 Kraków, Poland, e-mail: jrajchel@geolog.geol.agh.edu.pl; Konrad Wołowski, Department of Phycology, W. Szafer Institute of Botany, Polish Academy of Sciences, Lubicz 46, 31-512 Kraków, Poland, e-mail: wolowski@ib-pan.krakow.pl (received: August 7, 2001; accepted: March 6, 2002).

Key words: southern Poland, Carpathian Foredeep, sulphuric springs, microorganisms, schizomycetes, cyanoprokaryota, algae.

INTRODUCTION

Bacteria participate in processes associated with a migration of many elements in natural environment, including sulphur, manganese, nitrogen and iodine, a significant part of this migration taking place in groundwater. Particular taxa develop in groundwater under specific conditions determined by thermal regime, redox potential, pH and mineralisation; some require nutritional energy provided by organic matter (Macioszczyk, 1987).

Sulphurous waters are a typical environment of sulphur-fixing bacteria (Strzeszewski, 1913; Topińska-Luchter, 1951; Jarocka and Kłosowska, 1966; Kotelko *et al.*, 1979; Spadowska *et al.*, 1979) that often take part in hydrogeochemical transformations of sulphur (Rajchel *et al.*, 2000a, b). Denitrifying bacteria may develop in groundwater where sulphates are present but oxygen is lacking. They reduce the sulphates to hydrogen sulphide, evolving H₂S migrating up into zones with available oxygen and facilitating the development of aerobic bacteria. Desulphatising bacteria oxidise hydrogen sulphide and other sulphur compounds into sulphates or sulphuric acid, while thionic bacteria oxidise the reduced sulphur compounds into sulphates or sulphuric acid.

Microorganisms contained in sulphurous waters of southern Poland have been studied by numerous authors: Szafer (1910), Strzeszewski (1913), Namysłowski (1922), Turowska (1934), Topińska-Luchterowa (1951) and others. Only those by Strzeszewski (*op. cit.*) and Turowska (*op. cit.*) discuss the springs in Swoszowice and Rytro, the subject of the research described here.

We have identified the microflora in these sulphurous springs and compared the results with earlier determinations, and attempted to establish links between water chemistry and microorganism species composition and colony form.

METHODS

Samples of microflora were collected in seven springs, during November, February, April and July of 1998/1999. They were taken from the bottoms of the spring niches and from spring outflows, using plastic jars with a capacity of about 120 ml. Simultaneously, water samples for hydrogen sulphide analyses and other standard determinations (Witczak and Adamczyk, 1995) were collected. Water analyses were carried out using the IPC-AES method (Rajchel, 2000). Physico-chemical parameters were measured in each sample (Table 1). The taxonomic study

Table 1

Physico-chemical parameters of water (mg/dm³) from the springs investigated (after Rajchel, 1998, 2000)

Springs	t [°C]	pH	H ₂ S	Na ⁺	K ⁺	Li ⁺	Sr ²⁺	Ca ²⁺	Mg ²⁺	Fe ²⁺	Cl ⁻	HCO ₃ ⁻	SO ₄ ²⁻
Główne	10.2	6.7	62.8	46.80	8.50	0.09	5.97	547.0	92.00	0.01	62.05	564.4	1277.8
Napoleon	10.1	6.8	61.1	72.30	8.90	0.09	4.20	497.0	97.3	0.005	62.05	533.9	1247.8
Jacek	7.8	7.4	12.2	44.60	1.17	0.04	0.80	74.86	15.34	0.02	57.90	246.30	61.62
Paweł	8.8	7.1	15.5	6.06	1.62	0.01	0.20	79.70	28.93	0.01	8.50	362.30	39.62
Jerzy	8.8	7.5	10.2	10.75	1.69	0.02	1.04	51.70	17.84	0.09	3.60	228.10	22.11
Katarzyna	9.3	7.8	10.7	46.40	1.57	0.02	0.40	71.89	13.54	0.01	20.20	362.39	19.60
Stanisław	8.0	7.8	11.2	80.60	2.10	0.04	0.30	17.00	7.90	0.06	6.00	251.30	19.40

was based on live material, using an *ECLIPS 600 Nikon* light microscope. The identification of organisms was made following Häusler (1982), Kramer and Lange-Bertalot (1986a, b), Wołowski (1998) and Starmach (1966).

DESCRIPTION OF THE SPRINGS

The following springs from the Carpathians (Fig. 1) were selected for microbiological investigations: “Jacek” at Lipnica Wielka (Orawa), “Paweł” at Polichy near Ci kowice, “Katarzyna” at Rytro near Stary S cz, “Jerzy” at Wierchomla near Krynica and “Stanisław” at Łabowa near Krynica. Two

springs are situated in the Carpathian Foredeep: “Główne” and “Napoleon” at Kraków-Swoszowice (Rajchel, 1998, 2000; Rajchel and Rajchel, 1999; Rajchel *et al.*, 2000).

The “Jacek” spring at Lipnica Wielka is situated in the right, steep bank of the Lipniczanka stream and its small, right tributary, some 1.2 km upstream from the bridge across the former stream. It is a fracture spring with a diameter of about 0.25 x 0.35 m and a depth of 0.05 m, in an outcrop of thick-bedded sandstone that limits the niche from the slope. Incrustations and festoons of white desulphurising bacteria mark the streambed (Fig. 2a).

The “Katarzyna” spring at Rytro is sited in the scarp of the terrace, on the right bank of the Wielka Roztoka stream, ca. 300 m below a footbridge, opposite the last bus stop. Water




Fig. 1. The geological structure of the Carpathians and the Carpathian Foredeep (after Poprawa and Nem ok, 1988–1999; changed, simplified) and the location of analysed springs (after Rajchel, 2000)



Fig. 2. **a** — sulphuric bacteria with fringe-like appearance from the “Jacek” spring at Lipnica; **b** — violet mats of sulphuric bacteria in the “Katarzyna” spring at Rytro; **c** — violet mats of sulphuric bacteria in the “Pawel” spring at Polichy; **d** — sulphuric bacteria resembling thin filaments or spider-webs in the “Pawel” spring at Polichy; **e** — the outflow of the “Główne” spring at Swoszowice covered by purple sulphuric bacteria

flows from the fissure down the steep bank, covered by a thick layer of white bacteria, into an oval niche (Fig. 2b) with a diameter of 0.35 m and a depth of about 0.04 m, which has retreated because of backwards erosion. The niche bottom is covered with encrustations of white bacterial colonies.

The “**Jerzy**” spring at Wierchomla flows out from the large niche in the Izwór stream valley, on its left bank, ca. 1.8 km from the church, at the foot of a hill slope covered with a beech forest. The niche is oval with a diameter of 1.3 m, 0.5 m deep, its banks being lined with sandstone fragments and timbered. Water overflows the linings, leaving at the bottom of the spring niche a violet-coloured deposit of sulphurous bacteria, while bacterial colonies form white incrustations along the watercourse.

The “**Stanisław**” spring at Łabowa is found in a forest on the southern slope of Sapalska Hill, ca. 500 m from the forest edge. Its oval niche is 1.3 m in diameter and 1 m deep, the banks being lined with sandstone blocks up to a ground level. The stone lining is covered up to the water table with violet deposits composed of sulphurous bacteria.

The “**Paweł**” spring at Polichty is located on Sucha Hill, on the right bank of a stream. The niche is oval, 1.2 x 1.5 m across, and 0.3 m deep. Continuous, violet deposit of sulphurous bacteria cover its bottom (Fig. 2c), while the entire niche, filled with water, is covered with white bacteria that form a network of densely spaced (Fig. 2d) veinlets, 0.5 m long. Such veinlets have so far not been found anywhere else in the Carpathians.

The “**Główne**” spring at Kraków-Swoszowice is situated in the park, NW of the bath building, where the ground surface begins to slope toward the Wilga River valley. The spring takes the form of an artesian well, 10.2 m deep, lined with blocks of the Szydłowiec sandstone, situated inside a small wooden house resting on a stone foundation. Water flows out through a pipe, then along an open ditch and a short watercourse, emptying into the Wilga. The pipe is covered with thick encrustations of white sulphurous bacteria, while the bacteria deposited along the watercourse are purple (Fig. 2e).

The “**Napoleon**” spring at Kraków-Swoszowice is situated north of the “**Główne**” spring on the other side of the tarmac Swoszowice–Kraków road, at the bottom of a sandy gully. It is a natural outflow of sulphuric water from the mouth of a collapsed adit. The intake is in the form of a concrete ring, partly hidden inside the ground, and the water flows out from below the ring and down a narrow watercourse to the Wilga. White encrustations and purple covers of sulphurous bacteria can be seen along the stream.

RESULTS

About 31 taxa of microorganisms: Schizomycetes (16), Cyanophyceae — Cyanoprokaryota (3), Euglenophyceae (5), Bacillariophyceae (4) and Chlorophyceae (3) were identified in the seven springs investigated. Sulphuric bacteria belonging to genus *Chromatium* (Fig. 3a–d, j and l) occur abundantly and frequently in five of these springs. This type of bacteria occurs throughout the year in the “**Główne**” and the “**Napoleon**” spring at Swoszowice. During the summer and hot autumns of

1999 and 2000 these bacteria thrived as pinkish-red or pinkish-purple deposits. These springs have a high concentration of hydrogen sulphide, which favours the development of *Chromatium* species. In both these springs *Chromatium weissei* (Fig. 3c), *Ch. okenii* (Fig. 3j) occur abundantly. Also *Chlorobium vibrioforme* (Fig. 3h and i) formed masses here during the summer but it did not occur in the remaining springs investigated. More algal taxa occur in the “**Napoleon**” spring than in the “**Główne**” spring: *Pseudanabaena constricta* (Fig. 4j), *Lyngbya aeruginosa-coerulea* (Fig. 4i), *Navicula* sp. div., *Nitzschia linearis* (Fig. 5h), *Pinnularia viridis* (Fig. 5f), *Euglena mutabilis* (Fig. 5b), *E. viridis* (Fig. 5a) and *Lepocinclis ovum* (Fig. 5c) while *Euglena viridis* and *E. retronata* (Fig. 5d) developed abundantly during August 2000.

The taxa *Chromatium gracile*, *Ch. weissei* and *Thiocapsa roseopersicina* (Fig. 3k) occur as masses in the “**Paweł**” spring at Polichty, colouring the sediment pinkish-violet. *Achroonema spiroideum* (Fig. 4k) and *Chlorochromatium aggregatum* (Fig. 3m), which were not observed in the other springs, developed here.

In the springs “**Jerzy**” at Wierchomla, “**Katarzyna**” at Rytro, and “**Jacek**” at Lipnica Wielka filamentous bacteria of the genus *Thiothrix* developed well. *Thiothrix nivea*, *Th. tenuissima*, *Beggiatoa alba* and *B. minima* occurred abundantly. These bacteria turn the sediments white. *Chromatium weissei* and *Ch. vinosum*, which also developed abundantly among filamentous bacteria in the “**Jerzy**” spring during the summer, coloured the surface of the spring bottom whitish-pink.

Only 5 taxa were recognised in the “**Stanisław**” spring of these *Thiocapsa roseopersicina* (Fig. 3k) occurred abundantly, colouring the sediments violet-pinkish. Only in this spring were a few specimens of the colourless euglenophyte *Entosiphon sulcatus* (Fig. 5e) were found.

Among the springs investigated, only the “**Główne**” and “**Napoleon**” at Swoszowice, and “**Katarzyna**” at Rytro, have been investigated earlier. Strzeszewski (1913) gave a list of taxa from the “**Napoleon**” and “**Główne**” springs where many *Chromatium* taxa occurred. From the “**Główne**” spring he reported *Ch. minus* and *Thiothrix nivea*, *Th. nivea* var. *verticillata*, *Beggiatoa alba*, *B. leptomitiformis*, *Lamprocystis roseopersicina*, and several taxa of cyanobacteria (5), diatoms (13), and green algae (2), which have not been found during this study. He found a similar, but more diverse bacterial and algal flora in the “**Napoleon**” spring. A lower number of organisms in those springs was noted 20 years later by Turowska (1934), who also found *Chromatium okenii* fo *minutissima* and *Thiocapsa agilissima*. She listed taxa of cyanoprokaryota (*Anabaena constricta*, *Oscillatoria geminata* var. *sulphurea*, *Oscillatoria trichoides*), and green algae *Stigeoclonium subsecundum*, *St. pygmaeum*, *Ulothrix subtilis*, *Oedogonium* sp. which have not been observed in this study.

The present list of taxa (Table 2) shows only 11 taxa of bacteria and 11 taxa of other organisms, in the “**Główne**” and “**Napoleon**” springs, fewer than reported by Strzeszewski (1913, about 50 taxa), and Turowska (1934, about 19 taxa). This is probably connected with regulation of the outflow of the springs and local extinction of some populations. During Strzeszewski’s (1913) investigation, a small equalising tank

Table 2

Abundance of microflora in the springs investigated

Taxa	Springs/Sites						
	“Główne” Swoszowice	“Napoleon” Swoszowice	“Jerzy” Wier- chomla	“Katarzyna” Rytro	“Paweł” Polichy	“Stanisław” Łabowa	“Jacek” Lipnica Wielka
Schizomycetes							
<i>Achroonema spiroideum</i> Skuja	–	–	–	–	1	–	–
<i>Beggiatoa alba</i> Trevisan	–	–	–	1–2	–	–	1
<i>Beggiatoa minima</i> Vinogradskij	–	–	2	1–2	–	–	2
<i>Chlorochromatium aggregatum</i> Lauterborn	–	–	–	–	1	–	–
<i>Chlorobium vibrioforme</i> Pel’s	3	3	–	–	–	–	–
<i>Chromatium gracile</i> Strzeszewski	1	–	–	–	2	1	–
<i>Chromatium weissei</i> van Niel	3	3	1–3	–	3	–	–
<i>Chromatium vinosum</i> Vinogradskij	+	2	1	–	+	–	–
<i>Chromatium minus</i> Vinogradskij	2	–	–	–	–	–	–
<i>Chromatium okenii</i> Perty	2	2	–	–	–	–	–
<i>Thiocapsa roseopersicina</i> Vinogradskij	+	–	–	–	+	3	–
<i>Thiocystis violacea</i> Vinogradskij	+	–	1–2	–	–	–	–
<i>Thiothrix annulata</i> Molisch	–	–	2	–	–	1	–
<i>Thiothrix minutissima</i> Upehof	–	+	–	2	–	–	2
<i>Thiothrix nivea</i> Vinogradskij	–	1	3	3	–	–	3
<i>Thiothrix tenuissima</i> Vinogradskij	1–3	1	1–3	2	–	–	2–3
Cyanophyceae — Cyanoprokaryota							
<i>Pseudanabaena constricta</i> (Szafer) Lauterborn	1	1	–	–	–	1	–
<i>Lyngbya aerugineo-coerulea</i> (Kützing) Gomont	–	1	–	–	–	–	–
<i>Oscillatoria tenuis</i> Agardh	–	+	–	1	–	–	–
Euglenophyceae							
<i>Entosiphon sulcatus</i> Stein	–	–	+	–	–	+	–
<i>Euglena mutabilis</i> Schmitz	–	1	+	–	+	–	–
<i>Euglena retronata</i> Johnson	–	3	–	–	–	–	–
<i>Euglena viridis</i> Ehrenberg	–	1	1	–	–	–	–
<i>Lepocinclis ovum</i> Minkiewicz	–	+	–	–	+	–	–
Bacillariophyceae							
<i>Gomphonema parvulum</i> (Kützing) Grunov	–	1	–	–	–	–	–
<i>Navicula</i> sp. div.	1	1	–	–	–	1	–
<i>Nitzschia linearis</i> W. Smith	–	1	1	–	+	–	–
<i>Pinnularia viridis</i> Ehrenberg	–	1	1	–	+	–	1
Chlorophyceae							
<i>Klebsormidium tribonematoideum</i> Hindák	–	–	–	1	–	–	–
<i>Microthamnion kützingianum</i> Nägeli	–	–	–	+	–	–	–
<i>Stigeoclonium tenue</i> Kützing	–	–	–	–	1	–	1
Number of taxa ()	11	18	12	7	11	6	7

+ — single, 1 — sparse, 2 — frequent, 3 — very frequent

was lined with wood and overgrown by vascular plants, giving good conditions for the development of cyanobacteria and diatom species. At present, the tank is covered with concrete, there are no plants, and the H₂S concentration is similar to that in the spring, limiting the development of algae. We have found some taxa which were not observed (or recognised) during the earlier investigations: individual specimens of *Euglena viridis*, *E. mutabilis*, *E. retronata* and *Lepocinclis ovum*.

Comparing our results with those of Turowska (1934) on the “Katarzyna” spring it can be seen that the occurrence of bacteria in this spring is almost the same as it was 65 years ago. She found

Beggiatoa leptomitiformis, *B. arachnoidea* and a few taxa of *Oscillatoria* which the present authors did not find i.e. *Oscillatoria kützingii*, *O. limnetica* and *O. geminata* var. *sulphurea*. Neither we nor Turowska found species of *Chromatium* in the spring.

CONCLUSIONS

Altogether 16 taxa of bacteria, 3 taxa of cyanoprokaryota and 12 taxa of algae were found in the springs investigated. Sulphuric bacteria are the most frequently occurring organisms




Fig. 3. a, b — *Chromatium vinosum* (2400 x), c — *Chromatium weisseii* (1200 x), d — *Chromatium gracile* (4000 x), e–g — *Thiocystis violacea* (e, g — 650 x, f — 2400 x), h, i — *Chlorobium vibrioforme* (4000 x), j — *Chromatium okenii* (1200 x), k — *Thiocapsa roseopersicina* (1000 x), l — *Chromatium minus* (1400 x), m — *Chlorochromatium aggregatum* (1700 x); a–b — original, c–m — after Häusler (1982)

in the springs (Table 2). Usually they cover a hard surface forming thin filaments resembling a spider web or a soft coat plait.

Taxa of the genus *Chromatium* (Fig. 3a–d, j and l) were found in five springs out of the seven. They proliferated during the summer and autumn, giving an overall pinkish-red colour to the surface of leaves, twigs and stones. They were observed frequently and in abundance throughout the year in the “Główne” and “Napoleon” springs at Swoszowice. *Thiothrix* and *Beggiatoa* taxa occur abundantly in the “Katarzyna” and

“Jacek” springs, forming a greyish-white filamentous mat. Interesting taxa of *Chlorochromatium aggregatum* and *Chlorobium vibrioforme* (Fig. 3h and i) were occasionally found during summer at the “Pawel” and “Główne” springs.

Among the algal group, Cyanoprokaryota (3 taxa), Euglenophyceae (5), Bacillariophyceae (4) and Chlorophyceae (3) were found. All of them occurred as individual specimens. Only *Euglena viridis* and *E. retronata* (Fig. 5a and d) formed a light green bloom during the autumn of 2000 in the “Napoleon”




Fig. 4. a–c—*Thiothrix nivea* (a, b—1000 x, c—60 x), d—*Thiothrix annulata* (250 x), e—*Beggiatoa minima* (1000 x), f—*Beggiatoa alba* (900 x), g, h—*Oscillatoria tenuis* (g—2000 x, h—1000 x), i—*Lyngbya aerugineo-coerulea* (1000 x), j—*Pseudanabaena constricta* (1000 x), k—*Achroonema spiroideum* (2000 x); a–c, k—after Häusler (1982), d, f—after Huber-Pestalozzi (1938), e—original, g, h—after Starmach (1966)

spring. Most of the algal taxa occurred only at spring outlets, this being connected with a lower sulphur concentration in the water.

There are few diatom taxa relative to other freshwater springs (Namysłowski, 1922; Wołoszka, 1922; Foged, 1951; Round, 1960). Among other taxonomical groups: *Vaucheria*, *Tribonema*, *Mougeotia*, *Spirogyra*, *Drapanaldia* and *Batrachospermum* frequently occur. There is no doubt that characteristic groups of algae develop in different types of

springs, e.g. in springs with high concentrations of sulphides the community of algae consists of *Lyngbya* and *Phormidium* species; in oligohaline springs community of *Cladophora-Cocconeis-Enteromorpha* species usually occur (Kawecka and Eloranta, 1994).

In the springs investigated, when the H_2S concentration reached about 7 mg/dm^3 , a bacterial flora of *Thiothrix* and *Beggiatoa* occurred at the bottom and around the rim and outlet of the spring as single, very tiny filaments covering the surfaces




Fig. 5. **a** — *Euglena viridis* (2000 x), **b** — *Euglena mutabilis* (2000 x), **c** — *Lepocinclis ovum* (1500 x), **d** — *Euglena retronata* (1000 x), **e** — *Entosiphon sulcatus* (1000 x), **f** — *Pinnularia viridis* (1000 x), **g** — *Gomphonema parvulum* (2000 x), **h** — *Nitzschia linearis* (1000 x); **a**, **b**, **e** — after Wołowski (1998), **c** — after Starmach (1983), **f**–**h** — after Siemińska (1964)

with a white spider-web-like pattern. At higher concentrations of H_2S the white colonies of bacteria look like fringes or thin mats. Among these *Chromatium* sp. or *Thiocapsa* sp. locally developed, giving a pinkish colour to some parts of the sediments. In spring, when the concentration of H_2S was higher than 50 mg/dm^3 , pink mats of bacteria (*Chromatium weissei*, *Ch. vinosum*) cover the surface of the spring bottom as in the “Główne” and “Napoleon” at Swoszowice. The pinkish-violet bacteria represented by *Thiocapsa roseopersicina* occur at the

bottom of the “Stanisław” spring at Łabowa and the “Pawel” spring at Polichy. It was observed that sulphuric bacteria in the springs were rather stagnant throughout the year (Table 1). White filamentous and violet coccal bacteria occurred all the year round in equal amounts. Pink and purple bacteria developed well in late spring and early autumn. The colour and shape of agglomerated bacteria can indicate the level of hydrogen sulphide in a spring (Table 1 and 2).




Fig. 6. **a** — *Stigeoclonium tenue* (500 x), **b** — *Microthamnion kützingianum* (500 x), **c** — *Klebsormidium tribonematoideum* (1200 x); **a**, **b** — after Starmach (1972), **c** — after Hindák (1996)

Acknowledgements. The authors are grateful to Mrs C. Waligórska-Acheson for help with the English version of this work.

We would like to express our gratitude to Prof. dr hab. A. Giziński and Prof. dr hab. in . A. Sadurski for their review of the manuscript and valuable remarks and opinions.

This research has been supported financially by the Faculty of Geology, Geophysics and Environment Protection of the University of Mining and Metallurgy in Kraków — grant No. 11.11.140.808 and statutory research of the Chair of Stratigraphy and Regional Geology in 2002.

REFERENCES

- FOGED N. (1951) — The diatom flora of some Danish springs. Strandkaer. The Mols-laboratory. Nat. Jutlandica, **4**: 1–84.
- HÄUSLER J. (1982) — Schizomycetes. Bakterien. In: Süßwasserflora von Mitteleuropa (eds. H. Ettl, J. Gerloff and H. Heynig), **20**.
- HINDÁK F. (1996) — Key to the unbranched filamentous green algae (Ulortichineae, Ulotrichales, Chlorophyceae) (in Slovak). Bull. Slovenskej Botan. Spolecnosti SAV, Suppl. 1. Bratislava.
- HUBER-PESTALOZZI G. (1938) — Das Phytoplankton des Süßwassers Systematik und Biologie. Allgemeiner Teil Blaualgen. Bakterien. Pilze, B. **16** (1). E. Schweizerbart'sche Verlag. Stuttgart.
- JAROCKA A. and KŁOSOWSKA T. (1966) — Studia mikroflory swoistej wód leczniczych. Problemy Uzdrowskowe, **30** (3): 7–25.
- KAWECKA B. and ELORANTA P. V. (1994) — Zarys ekologii glonów wód słodkich i rodowisk l dowych. PWN. Warszawa.
- KOTELKO K., SEDLACZEK L. and LACHOWICZ T. (1979) — Biologia bakterii. PWN. Warszawa.
- KRAMER K. and LANGE-BERTALOT H. (1986a) — Bacillariophyceae. Naviculaceae. In: Süßwasserflora von Mitteleuropa (eds. H. Ettl, J. Gerloff and H. Heynig), **2** (1).
- KRAMER K. and LANGE-BERTALOT H. (1986b) — Bacillariophyceae. Bacillariaceae, Epithemiaceae, Suriellaceae. In: Süßwasserflora von Mitteleuropa (eds. H. Ettl, J. Gerloff and H. Heynig), **2** (2).
- MACIOSZCZYK A. (1987) — Hydrogeochemia. Pa stw. Inst. Geol. Warszawa.
- NAMYSŁOWSKI B. (1922) — La microflore des sources subalpines (in Polish with French summary). Kosmos, **47** (1–3): 204–232.
- POPRAWA D. and NEMOK J. (1988–1999) — Geological Atlas of the Western Outer Carpathians and their Foreland. Pa stw. Inst. Geol. Warszawa.
- RAJCHEL L. (1998) — Wody mineralne i akrotopegi Krakowa. Prz. Geol., **46** (11): 1139–1145.
- RAJCHEL L. (2000) — Źródła wód siarczkowych w Karpatach polskich. Geologia, **26** (3): 309–373.
- RAJCHEL J., MARSZAŁEK M. and RAJCHEL L. (2000a) — Deposits of sulphurous spring waters from the Carpathians and the Carpathian Foredeep (southern Poland) (in Polish with English summary). Prz. Geol., **48** (12): 1174–1180.

- RAJCHEL J., MARSZAŁEK M. and RAJCHEL L. (2000b) — Minerals of sulphurous spring sediments from the Carpathians and Carpathian Foredeep. *Acta Miner.-Petrol.*, **41**, suppl.: 91.
- RAJCHEL L. and RAJCHEL J. (1999) — Carpathian springs of mineral and therapeutic waters — monuments of inanimate nature (southern Poland) (in Polish with English summary). *Prz. Geol.*, **47** (10): 911–919.
- RAJCHEL L., RAJCHEL J. and WOŁOWSKI K. (2000) — Mikroflora wód siarczkowych. XVII Konf. Paleont. Kraków, 21–23. IX. 2000. Historia basenów sedimentacyjnych a zapis paleontologiczny: 62–64.
- ROUND F. E. (1960) — A note of the diatom flora of some springs in the Malham Tarn area of Yorkshire. *Arch. Protistenk.*, **104**: 515–526.
- SIEMIŃSKA J. (1964) — Chrysophyta II. Diatoms — Bacillariophyceae. *Freshwater Flora of Poland*. PWN. Warszawa. Kraków.
- SPANDOWSKA S., DANIELAK K. and ZIEMOWSKA A. (1979) — Metodyka bakteriologicznego badania wód podziemnych i gruntów. *Inst. Geol. Warszawa*.
- STARMACH K. (1966) — Cyanophyta — sinice, Glaucophyta — glaukofity. In: *Flora słodkowodna Polski* (eds. K. Starmach and J. Siemińska), **2**. PWN. Warszawa.
- STARMACH K. (1972) — Chlorophyta III. Zielenice nitkowate. In: *Flora słodkowodna Polski* (eds. K. Starmach and J. Siemińska), **10**. PWN. Warszawa.
- STARMACH K. (1983) — Euglenophyta — eugleniny. In: *Flora słodkowodna Polski* (eds. K. Starmach and J. Siemińska), **3**. PWN. Kraków.
- STRZESZEWSKI B. (1913) — Beitrag zur Kenntnis der Schwefelflora in der Umgebung von Krakau. *Bull. Intern. Acad. Sci., Cracovie, Cl. Sci. Math. Nat., Sér. B., Sci. Nat.*, **6**: 309–334.
- SZAFER W. (1910) — Zur Kenntnis der Schwefelflora in der Umgebung von Lemberg. *Bull. Intern. Acad. Sci., Cracovie, Cl. Sci. Math. Nat., Sér. B., Sci. Nat.*, **3**, B: 161–167.
- TOPIŃSKA-LUCHTER A. (1951) — Problems concerning the relations between the microflora and the chemism of the waters in sulphur sources. *Pr. Komis. Farmac., Pol. Akad. Umiej.*, **3**: 225–299.
- TUROWSKA I. (1934) — Études sur la microflore des sources sulfureuses en Pologne. Contribution à l'étude des sulfobactéries. *Bull. Intern. Acad. Pol. Sci. Lettres, Cl. Sci. Math. Nat., Sér. B 1: Sci. Nat.* (1933): 45–66.
- WITCZAK S. and ADAMCZYK A. (1995) — Katalog wybranych fizycznych i chemicznych wskaźników zanieczyszczeń wód podziemnych i metod ich oznaczania. *Biblioteka Monitoringu środowiska*. Państw. Inst. Ochr. środowiska. Warszawa.
- WOŁOSZYŃSKA J. (1922) — Winter-Flora der Morenenguel des Wigrysees (in Polish with German summary). *Kosmos*, **47**: 305–326.
- WOŁOWSKI K. (1998) — Taxonomic and environmental studies on euglenophytes of the Kraków-Cz. stochowa Upland (Southern Poland). *Fragm. Flor. Geobot., suppl.*, **6**: 3–192.