

Marta KANDZIORA¹, Ryszard CIEPAŁ¹
and Aleksandra NADGÓRSKA-SOCHA¹

**HEAVY METALS AND SULPHUR ACCUMULATION
IN THE *Picea abies* L. Karst. NEEDLES AND SOIL
OF THE FOREST PROMOTIONAL COMPLEX
“LASZY BESKIDU ŚLĄSKIEGO”**

**AKUMULACJA METALI CIĘŻKICH I SIARKI
W SZPILKACH *Picea abies* L. Karst I GLEBIE NA TERENIE LEŚNEGO
KOMPLEKSU PROMOCYJNEGO „LASZY BESKIDU ŚLĄSKIEGO”**

Abstract: The investigation of zinc, cadmium, lead, copper and sulphur contents was carried out in the soil and in the needles of *Picea abies* L. Karst. in the Forest Promotional Complex “Lasy Beskidu Śląskiego”. Samples of soil (from the level 0–10 cm) and annual, biennial and triennial needles of spruce were collected in autumn 2005 and 2006 from top parts of mountains: Rownica, Barania Góra and Czantoria. The highest heavy metals concentration were determined in triennial needles of *Picea abies* collected from Czantoria and Barania Góra. The lowest content of sulphur and investigated metals was found in soil samples and heavy metals in plant material from the Równica. Cadmium content (1.2 µg/g) was higher than normal level (0.2–0.8 µg/g) only triennial needles collected from Czantoria. The lead concentration (118–165 µg/g) in the soil of all investigated areas was exceeded, lead level considered as allowable. Relatively low microelement concentrations in the soil and the needles of *Picea abies* L. Karst. indicated poor plant nutrition.

Keywords: heavy metals, sulphur, *Picea abies* L. Karst., Beskid Śląski

Introduction

Spruce (*Picea abies* L. Karst.) is one of the most endangered species of trees in Europe, with the highest percentage of damaged trees (30 %). There is a close relation between the amount of trees damaged and the level of pollution in a given area. The chemical composition of the assimilation apparatus of spruce does not only indicate the contents of nutrients but also shows the degree of the environmental pollution [1].

¹ Department of Ecology, Silesian University, ul. Bankowa 9, 40-007 Katowice, Poland, email: marta.kandziora@yahoo.pl

Thus, the most endangered spruce are those from Poland, the Czech Republic and Germany, which is a result of high deposition of pollution in those countries [2].

Strong anthropopressure is characteristic of Forest Promotional Complex “Lasy Beskidu Śląskiego”. It results from the vicinity of large industrial areas of the Upper Silesia, Ostrava and Karvina District in the Czech Republic [2], as well as from the pollution emitted by transport and agriculture. On the basis of the analysis of assimilation apparatus, spruce trees from Beskid Śląski have been categorized into the 1st and 2nd classes of damage [2]. It is assumed that the high level of pollution until mid-90s had a huge impact on the condition and vitality of the trees. The first symptoms of the spruce forests dying out were observed in the 50s in Beskid Mały, in the 70s the problem was discovered in Beskid Śląski, and later also in Beskid Żywiecki [4]. In 2006 the amount of dead wood and coarse woody debris was estimated at 118,000 m³ in the forests of Beskid Śląski, which was ten times more than in the previous years. In addition, every year the number of sanitary cuts increases which is caused by the unfavourable weather conditions and the increasing threat of the insect pests [5].

Dying out of the spruce forests cannot of course be attributed just to one disadvantageous factor. The bad condition of the stands results from a variety of causes, accumulated over a period of time, which overlap and strengthen the final effect – dying out of a forest. Among diverse factors which have a negative influence on the stand, the industrial pollution, mostly heavy metals and sulphur, is the most significant factor in this process.

The research aimed at estimating heavy metals (Cd, Pb, Zn and Cu) as well as sulfur accumulation in the upper levels of the forest soil and in the accumulation apparatus of spruce in the stands of Forest Promotional Complex “Lasy Beskidu Śląskiego” in relation to dying of the spruce forest.

Material and methods

The investigation was carried out in the soil and needles of *Picea abies* L. Karst. in the Forest Promotional Complex “Lasy Beskidu Śląskiego”. Samples of soil (from the level 0–10 cm) and annual, biennial and triennial needles of *Picea abies* were collected in autumn 2005 and 2006 from top parts of mountains: Rownica, Barania Góra and Czantoria). In order to determine the heavy metal concentration, spruce needles were divided into three groups (one-, two- and three-year-old), dried at 105 °C to a constant weight, ground to a powder, then mineralized and dissolved in 10 % HNO₃. After filtration Cd, Pb, Zn and Cu content were determined using flame Atomic Absorption Spectrometry (AAS) [6]. Soil was air dried and extracted with 10 % HNO₃ and measurements were made using the conventional Atomic Absorption Spectrometry (AAS) [6]. The sulphur (total sulphur) content in plants and soil was determined with colorimetric method [6, 7]. The quality of analytical procedures was controlled by using the reference material (Certified Reference Material CTA–OTL–1 Oriental Tobacco Leaves). The data were processed using software Statistica.

Results and discussion

Results are shown in Table 1, 2.

Table 1

Content of cadmium, lead, zinc, cooper [$\mu\text{g/g}$] and sulphur [%] in the needles *Picea abies* L. Karst.
Values with the same letter are statistically the same for $p < 0,05$

Location	Sample	Cd [$\mu\text{g/g}$]	Pb [$\mu\text{g/g}$]	Zn [$\mu\text{g/g}$]	Cu [$\mu\text{g/g}$]	S [%]
Barania Gora	annual needles	0.27b	0.53a	25.2a	1.75a	0.052a
	biennial needles	0.325a	0.38a	25.045a	1.685a	0.045a
	triennial needles	0.354a	1.2a	24.115a	1.29a	0.026a
Czantoria	annual needles	0.755a	0.535a	22.415a	1.285a	0.087b
	biennial needles	0.49b	1.155a	25.46b	2.298a	0.043a
	triennial needles	1.175b	1.625b	54.225b	4.237a	0.065b
Rownica	annual needles	0.54a	0.32a	15.67b	1.32a	0.063a
	biennial needles	0.43c	1.01a	11.23a	1.14b	0.042a
	triennial needles	0.76c	0.37c	9.435c	1.7b	0.060b

Table 2

Content of cadmium, lead, zinc, cooper and sulphur [$\mu\text{g/g}$] and in the upper layer of the soil
Values with the same letter are statistically the same for $p < 0,05$

Location	Cd [$\mu\text{g/g}$]	Pb [$\mu\text{g/g}$]	Zn [$\mu\text{g/g}$]	Cu [$\mu\text{g/g}$]	S [$\mu\text{g/g}$]
Barania Gora	1.62a	165.6a	37.4a	9.4a	147.3a
Czantoria	1.24b	155.6b	26.9b	6.9b	100.1b
Rownica	0.96c	118.6b	11.5c	6.6b	81.0b

The concentration of cadmium, which is considered highly toxic, fluctuates in the examined material from 0.27 $\mu\text{g/g}$ dry mass in the needles of one-year-old spruce from Barania Gora to 1.175 $\mu\text{g/g}$ in the needles of three-year-old from Czantoria. Cadmium concentration in the soil remains within the normal range, from 0.96 $\mu\text{g/g}$ on Rownica to 1.62 $\mu\text{g/g}$ on Barania Gora. In the plant material from Czantoria and in the soil from Barania Gora the content of cadmium was exceeded, 0.2–0.8 $\mu\text{g/g}$ Cd in plants and to 1 $\mu\text{g/g}$ Cd in the soil [8].

Lead is also considered a very toxic element and its content in the plants shows the influence of anthropogenic factors. The concentration of lead in the analyzed material falls between 0.32 $\mu\text{g/g}$ in one-year-old needles from Rownica and 1.625 $\mu\text{g/g}$ in three-year-old needles from Czantoria, and in the soil between 118.6 $\mu\text{g/g}$ on Rownica to 165.6 $\mu\text{g/g}$ on Barania Gora. Hence, it can be concluded that in all investigated areas, the normal range of lead (100 $\mu\text{g/g}$ Pb) [8] has been exceeded.

Zinc is an element which is indispensable to a proper functioning of living organisms. Its content may fluctuate in a wide range from 10 to 100 $\mu\text{g/g}$ [8]. The concentration of

zinc in the examined samples of plant material was in the range from 9.435 $\mu\text{g/g}$ (in the three-year-old spruce needles from Rownica) to 54.225 $\mu\text{g/g}$ (in the three-year-old spruce needles from Czantoria) and in the soil from 11.5 $\mu\text{g/g}$ on Rownica to 37.4 $\mu\text{g/g}$ on Barania Gora, which does not exceed the norms and even shows zinc deficiency.

The concentrations of copper in the analysed samples are within the normal ranges from 1.14 $\mu\text{g/g}$ in the two-year-old spruce needles from Czantoria and do not exceed the norms (2–20 $\mu\text{g/g}$ Cu) [8]. Similarly, the content of copper in the soil from 6.6 $\mu\text{g/g}$ on Rownica to 9.4 $\mu\text{g/g}$ on Barania Gora are within the normal range of concentration (1–140 $\mu\text{g/g}$ Cu) [8].

Sulphur, similarly to zinc and copper, is absolutely crucial for the development of living forms. Its higher concentration in the plants is often a result of air pollution caused by sulphur dioxide. The content of sulphur fluctuated between 0.026 % in the three-year-old needles from Barania Gora to 0.087 % in one-year-olds from Czantoria. Sulphur content in the soil falls between 81.0 % (Rownica) and 147.3 % (Barania Gora).

The research showed, that heavy metals concentration in spruce needles and soil was similar to those of Zwolinski study (Cd 0.09–0.46 $\mu\text{g/g}$, Pb 1.9–2.5 $\mu\text{g/g}$, Zn 11.0–28.2 $\mu\text{g/g}$, Cu 2.0–2.8 $\mu\text{g/g}$, S 0.096–0.12 % in spruce needles and Zn 20–56 $\mu\text{g/g}$, Cu 7–13 $\mu\text{g/g}$, Cd 0.6–1 $\mu\text{g/g}$ in soil) [3]. Only lead concentrations (118.6–165.6 $\mu\text{g/g}$) in soil were considerably higher than Zwolinski results. This study differed from data for 'Male Pieniny' [1], only concentration of zinc (12.5–43.5 $\mu\text{g/g}$) was similar but concentrations of cadmium (0.01–0.1 $\mu\text{g/g}$) and sulphur (mean 0.023 %) were lower and the content of lead (1.0–2.0 $\mu\text{g/g}$) was higher than in this study. Similar situation was reported in the research conducted in Podhale Region. The concentrations of zinc (31.8–34.4 $\mu\text{g/g}$) and lead (1.6–1.7 $\mu\text{g/g}$) were lower and the content of cadmium (0.24–0.34 $\mu\text{g/g}$) was higher [1, 9]. The content of sulphur in Forest Promotional Complex "Lasy Beskidu Śląskiego" was lower than previously reported by Barszcz and Malek (annual needles 0.103–0.138 %, biennial needles 0.094–0.121 %) [10]. The concentrations of zinc in annual, biennial and triennial needles were similar to German research results (annual – 30.7 $\mu\text{g/g}$, biennial 27.6 $\mu\text{g/g}$, triennial 24.8 $\mu\text{g/g}$) [11]. The contents of cadmium and lead were higher than in Norway (0.011–0.173 $\mu\text{g/g}$ Cd, 0.03–0.89 $\mu\text{g/g}$ Pb), but lower than in Slovakia (1.19 $\mu\text{g/g}$ Cd, 1.73 $\mu\text{g/g}$ Pb) [11, 12].

The spruce forests in Beskid Śląski do not show an excessive accumulation of heavy metals in the needles. The content of metals both in the soil and in the needles is within the normal range. Only lead displays excessive concentration in the soil, which indicates that the soil is polluted with this metal. It is also noticeable that those forests are deficient in the basic nutrients, which has been noticed also in his studies by Zwolinski [2].

References

- [1] Panek E. and Szczepańska M.: *Metale śladowe i siarka w wybranych gatunkach roślin w Małych Pieninach*. Gospod. Surowc. Mineral., 2005, **21**(1) 89–109.
- [2] Zwoliński J.: *Ocena zagrożenia lasów świerkowych w Beskidzie Śląskim przez zanieczyszczenia powietrza atmosferycznego*. Pr. Inst. Bad. Leśn. A, 2003, 1(951) 53–68.
- [3] Lech P. and Żółciak A.: *Uwarunkowania występowania opieńkowej zgnilizny korzeni w lasach Beskidu Śląskiego*. Leśne Pr. Badaw., 2006, **2**, 33–49.

- [4] Fronczak K.: *Dlaczego wymierają świerki*. Przynr. Polska, 2006, **10**, 25–27.
- [5] Fronczak K.: *W trosce o beskidzkie świerczyny*. Głos lasu, 2007, **1(439)**, 2–5.
- [6] Ostrowska A., Gawliński S. and Szczubiałka Z.: *Metody analizy i oceny właściwości gleb i roślin*. IOŚ, Warszawa 1991.
- [7] Saalbach E., Essen G. and Judel G.K.: Pflanz. Dung. Bodenkunde, 1961, **93**, 17–26.
- [8] Kabata-Pendias A. and Pendias H.: *Biogeochemia pierwiastków śladowych*. PWN, Warszawa 1993.
- [9] Panek E. and Józefko U.: *Trace metals (Cd, Cr, Pb, Zn) and sulphur in spruce Picea abies L. of the road-side in the Podhale region, southern Poland*. Macro and Trace Elements, 21 workshop. 18–19.10.2002 Jena, Gemany, s. 249–255.
- [10] Barszcz J. and Małek S.: *Stability of Norway spruce (Picea abies L. Karst) stands in the Beskid Śląski and Beskid Żywiecki Mts. From the aspect of their nutrition status*. J. Forest Sci., 2008, **54(2)**, 41–48.
- [11] Trimbacher C. and Weiss D.: *Needle surface characteristic and element contents of Norway spruce in relation to the distance of emission sources*. Environ. Pollut., 1999, **105**, 111–119.
- [12] Čeburus D. and Steinnes E.: *Conifer needles as biomonitors of atmospheric heavy metal deposition comparison with mosses and precipitation, role of the canopy*. Atmos. Environ. 2000, **34**, 4265–4271.

**AKUMULACJA METALI CIĘŻKICH I SIARKI W SZPILKACH *Picea abies* L. Karst.
I GLEBIE NA TERENIE LEŚNEGO KOMPLEKSU PROMOCYJNEGO
„LASY BESKIDU ŚLĄSKIEGO”**

Katedra Ekologii, Uniwersytet Śląski

Abstrakt: Prowadzono badania dotyczące zawartości metali ciężkich (Zn, Pb, Cu, Cd) w glebie i szpilkach *Picea abies* L. Karst. na terenie Leśnego Kompleksu Promocyjnego „Lasy Beskidu Śląskiego”. Próbki gleby (z górnego poziomu) oraz szpilki jedno-, dwu- i trzyletnie świerka pobierano ze szczytowych partii gór: Równicy, Baraniej Góry i Czantorii jesienią 2005 i 2006 r. Stwierdzono najwyższe stężenia metali ciężkich w igłach trzyletnich zbieranych na Baraniej Górze i Czantorii. Najniższe koncentracje badanych metali i siarki zanotowano w próbkach gleby oraz w materiale roślinnym z Równicy. Zawartość Cd (1.2 µg/g) w trzyletnich igłach świerka pobieranych na Czantorii była wyższa od określanych jako stężenie normalne (0.2–0.8 µg/g). Stwierdzono przekroczenie dopuszczalnego poziomu dla Pb w glebie wszystkich badanych powierzchni (118–165 µg/g). Stosunkowo niskie stężenia mikroelementów zarówno w glebie, jak i igłach świerka wskazywać mogą na zły stan odżywienia roślin.

Słowa kluczowe: metale ciężkie, siarka, świerk pospolity *Picea abies* L. Karst., Beskid Śląski