

Molluscan immigrations via biogeographical ecotone of the Middle Russian Sea during the Jurassic

Mikhail ROGOV¹, Viktor ZAKHAROV¹ and Dmitry KISELEV²

¹ Geological Institute of Russian Academy of Sciences, Pyzhevskii Lane 7, Moscow – 109017, Russia;
e-mail: rogov_m@rambler.ru; zakharov@ginras.ru

² Yaroslavl State Pedagogical University, Kotorosl'naya nab. 46, Yaroslavl - 150000, Russia; e-mail: dnkiselev@mail.ru

Key-words: paleobiogeography, paleobiology, ecotone, Middle-Late Jurassic, Middle Russian Sea, stratigraphy, immigrational patterns.

ABSTRACT: Patterns of molluscan immigration via the biogeographical ecotone zone of the Middle Russian Sea during the Middle – Late Jurassic have been studied in detail. Firstly, the migration routes and ranges of recent molluscs are briefly reviewed, with some comments about their applicability to the Jurassic. Secondly, the migrational events have been classified by their direction, duration and intensity. We recognize two main types of immigration by their direction (unidirectional and multidirectional) and by their intensity (mass immigrations and isolated strayings). Recent progress in infrazonal ammonite biostratigraphy leads us to the more precise recognition of paleobiogeographical events and to the understanding of the immigration patterns that can produce more precise correlations.

INTRODUCTION

Jurassic molluscs, especially ammonoids, are the only group of fossils which has been traditionally used to elaborate new approaches in biostratigraphy such as the concept of zones by Oppel (1856-58) and the hemerae of Buckman (1893) as well as in paleobiogeography (Neumayr 1878; Uhlig 1911) and, recently, Sachs *et al.* (1971), Westermann (2000).

This paper summarises the results of recent detailed investigations of the biostratigraphy and paleobiogeography of the Middle Russian Sea (Fig. 1) during Middle-Late Jurassic time. This water body was chosen mostly because of its position as an ecotone between two main Superrealms during the Middle-Late Jurassic (Zakharov and Rogov 2004).

The so-called Middle Russian Sea flooded the East-European Platform during Bajocian-Bathonian time. After that for ca. 30 Myr this sea repeatedly changing its configuration, but always remaining connected with Boreal Basin via the Timan-Pechora Sea to the north. The southern connection with Peri-Tethyan (Caucasian) seas was not permanent and sometimes disappeared.

MIGRATION PATTERNS IN MODERN CEPHALOPODS AND BIVALVES AS KEY TO THE PAST

Modern cephalopods are represented by different living forms which inhabit all seas and oceans from tidal areas to the hadal zone. The taxa include both neritic and oceanic species, and both assemblages include pelagic and bottom dwellers,




Fig. 1. Middle Russian Sea and adjacent areas during the Late Jurassic (example of Late Kimmeridgian – Early Volgian): 1 – land areas; 2 – oceanic basins; 3 – shallow epicontinental seas; 4 – coastal areas, sometimes flooded by sea. Palaeogeography and paleolatitudes are from Thierry and Barrier (2000), corrected.

with the maximum diversity in the tropics and subtropics (Nesis 2003). They occur in water of normal marine salinity, between 27 and 37‰ (Jereb *et al.* 2005) with some rare exclusions. The bulk of modern cephalopods is characterized by very fast growth and short life-cycles with terminal spawning. Cephalopods are voracious, active predators (Nesis 1985; Jereb *et al.* 2005).

The major factors which influence cephalopod distribution are temperature, food and currents. Oceanic species usually undergo daily vertical migrations, occurring at depths of about 200 to 700 m during the day, and ascending into the uppermost 200 m for the night (Jereb *et al.* 2005). Some cephalopods are characterized by ontogenetic and/or seasonal migrations, either active (Ommastrephidae) or passive, through areas with closed circulation (*Spirula*) (Nesis 1985). Some neritic cephalopods have small natural habitats and some of them have very wide distribution, for example *Octopus vulgaris* (Nesis 1987). Oceanic coleoids usually have latitudinal distribution, but also pan-boreal, bi-subtropical amphi-atlantic ranges occur

(Nesis 2003). Post-mortal drift is common for Sepiidae and Spirulidae shells.

Cephalopods are very prone to expand and to reduce their species ranges quickly. After the opening of Suez Channel more than 200 different hydrobionts penetrated into the Mediterranean Sea, including the Indo-Pacific *Octopus kagoshimensis* (Salman *et al.* 2005). Recent warming of the climate (the so-called “Global warming”) has provoked a gradual disappearance of the squid *Loligo forbesi* from Portugal (Chen *et al.* 2006) and, presumably, well before this, in Morocco, where it was recorded in the past but never met during intensive investigations in 1995-1998 (Laptikhovsky, pers. comm). The influence of interspecific competition on the ranges of extant cephalopods is still insufficiently known. Inverse patterns of abundance of two commercial squids *Illex argentinus* and *Loligo gahi* has been suggested as the possible result of interspecific competition (Arkhipkin and Middleton 2002), and the same causal hypothesis was invoked to explain the strong depth segregation in some North African coleoids (Arkhipkin and Laptikhovski 2005). Nevertheless, numerous occasions of the coexistence of morphologically close species are known, amongst commercial taxa (Bonaud *et al.* 1998; Herke and Foltz 2002).

With respect to cephalopods, bivalves are slow-moving when adult, (except when attached to algae or wood) and their migratory ranges are restricted up to several kilometers, such as in *Macoma baltica* (Hiddink 2002). However, many species have planktonic larvae that can be widely dispersed by oceanic currents. The major factor limiting their spreading is water temperature during spawning. Temperature limits for gonad maturation and hatching could be much narrower in comparison with those of adult animals (Scarlatto 1981). Another important factor is the presence/absence of a suitable substrate for juvenile settling. Gastropod taxa with pelagic larvae are widely distributed along the west coast of South America, where the necessary rocky substrates are abundant. In contrast to this, the east coast of the continent is represented almost exclusively by sandy substrates, and species with direct development predominate from the subtropics

to Antarctica (Gallardo and Penchaszadeh 2001). Bivalves probably could suffer the same problem.

The critical temperatures differ between species and populations depending on latitude or seasonal temperature acclimatization. Peck and Conway (after Pörtner 2001) suggested that tropical and temperate species are more eurythermic than polar ones.

The polarbound limit of the species range in summer is determined in bivalves by the minimum temperature suitable for reproduction, and in winter by the minimum survival temperature. The equator-ward limit of the species distribution in winter is determined by the maximum temperature of reproduction whereas the maximum survival temperature restricts this boundary in summer (Scarlato 1981). These adaptive patterns permit the bivalves of high latitudes to inhabit the deep waters of the temperate zone, whereas tropical or subtropical species are able to penetrate far into high latitudes in summer. For example, a Boreal species *Astarte crenata* inhabits shallow areas in both the Barents Sea and the North Sea, and occurs in deep biotopes of the Gulf of Biscay. Some tropical-subtropical (for example, *Solen corneus*) and numerous subtropical relatively deep-water bivalves occur in the shallow habitats of Possjet Bay (42° N, Japan Sea) (Golikov and Scarlato 1967; Scarlato 1981).


These biological features controlling the geographical ranges of bivalves in modern seas, were important for bivalve spreading in the past. That is why, in our opinion, the southward spreading of natural habitats of Boreal bivalves (for example, belonging to the genera *Buchia* and *Retroceramus*) and the northward penetration of Tethyan Trigoniidae were caused by movement of larvae by currents. Another explanation of these events is the movement of water masses caused by transgressive-regressive events. In comparison to the situation with ammonites and other active poikilotherm animals, we should suppose more developed eurythermy within bivalves (Scarlato 1981). In spite of the planktonic mode of life of bivalve larvae, sometimes "ecological opportunity" rather than current direction influenced their dispersion (Amano 2005).

DISCUSSION ABOUT THE AMMONITE MODE OF LIFE AND THEIR MIGRATIONAL PATTERNS

The biogeography, life habitat and migrational patterns of ammonites were intensively studied

during the last decades (Kennedy and Cobban 1976; Olóriz 1990; Cecca 1992; Westermann 1996; Westermann and Tsujita 1999, among others). In a similar manner to modern cephalopods the ammonoids were possibly animals with different modes of life, from planktonic to quasi-nektonic and nekto-benthic ones. The ammonoids inhabited different biomes, and the oceanic groups of ammonites are usually different from those of neritic areas. Perhaps their penetration from shelf areas to the oceans, in contrast to coleoids, happened to planktonic, non nektonic, forms. It is very possible, that many ammonoids (especially heteromorphs) were vertical migrants (Westermann and Tsujita 1999), while others mostly swam above the bottom. The bulk of ammonoids was restricted to waters with normal oceanic salinity, with some exceptions like *Ceratites* which lived in restricted carbonate platforms surrounded by evaporate basins (Westermann and Tsujita 1999). Ontogenetic migrations leading to spatial segregation of juvenile and adult growth stages are also known in ammonoids (Westermann 1996).

Migrations which are related to long-time changes in the natural habitat, sometimes accompanied by diversification or sharp changes in fossil (ammonite) communities, should be separated from cyclic and repeated vertical, ontogenetic and seasonal migrations. Some of these types of migration in ammonites were recently reviewed by Reyment (2005) with some additions concerning necroplanktonic drift and "spurious migrations", caused by tectonics (continental drift, terrane movement, etc). Long-time and non-cyclic migrations we could call "immigrations", following the earlier usage of this term in ammonite biogeography (Westermann 1996; Fernandez-Lopez and Melendez 1996; Schweigert 2000; Navarro *et al.* 2005) and study of recent coleoids (Neige 2003), or "invasions" (Sei and Kalacheva 1983). The term "invasion" usually means a strong event, leading to the alteration of ecological relationships (*i.e.* mass migrations), while "immigration" could be used for all the characters of ammonite range movement. The analysis of ammonite immigrations involves difficulties due to problems in the precise outlining of their natural habitat (Bengtson and Kakabadze 1999). Ammonoids could be assigned to biogeographic categories (eudemic, midemic and parademic, carrying by currents, and so on (Fernandez-Lopez and Melendez 1996)), but even analyzing modern cephalopod areas we sometimes could



confuse these categories. Examples of post-mortem shell distribution published in the last decades were peculiar to the some groups of ammonoids only (Westermann 1996), especially from the oceanic biome.

In comparison to most of juvenile coleoids, very small ammonite hatchlings with a spherical shell filled by gas perhaps were easily scattered by currents. After growing and maturation, some ammonites remained planktonic while other began near-bottom or demersal life styles. There are different ways to reconstruct the mode of life in ammonites, from morphofunctional studies to facies and stable isotope analysis and to the studying of palaeopathologies and stable isotope signals (Ziegler 1967; Keupp and Ilg 1992; Westermann 1996; Anderson *et al.* 1994; Zakharov *et al.* 2006). We have to admit the high diversity in ecology of different ammonites.

Following Rawson (1973) we have recognized two different types of immigration according to intensity: 1) mass immigrations (or expansions) which lead to the movement of high-ranked biogeographic boundaries and alter the characteristics of ammonite communities, and 2) isolate straying, when immigrants are scarce and didn't give rise to new stocks. These types sometimes could be similar both spatially and temporally. One of the most unusual examples of the considerable distance travelled by small ammonite populations is shown by the aspidoceratids of latest Kimmeridgian age. Already living in the Subboreal seas of Poland and central areas of the Russian Platform, aspidoceratids of the late Autissiodorensis Chron are extremely rare (Kutek and Zeiss 1997; Rogov 2004). But some representatives of this group travelled as far as the Pechora River Basin and even the Subpolar Ural (Mesezhnikov 1984; Zakharov *et al.* 2005).

There are also two main types of immigrations which differ in their direction.

Unidirectional immigrations that usually coincide with movements of the biochore boundaries and drastically change ammonite communities.

Multidirectional (bidirectional) immigrations reflecting the simultaneous southward penetration of the Boreal taxa and the northward of Tethyan taxa. Perhaps, bidirectional immigrations occur rarely and reflect mainly a high sea-level stand. The biogeographical ecotone is usually enlarged at this time (Zakharov and Rogov 2004).

The ranges of ammonite species were influenced (with the exception of evolution) by

the following major factors: climate, geographical barriers and currents. Climate and geographical barriers often were interrelated, because an appearance or a disappearance of a barrier usually lead to a change in the water mass circulation and hence to climate change. The effect of currents was expressed mostly in the differences between fossil assemblages within different parts of a single basin with constant paleolatitude and sedimentation. A good example of sign of a cold paleocurrent via the Middle Russian Sea along the Urals during the Middle-Late Jurassic has been suggested by Kiselev (2004) due to the strong prevalence of Boreal ammonites. The same current also persisted during the Volgian (Rogov 2004).

MOLLUSCAN IMMIGRATIONS THROUGH THE MIDDLE RUSSIAN SEA

Any paleobiogeographical reconstruction, in contrast to modern biogeography, needs preliminary correlation and the revealing of the bio-events. It could be carried out accurately through the faunal horizons and their temporal equivalents, so-called hemerae (~ca. 0.08-0.2 My, after Page 1995). Only sometimes important changes in relative abundance of taxa were widely traced within the smaller part of faunal horizons.

The Middle Russian Sea existed on the boundary zone of two climatic belts, Boreal and Tethyan. Analysis of the biogeographical distribution of mollusks shows a permanent location of the biogeographical ecotone in this region (Zakharov and Rogov 2004). This is shown by the contemporaneous existence of the mixed Boreal, Subboreal and Submediterranean taxa in the bulk of the Middle-Upper Jurassic zonal assemblages (Hantzpergue *et al.* 1998; Mitta and Seltzer 2002; Rogov 2004). The Middle Russian Sea was influenced by Arctic and Peri-Tethyan water masses. This is demonstrated by the temporal dynamics of the Boreal and Tethyan elements in molluscan faunas (Figs 2-3).

Boreal taxa were dominant and widely distributed at the end of the Early (?) and the Late Bathonian, the earliest Callovian (when multidirectional migrations were important), in the latest Callovian and beginning of the Oxfordian, in the Late Oxfordian, in the beginning of the Middle Volgian and near to the Jurassic-Cretaceous boundary. Prolonged Tethyan influences occurred in the earliest Bathonian, in the beginning of




Fig. 3. Latitudinal and most important longitudinal immigrations of Boreal, Subboreal and Submediterranean molluscs during the Late Jurassic through Middle Russian Sea (see Fig. 2 for explanation).

the Late Callovian and during the Late Kimmeridgian – Early Volgian. Also two short but strong Tethyan events were restricted to the Oxfordian: in the Early Oxfordian (*Cordatium* Chron, *baccatum* hemera) and the Middle Oxfordian (*Densiplicatum*–*Tenuiserratum* chrons). The Late Kimmeridgian – Early Volgian penetration of the Peri-Tethyan ammonoids, such as the OPELLIIDAE and the ASPIDOCERATIDAE were mostly long termed in comparison with other Middle – Late Jurassic invasions. But even during this time the predominance of Peri-Tethyan taxa was combined with the presence of numerous Boreal genera (Fig. 3). In summary, the analysis of the fossil assemblages shows strong Boreal influence during the Middle-Late Jurassic. In spite of the mixed character of the faunas, Boreal taxa prevailed during the Callovian, Oxfordian, Early Kimmeridgian and Middle-Late Volgian. Nevertheless, by detailed investigation of the outcrops situated in the East-European Platform, representatives of both Peri-Tethyan and Boreal taxa are encountered within nearly all the ammonite zones.

Among the most important discoveries of the last few years influencing our paleobiogeographic knowledge, were the finding of true Boreal molluscs of Early Bathonian age in the Saratov Volga area (~37° N in paleolatitudes). These are the cardioceratid ammonites *Arcticoceras* and *Arctocephalites* and also the bivalve *Retroceramus* (Mitta *et al.* 2004). During the Late Bathonian the first signs of Boreal influence appeared in West Europe and, at the same time, in the Northern Caucasus, where recently some Bathonian *Kepplerites* have been discovered (Mitta 2003). The beginning of the Callovian coincides with the opening of the long-term Boreal-Tethyan connection of the Arctic and Tethys seas through the Middle Russian Sea perhaps caused by sea-level rise. Simultaneously Boreal *Paracadoceras* penetrated to the Northern Caucasus (and even its south slope, in Azerbaijan), while the Tethyan Macrocephalitidae (*M. jaquoti*) travelled as far as the Pechora region (Gulyaev 2005). During the Callovian some oscillations in the predominance of the Boreal, Subboreal and (scarcely) Submediterranean ammonites occurred. Major events of Tethyan ammonite immigration were observed in the late Early Callovian, when *Oxycerites* and *Parapatoceras* penetrated northwards to the Unzha River Basin (~46° N in paleolatitudes; Gulyaev 2002; Mitta 2004b), and in the beginning of the Late Callovian. The turn of the Callovian and Oxfordian has been marked

by the vast Boreal transgression connected with the southward migration of Boreal ammonites, especially cardioceratids (Boreal Spread of Arkell, 1956). These ammonoids mostly predominated during the Oxfordian, but during short warming episodes Tethyan cephalopods also inhabited the Middle Russian Sea (Fig. 3). The *Tenuiserratum* Chron could be listed among the strongest Tethyan faunistic influence intervals, when oppeliids (*Richeiceras*, *Creniceras*, *Glochiceras*, *Neoprioceras*) have penetrated to the Unzha River. The Early Kimmeridgian also was characterized by a few short events of the penetration of Submediterranean ammonites (Zakharov and Rogov 2004), while the Late Kimmeridgian shows even a predominance of these ammonites (aspidoceratids are common in the Eudoxus and earliest Autissiodorensis Chron, whereas oppeliids became numerous during the Autissiodorensis Chron and persist till the Early Volgian). Nevertheless, some strata in the Upper Kimmeridgian and Lower Volgian are characterized by numerous Boreal fossils (*volgae* horizon of the basal Autissiodorensis Zone; *pavida* horizon of the Sokolovi Zone). At the beginning of the Middle Volgian numerous Boreal *Pavlovia* penetrated to the Orenburg area (~36° N in paleolatitudes) and possibly even to the Northern Caucasus (Rogov 2004). The *Regularis* hemera of the Middle Volgian shows a wide spreading of *Zaraiskites*, from the Pechora area to the Pericaspian region. During the Middle-Late Volgian (*Virgatus* Chron and later) Submediterranean mollusks were very rare; only the bivalve family Trigonidae is known. At that time Boreal bivalves (*Buchia*) penetrated very far south. Recently some Middle-Upper Volgian *Buchia* have been recorded from the Minor Caucasus (~28° N in paleolatitudes; Zakharov and Kasumzadeh 2005). Latitudinal immigrations of Boreal ammonites also were common, especially around the Middle-Late Volgian transition. The latest Volgian deposits are characterized by the appearance of the Arctic *Volgidiscus* in the Yaroslavl region (Kiselev 2003). Significant Boreal influence also persisted through to the earliest Ryazanian, where ammonites of the genus *Praetollia* are known (Mitta 2004a).

CONCLUSION

The precise study of molluscan immigration leads to the more accurate recognition of paleogeography, while contemporaneous records of

fossils with different paleogeographic affinities permits the correlation of different biostratigraphic schemes. The European part of Russia seems to be one of the most suitable areas for the recognition of both the Boreal and Tethyan influences in the Middle-Late Jurassic due to its unique position between the Arctic Basin, Tethys and the West-European shallow seas.

Acknowledgements

This study has been supported by RFBR grants no.06-05-64284 and 09-05-00456, Russian Science Support Foundation and grants MK-3235.2006.5 and MK-856.2008.5 of the President of Russian Federation. We warmly thank Dr. V. Laptikhovskiy for his valuable comments about the migrational patterns of modern molluscs and their ecology and some references provided, and reviewers (G. Pavia, A. Wierzbowski and anonymous reviewer) for their comments.

REFERENCES

- Amano K. 2005. Migration and adaptation of late Cenozoic coldwater molluscs in the North Pacific. *In: A. T. Elewa (Ed.), Migration of organisms. Climate, geography, ecology*, 127-150. Springer.
- Anderson T. F., Popp B. N., Williams A. G., Ho L.-Z. and Hudson J. D. 1994. The stable isotopic records of fossils from the Peterborough Member, Oxford Clay Formation (Jurassic), UK: palaeoenvironmental implications. *Journal of Geological Society, London*, **151**: 125-138.
- Arkell W. J. 1956. *Jurassic Geology of the World*, I-XV, 1-806. Oxford, Oliver & Boyd.
- Arkhipkin A. and Laptikhovskiy V. 2005. Allopatric speciation of the teuthid fauna on the shelf and slope of Northwestern Africa. *2nd Intern. Symp. Coleoid Cephalopods through time, Prague 2005. Short papers/Abstract volume*, 7-10.
- Arkhipkin A. I. and Middleton D. A. J. 2002. Inverse patterns in abundance of *Illex argentinus* and *Loligo gahi* in Falkland waters: possible interspecific competition between squid. *Fisheries Research*, **59**: 181-196.
- Bengtson P. and Kakabadze M. V. 1999. Biogeography of Cretaceous ammonites – a review of procedures and problems. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, **212**: 221-239.
- Bonnaud L., Rodhouse P. G. and Boucher-Rodoni R. 1998. A phylogenetic study of the squid family Onychoteuthidae (Cephalopoda: Oegopsida). *Proceedings of the Royal Society of London, Series B*, **265**: 1761-1770.
- Buckman S. S. 1893. The Bajocian of the Sherborne District: its Relation to Subjacent and Superjacent Strata. *Quarterly Journal of the Geological Society*, **9**: 479-522.
- Cecca F. 1992. Ammonite habitats in the Early Tithonian of Western Tethys. *Lethaia*, **25**: 257-267.
- Chen C. S., Pierce G. J., Wang J., Robin J.-P., Poulard J.-C., Pereira J., Zuur A. F., Boyle P. R., Bailey N., Beare D. J., Jereb P., Ragonese S., Mannini A. and Orsi-Relini L. 2006. The apparent disappearance of *Loligo forbesi* from the south of its range in the 1990s: trends in *Loligo* spp. abundance in the northeast Atlantic and possible environmental influences. *Fisheries Research*, **78**: 44-54.
- Fernández-López S. and Meléndez G. 1996. Phylloceratina ammonoids in the Iberian Basin during the Middle Jurassic: a model of biogeographical and taphonomic dispersal related to relative sea-level changes. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **120**: 291-302.
- Gallardo C. S. and Penchaszadeh P. E. 2001. Hatching mode and latitude gradients in marine gastropods: revising Thorson's paradigm in the Southern hemisphere. *Marine Biology*, **138**: 547-552.
- Golikov A. N. and Scarlato O. A. 1967. Ecology of bottom biocoenoses in the Possjet Bay (the sea of Japan) and the peculiarities of their distribution in connection with physical and chemical conditions of the habitat. *Helgoländer Meeresunters*, **15**: 193-201.
- Gulyaev D. B. 2002. The first find of heteromorph ammonites in the Lower Callovian of European Russia. *Paleontological Journal*, **36**: 606-608.
- Gulyaev D. B. 2005. Stages of the ammonite fauna development during the early phases of the formation of the East-European sea basin (Late Bathonian – Early Callovian). *In: V. A. Zakharov, M. A. Rogov and O. S. Dzyuba (Eds), Materials of the first All-Russian Meeting "Jurassic System of Russia: problems of stratigraphy and paleogeography"*, 78-80, Geological Institute of RAS. (*in Russian*)
- Hantzpergue P., Baudin F., Mitta V., Olfieriev A. and Zakharov V. 1998. The Upper Jurassic of

- the Volga basin: ammonite biostratigraphy and occurrence of organic-carbon rich facies. Correlations between Boreal-Subboreal and Submediterranean provinces. *Mémoires du Muséum National d'Histoire Naturelle*, **179**: 9-33.
- Herke S. W. and Foltz D. W. 2002. Phylogeography of two squids (*Loligo pealei* and *L. plei*) in the Gulf of Mexico and northwestern Atlantic Ocean. *Marine Biology*, **140**: 103-115.
- Hiddink J. G. 2002. The adaptive value of migrations for the bivalve *Macoma baltica*. PhD. Thesis. Department of Marine Biology, Groningen, The Netherlands (available at link <http://www.bangor.ac.uk/~ossc06/infoJG.html>)
- Jereb P., Roper C. F. E. and Vecchione M. 2005. Introduction. In: P. Jereb and C. F. E. Roper (Eds), Cephalopods of the world. An annotated and illustrated catalogue of species known to date. Volume 1. Chambered nautilus and sepioids (Nautilidae, Sepiidae, Sepiolidae, Sepiariidae, Idiosepiidae and Spirulidae). *FAO Species Catalogue for Fishery Purposes*, **4**: 1-19.
- Kennedy W. J. and Cobban W. A. 1976. Aspects of ammonite biology, biogeography and biostratigraphy. *Special Papers in Paleontology*, **17**: 1-74.
- Keupp H. and Ilg A. 1992. Paläopathologie der Ammonitenfauna aus dem Obercallovium der Normandie und ihre palökologische Interpretation. *Berliner Geowissenschaftliche Abhandlungen, Reihe E*, **3**: 171-189.
- Kiselev D. N. 2003. Seltzo-Voskresenskoe. In: Atlas of the Yaroslavl region geological monuments, 58-62, Yaroslavl State Pedagogical University.
- Kiselev D. N. 2004. The thermal regime dynamics in Callovian-Oxfordian seas of Northwestern Eurasia: implications of relative paleotemperature data. *Stratigraphy and Geological Correlation*, **12** (4): 32-53.
- Kutek J. and Zeiss A. 1997. The highest Kimmeridgian and Lower Volgian in Central Poland; their ammonites and biostratigraphy. *Acta Geologica Polonica*, **47** (3-4): 107-198.
- Mesezhnikov M. S. 1984. Kimmeridgian and Volgian Stages of north of the USSR, 1-224, Nedra, Leningrad. (in Russian)
- Mitta V. V. 2003. On the Callovian-Oxfordian boundary beds of the Volga basin. *Vernadski Museum-Novitates*, **11**: 1-21 (in Russian).
- Mitta V. V. 2004 a. On the ammonite succession in the Jurassic-Cretaceous boundary beds of the Moscow Syncline. *Paleontological Journal*, **38**: 483-491.
- Mitta V. V. 2004 b. On the new publications on Jurassic ammonites and stratigraphy. *Bulletin of the Society of Naturalists of Moscow, Section of Geology*, **79** (1): 90-98 (in Russian).
- Mitta V. V., Barskov I. S., Gründel J., Zakharov V. A., Seltzer V. B., Ivanov A. V., Rostovtseva Yu. A. and Tarasova L. O. 2004. Upper Bajocian and Lower Bathonian in the section near Saratov. *Vernadski Museum-Novitates*, **12**: 1-39 (in Russian).
- Mitta V. V. and Seltzer V. B. 2002. First finds of Aretocephalitinae (Ammonoidea) in the Jurassic of the south-eastern Russian Platform, and the correlation of the Boreal Bathonian Stage with the standard scale. *Transactions of the Scientific Research Geological Institute of the N. G. Chernyshevskii Saratov State University. New Series*, **10**: 12-39 (in Russian).
- Navarro N., Neige P. and Marchand D. 2005. Faunal invasions as a source of morphological constraints and innovations? The diversification of the early Cardioceratidae (Ammonoidea; Middle Jurassic). *Paleobiology*, **31**: 98-116.
- Neige P. 2003. Spatial patterns of disparity and diversity of the Recent cuttlefishes (Cephalopoda) across the Old World. *Journal of Biogeography*, **30**: 1125-1137.
- Nesis K. N. 1985. Oceanic cephalopods: distribution, life forms, evolution, 1-285, Nauka, Moscow (in Russian).
- Nesis K. N. 1987. Cephalopods of the World. Squids, cuttlefishes, octopuses and allies. T. F. H. Publications Inc., Neptune City NJ, 351 pp.
- Nesis K. N. 2003. Distribution of Recent Cephalopoda and implications for Plio-Pleistocene events. *Berliner Paläobiologisches Abhandlungen*, **3**: 199-224.
- Neumayr M. 1878. Ueber untermittelt auftretende Cephalopodentypen im Jura Mitteleuropas. *Jahrbuch der Kaiserlich-Königlichen Geologischen Reichsanstalt*, **28** (1): 37-80.
- Olóriz F. 1990. Ammonite phenotypes and ammonite distributions. Notes and comments. In: G. Pallini, F. Cecca, S. Cresta and M. Santantonio (Eds), Fossili, Evoluzione, Ambiente. *Atti II Conv. Int. F.E.A. Pergola, 1987*: 417-426.
- Oppel A. 1856-1858. Die Juraformation Englands, Frankreichs und des südwestlichen Deutschlands. *Jahresberichte Jahreshefte des Vereins für Vaterländische Naturkunde in Württemberg*, **12-14**: 1-857.
- Page K. N. 1995. Biohorizons and zonules: infra-subzonal units in Jurassic ammonite stratigraphy. *Palaeontology*, **38**: 801-814.

- Pörtner H. O. 2001. Climate change and temperature-dependent biogeography: oxygen limitation of thermal tolerance in animals. *Naturwissenschaften*, **88**: 137–146.
- Rawson P. F. 1973. Lower Cretaceous (Ryazanian-Barremian) marine connections and cephalopod migrations between the Tethyan and Boreal realms. *Geological Journal Special Issue*, **5**: 131-144.
- Reyment R. A. 2005. Active migration and passive transport of marine organisms in the fossil record. *In*: A. M. T. Elewa (Ed.), Migration of organisms. Climate, geography, ecology, 7-24. Springer.
- Rogov M. A. 2004. The Russian Platform as a key region for Volgian/Tithonian correlation: A review of the Mediterranean faunal elements and ammonite biostratigraphy of the Volgian stage. *Rivista Italiana di Paleontologia e Stratigrafia*, **110** (1): 321-328.
- Sachs V. N., Basov V. A., Dagens A. A., Dagens A. S., Zakharov V. A., Ivanova E. F., Meledina S. V., Mesezhnikov M. S., Nalnyaeva T. I. and Shulgina N. I. 1971. Paleozoogeography of the seas of the Boreal Belt in the Jurassic and Neocomian, *In*: Problems of general and regional geology, 179-211, Nauka, Novosibirsk. (*in Russian*)
- Salman A., Laptikhovskiy V. and Katagan T. 2005. Male and female fecundity of the Indo-Pacific octopus, *Octopus kagoshimensis* Ortmann, 1888 (Cephalopoda: Octopodidae) in the East Mediterranean. *Zoologicheskii Zhurnal*, **84** (2): 269-271. (*in Russian*)
- Scarlato O. A. 1981. Bivalve mollusks of the temperate latitudes of the western part of the Pacific Ocean. 1-479, Nauka, Leningrad. (*in Russian*)
- Schweigert G. 2000. Immigration of amoeboceratids into the Submediterranean Upper Jurassic of SW Germany. *In*: R. L. Hall and P. L. Smith (Eds), Advances in Jurassic Research 2000. *GeoResearch Forum*, **6**: 203-209. TransTech Publishing.
- Sey I. I. and Kalacheva E. D. 1983. On invasions of Tethyan ammonites in Boreal Late Jurassic basins of the East of USSR. *In*: V. A. Zakharov and T. I. Nalnyaeva (Eds), Mesozoic of the Soviet Arctic, 61-74, Nauka, Novosibirsk (*in Russian*).
- Thierry J. and Barrier E. 2000. Early Tithonian. *In*: J. Dercourt, M. Gaetani, B. Vrielynck, E. Barrier, B. Biju-Duval, M. F. Brunet, J. P. Cadet, S. Crasquin and M. Sandulescu (Eds), Atlas Peri-Tethys. Palaeogeographical maps, Paris. Commission de la Carte Géologique du Monde/Commission for the Geological Map of the World.
- Uhlig V. 1911. Die marinen Reiche des Jura und der Unterkreide. *Mitteilungen der Geologisches Gesellschaft Wien*, **4** (3): 329-448.
- Westermann G. E. G. 1996. Ammonoid life and habitat. *In*: N. Landman, K. Tanabe and R. A. Davis (Eds), Ammonoid paleobiology, Topics in geobiology, **13**: 607-707. N.-Y, Plenum Press.
- Westermann G. E. G. 2000. Biochore classification and nomenclature in paleobiogeography: an attempt at order. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **158**: 1-13.
- Westermann G. E. G. and Tsujita C. J. 1999. Life habits of ammonoids. *In*: E. Savazzi (Ed.), Functional morphology of the invertebrate skeleton, 299-325. London, John Wiley & Sons Ltd.
- Zakharov V. A. and Rogov M. A. 2004. The Boreal-Tethyan biogeographical mollusk ecotone in Europe during the Jurassic-Cretaceous transition. *Rivista Italiana di Paleontologia e Stratigrafia*, **110**, 1: 339-444.
- Zakharov V. A. and Kasumzadeh A. A. 2005. On the Boreal genus *Buchia* (Bivalvia) from the Tithonian of the Lesser Caucasus. *Stratigraphy and Geological Correlation*, **13**, 6: 606-612.
- Zakharov V. A., Baudin F., Dzyuba O. S., Daux V., Zverev K. V. and Renard M. 2005. Isotopic and faunal record of high paleotemperatures in the Kimmeridgian of Subpolar Urals. *Russian Geology and Geophysics*, **46**: 1-19.
- Zakharov Y. D., Smyshlyaeva O. P., Shigeta Y., Popov A. M. and Zonova T. D. 2006. New data on isotopic composition of Jurassic – Early Cretaceous cephalopods. *Progress in Natural Science*, **16** (13): 50-67.
- Ziegler B. 1967. Ammoniten-Ökologie am Beispiel des Oberjura. *Geologische Rundschau*, **56** (2): 439-464.