



New approach to stratigraphy of palaeolake and glacial sediments of the younger Middle Pleistocene in mid-eastern Poland

Leszek LINDNER, Leszek MARKS



Lindner L., Marks L. (1999) — New approach to stratigraphy of palaeolake and glacial sediments of the younger Middle Pleistocene in mid-eastern Poland. *Geol. Quart.*, 43 (1): 1–8. Warszawa.

Basing on 10 key sections from mid-eastern Poland, the three interglacials (Mazovian, Zbójnian and Lubavian, which follow the Sanian 2 (Wilgian) Glaciation and precede the Wartanian, are separated from one another by two glacial episodes. The older is the Liviecian Glaciation and the younger is the Krznanian Glaciation (named after the River Krzna). The Krznanian Glaciation occupies a stratigraphic location of the previous Odranian Glaciation. The latter is postulated to be incorporated into the Wartanian Glaciation as the maximum stadial. Such stratigraphic subdivision of the younger Middle Pleistocene of Poland indicates a very close relationship to the most recent stratigraphic schemes of glacial and interglacial units in Germany and in the Russian Plain.

Leszek Lindner, Institute of Geology, Warsaw University, Żwirki i Wigury 93, 02-089 Warszawa, Poland; Leszek Marks, Polish Geological Institute, Rakowiecka 4, 00-975 Warszawa, Poland, e-mail: lmar@pgi.waw.pl (received: November 19, 1998; accepted: December 30, 1998).

Key words: mid-eastern Poland, Middle Pleistocene, Mazovian Interglacial, Krznanian Glaciation, stratigraphy, palaeogeography.

INTRODUCTION

The paper presents a new approach to stratigraphy of the younger Middle Pleistocene of Poland, particularly to a number and age of the Middle Polish Glaciations (Saalian). This idea is based on denial of the Eemian age of sediments of the so-called Sztum Sea (*cf.* A. Makowska, 1994), as well as on numerous new evidence of a more southern limit of a till, geochronological and lithological parameters of which are typical for the Wartanian Glaciation (among others J. Nitychoruk, 1994; L. Marks *et al.*, 1995; L. Lindner, 1996, 1998; L. Lindner, S. Fedorowicz, 1996; B. Jaśkowski, B. J. Kowalski, 1997, 1998; S. Lisicki, 1998a). Significant role in this stratigraphic discussion has been played by new sites of the Zbójnian Interglacial in Poland (*cf.* P. Woźniak, 1989; Z. Janczyk-Kopikowa, 1991, 1998; T. Kuszell, 1997). All these facts supplement the previous conclusions, drawn from correlation of individual units of the Saalian Glaciation in mid-western Europe (*cf.* L. Marks, 1991), and the more recent opinions on evidence for four interglacials after the Sanian 2 Glaciation (Elsterian 2) in Europe (*cf.* L. Lindner, B. Marciński, 1998).

YOUNGER MIDDLE PLEISTOCENE

Limits of the Middle Pleistocene are accepted in this paper by D. Q. Bowen (1978), i.e. the boundary Matuyama/Brunhes at the base (about 700 ka) and a beginning of the Eemian Interglacial at the top (about 128 ka). The younger Middle Pleistocene begins with the Sanian 2 Glaciation (=Wilgian) and terminates with the Wartanian Glaciation. Description of this interval is based among others on analysis of 10 sections of the Quaternary deposits (Grabówka, Nowiny, Goleń, Rączki Wielkie, Losy, Nidzica, Krępiec, Ossówka, Zbójno and Bedlno) from mid-eastern Poland (Figs. 1 and 2).

SANIAN 2 GLACIATION

Ice sheet of the Sanian 2 Glaciation, correlated with the oxygen isotope stage 12, reached northern slopes of the Carpathians and the Sudetes (L. Lindner, L. Marks, 1995). It deposited commonly a single or two tills. In the section Goleń (Figs. 1 and 2), between these tills there are lake sediments with a fragmentary pollen succession (*cf.* H. Winter, S. Lisi-




Fig. 1. Location of the study area in mid-eastern Poland

1 — main sections of interglacial sediments; maximum limit of ice sheets during the Scandinavian glaciations; 2 — Liviecian, 3 — Krznanian, 4 — Wartanian (Odranian Stadial), 5 — Wartanian (Wartanian s.s. Stadial), 6 — Vistulian; 7 — Late Glacial and Holocene river valleys

cki, 1998), components and location of which speak for a warm interstadial or a new interglacial, named recently the Mrongovian (cf. H. Winter, 1998; S. Lisicki, 1998b). Till of the Sanian 2 Glaciation occurs also at Krępiec (M. Harasimiuk, A. Henkiel, 1981; M. Harasimiuk *et al.*, 1988). According to L. Lindner (1992), it represents the youngest of the South Polish Glaciations (Fig. 2).

MAZOVIAN INTERGLACIAL

It is the oldest climatic warming within the so-called Great Interglacial (Fig. 2) according to S. Z. Różycki (1964, 1972), and is among the better examined climatostratigraphic units of the Pleistocene of Poland. In the presented area it occurs commonly, among others in the sections of lake sediments at Krępiec, Goleń and Ossówka (Figs. 1 and 2). Palynological analysis of the section Krępiec proved (cf. Z. Janczyk-Kopikowa, 1981) that deposition occurred during four vegetation periods, a climatic optimum during the third one with *Vitis*, and predominating *Carpinus* and *Abies*. Diatom analysis of these sediments indicates (cf. B. Marciniak, 1980, 1983) prevalence of nannoplanktonic diatoms in the bottom, partly replaced in the middle by littoral and reophilic species, and in

the top by specimens, typical for shallowing and overgrowing of the lake with macrophytes. From a biostratigraphic point of view, these sediments are to be easily correlated with deposits of the Holsteinian Interglacial in Germany (cf. B. Urban, 1997) and the Likhvinian Interglacial in the Russian Plain (cf. N. S. Bolikhovskaya, N. G. Sudakova, 1996). They correspond to the oxygen isotope stage 11 in deep-sea sediments (cf. L. Lindner, B. Marciniak, 1998).

In the section Goleń, this interglacial is represented (H. Winter, S. Lisicki, 1998) by lake sediments, underlain by a till with petrographic coefficients typical for the Liviecian(?) Glaciation and covered by a till of the Odranian Glaciation. In the Lower Vistula Valley, bottom parts of tills of the Middle Polish Glaciations contain glacial rafts with marine sediments of the Mazovian (= Holsteinian) Interglacial (cf. A. Makowska, 1986). They are mainly sands and silts with pieces of *Cardium* sp., *C. echinatum*, *C. edule*, as well as forams of the Holstein Sea that undoubtedly occurred in the southern part of the Baltic Basin (cf. O. Kondratienė, V. Gudelis, 1983; L. Marks, 1988, 1994).

Fluvial pattern of the Mazovian Interglacial in Poland has been considered to be reflected by relief of the Quaternary substrate (S. Z. Różycki, 1965; E. Rühle, 1955, 1967, 1973). They were L. Lindner *et al.* (1982) who postulated a present-like fluvial pattern of the Mazovian Interglacial. Lack of reliable dating methods makes the assumed water level of the Holstein Sea to be the most important index for reconstruction of a fluvial pattern of the Mazovian Interglacial. According to the studies in the Kaliningrad District of Russia and in the Hamburg region of northwestern Germany, this level is expected to be close to a sea level of the present Baltic Sea (cf. L. Marks, 1994, 1995). Therefore, outside the areas with undoubted glaciotectionic deformations or neotectonic movements, beds of the buried river valleys of the Mazovian Interglacial are to be located at similar altitudes as the beds of contemporary rivers in Central Europe (cf. Fig. 3).

LIVIECIAN GLACIATION

The Liviecian Glaciation in mid-eastern and northeastern Poland is represented by a separate till (L. Lindner, L. Marks, 1995; S. Lisicki, 1997), preserved among others in the section Raczki Wielkie (Figs. 1 and 2), and possibly also as a lowest till at Grabówka (Figs. 1 and 2). In other sections this till is absent (cf. Fig. 2), presumably due to glacial erosion. Close to its southernmost limit, a till of this glaciation was found in the vicinity of Koziernice and Dęblin, central Poland (M. Żarski, 1994). In the section Zbójno there are only silts, deposited in the extraglacial zone (L. Lindner, E. Brykczyńska, 1980). Geological setting of sediments of the Liviecian Glaciation suggests their correlation with the oxygen isotope stage 10 (cf. L. Lindner, L. Marks, 1995).

ZBÓJNIAN INTERGLACIAL

This interglacial is a successively younger, warm climatostratigraphic unit within the Great Interglacial (Fig. 2). In the presented area, its lake sediments are known from the

section Zbójno and presumably also from the section Raczki Wielkie (Figs. 1 i 2). In the section Zbójno, four periods of vegetation development were distinguished, with climatic optimum during the second period when drop of *Pinus* was accompanied at first by rapid development of *Tilia* (to 48%) and then, drop of *Tilia* (to 20%) with rise of *Alnus*, *Carpinus*, *Picea* and *Corylus*, accompanied by *Quercus* (L. Lindner, E. Brykczyńska, 1980). These sediments are represented by peat on silts of the Liviecian Glaciation and the underlying alluvia of the Mazovian Interglacial. They are covered by silts, sands and a till of the younger glaciation (L. Lindner, B. Marciniak, 1998). In the section Raczki Wielkie, deposits of this interglacial are presumably composed of two layers of peat, a pollen analysis of which was done by Z. Janczyk-Kopikowa (cf. P. Woźniak, 1989). The Zbójnian Interglacial is generally correlated with the oxygen isotope stage 9 (cf. L. Lindner, B. Marciniak, 1998).

KRZANIAN GLACIATION

Geological data on the Pleistocene of mid-eastern Poland prove that the first post-Zbójnian glacial sediments are located at stratigraphic position of the previously distinguished Odranian Glaciation (Fig. 4). These deposits are, however, less widespread if compared with extents of the hitherto distinguished Middle Polish Glaciations (i.e. Odranian and Wartanian). They are connected with the Scandinavian ice sheet that presumably passed across the meridional reaches of the Pilica and Wieprz Rivers, as far south as the Krzna River (Krzna Stadial after E. Rühle, 1970) and therefore, should be named the Krznanian Glaciation. It could, however, even reach the northern slopes of the South Polish Uplands and had been miscorrelated with the Końskie (pre-maximum) Phase of the Odranian Glaciation (cf. L. Lindner, 1971, 1995, 1998; L. Lindner, S. Fedorowicz, 1996; L. Lindner, J. Semil, in print). In the whole mid-eastern Poland, the Krznanian Glaciation is represented by a till, preserved in most sections (Figs. 1 and 2). At Grabówka, this till presumably underlies the interglacial lake series, examined by A. Makowska (1977). At Nowiny, it is overlain by varved clays and silts that pass gradually upwards into sediments of the so-called Sztum Sea (cf. A. Makowska, 1979, 1986). This till occurs presumably also at Raczki Wielkie (P. Woźniak, 1989), whereas at Losy it is probably replaced by sands and gravels beneath sediments of the Lubavian Interglacial (cf. K. M. Krupiński, L. Marks, 1986). Thus, the Krznanian Glaciation seems to correspond to the oxygen isotope stage 8, i.e. to this global cooling which was previously correlated with the Odranian Glaciation (cf. L. Lindner, L. Marks, 1995).

LUBAVIAN INTERGLACIAL

The Lubavian Interglacial is best represented by a lake marl in the section Losy near Lubawa (Figs. 1 and 2), geological setting and pollen diagram of which were examined by K. M. Krupiński and L. Marks (1985, 1986). The lake sediments from Grabówka, examined in a few samples only by Z.




Fig. 2. Stratigraphic-palaeogeographic scheme of glaciations and interglacials during the younger Middle Pleistocene and the Late Pleistocene of Poland

1 — main tills; 2 — interglacial sea transgressions; 3 — probable interglacial sea ingressions (on the basis of re deposited marine sediments); 4 — section fragments with palynologic analysis; 5 — section fragments with diatom analysis; interglacial sites: Grabówka (A. Makowska, 1977), Nowiny (A. Makowska, 1986), Goleń (H. Winter, S. Lisicki, 1998), Raczki Wielkie (P. Woźniak, 1989), Losy (K. M. Krupiński, L. Marks, 1986), Nidzica (Z. Michalska in: B. Marciniak, W. W. Kowalski, 1978), Krępiec (M. Harasimiuk, A. Henkiel, 1981), Ossówka (B. Marciniak in: L. Lindner *et al.*, 1990; K. M. Krupiński, 1995), Zbójno (L. Lindner, E. Brykczyńska, 1980), Bedlno (A. Środoń, M. Gołąbowa, 1956)

Janczyk-Kopikowa (cf. A. Makowska, 1977), are presumably of the same age. The same interglacial seems to be represented by marine sediments with Lusitanian mollusc shells (cf. A. Makowska, 1986) of the so-called Sztum Sea at Nowiny (Figs. 1 and 2). At Losy, four periods of vegetation development were distinguished. The first one was predominated by birch shrubs or forest with *Pinus* and *Larix*. *Picea*, *Quercus* and *Ulmus* were also present during the second, and *Tilia* during the third period. The fourth period indicates development of *Corylus*, accompanied by *Ulmus*, *Alnus* and *Tilia*. Diatoms in these deposits are different from the typical ones of the Eemian (B. Marciniak, pers. inform.). Deposits of the Lubavian Interglacial at Losy are overlain by glaciofluvial sands with gravel and till of the Wartanian Glaciation (K. M. Krupiński, L. Marks, 1986). The lake sediments at Losy represent a warming which presumably corresponds to the oxygen isotope stage 7 (cf. L. Lindner, B. Marciniak, 1998).




Fig. 3. Longitudinal section of the Middle Vistula valley from Sandomierz to Warsaw

WARTANIAN GLACIATION

Sediments of this glaciation comprise presumably the pre-maximum (Odranian?) and the maximum stadial (Wartanian *s.s.*), and the till occurs commonly in the presented sections Grabówka, Goleń, Raczki Wielkie and Losy (Figs. 1 and 2). In the Bechatów region, two tills of this age are

interbedded with sands (*cf.* L. Lindner, 1984). Recent examination of this till in Poland suggests further southward limit of the ice sheet during the Wartanian Glaciation or even its correspondence to the maximum (or close to maximum) limit of the Scandinavian ice sheet during the Middle Polish Glaciations (*cf.* L. Marks *et al.*, 1995; L. Lindner, S. Fedorowicz, 1996; B. Jaśkowski, B. J. Kowalski, 1997, 1998). Basing on petrographic composition of tills in the area occupied by the ice sheet of the Wartanian Glaciation and in its forefield, a traditional concept of the ice sheet limit in that time seems questionable (*cf.* J. Winnicki, 1997; S. Lisicki, 1998a). The Wartanian Glaciation corresponds to the oxygen isotope stage 7 (*cf.* L. Lindner, L. Marks, 1995).




Fig. 4. Possible variants of correlation of the younger Middle Pleistocene in Poland

FINAL REMARKS

The presented facts postulate a considerable revision of predominant concepts (e.g. L. Lindner, K. Grzybowski, 1982; L. Lindner, 1992; J. E. Mojski, 1993) of the Middle Polish Glaciations in Poland (Fig. 2). They suggest also a more reliable correlation of the main younger Middle Pleistocene climatostratigraphic units in Poland with similar units in Germany and the Russian Plain (Fig. 5).

Therefore, the recently distinguished two new interglacials, i.e. Reinsdorf and Schöningen (Fig. 5) at Schöningen in central Germany (*cf.* B. Urban, 1995, 1997), should be, however, considered for equivalents of the Zbójnian and the Lubavian (*cf.* L. Lindner, B. Marciniak, 1998), although they are not separated by tills from each other and from the prece-

ding Holsteinian Interglacial. At Schöningen, the lake sediments of the Holsteinian Interglacial, together with the preceding and the following stadials and interstadials, are underlain by a till of the Elsterian 2 Glaciation. Deposits of the youngest pre-Saalian Interglacial in this section are overlain by a till of the Drenthe Stage, mantled in turn with extraglacial(?) deposits of the Warthe Stage and by the Eemian series (B. Urban, 1997).

In the section Chekalin of the Russian Plain (cf. N. S. Bolikhovskaya, N. G. Sudakova, 1996), there are also two new interglacials examined recently i.e. Chekalinian and Cherepietskian, younger than the Likhvinian Interglacial and also without any separation by glacial deposits. Deposits of the Likhvinian Interglacial in this section are underlain by a till of the Okanian Glaciation and this in turn, by deposits of the Byelovezhian Interglacial (*op. cit.*). The mentioned three younger interglacials in this section are separated by deposits of the Kaluganian and Zhizdrinskian climatic coolings (Fig. 5). Deposits of the youngest interglacial (Cherepietskian) are covered by two tills of the Dnieperian Glaciation (N. S. Bolikhovskaya, N. G. Sudakova, 1996).

All these data suggest considerable similarity in a number and climatostratigraphic rank of units of the younger Middle Pleistocene in Germany, Poland and the Russian Plain. Both in Germany and in the Russian Plain, these units remarkably resemble a sequential warming and coolings which were identified in Poland as the Great Interglacial *sensu* S. Z. Różycki (1964, 1972) or the Mazovian Interglacial *s.l.* of E. Rühle (1973). A concept of this rhythm seems to be still acceptable for the territory of Poland but only in the case of buried depressions, influenced by neotectonic processes and fluvial activity (M. D. Baraniecka, 1975; D. Krzyszkowski, 1991; W. Pożaryski *et al.*, 1994). At the ancient morainic plateaux in Poland, there survived tills that separate deposits of the Mazovian, Zbójnian and Lubavian Interglacials, suggesting ice sheet advance at least twice, i.e. during the Liviecian and the Krznanian Glaciations. These advances could be possible due to the most favourable conditions for ice sheet development to the south of the Baltic Sea.

AGE	OXYGEN ISOTOPE STAGES (D.Q. Bowen <i>et al.</i> , 1986)	GERMANY	POLAND	RUSSIAN PLAIN	
		(B. Urban, 1997)		(N. S. Bolikhovskaya, N. G. Sudakova, 1996)	
QUATERNARY	HOL.	1	HOLOCENE	HOLOCENE	
	UPPER PLEISTOCENE	2-5d	WEICHSELIAN	VISTULIAN	VALDAYNIAN
5e		EEMIAN	EEMIAN	MIKULINIAN	
MIDDLE PLEISTOCENE	6	WARTHE ORENTHE	SAALE COMPLEX	WARTANIAN ODRANIAN	MOSCOWIAN DNEIPERIAN
	7	SCHÖNINGEN		LUBAVIAN	CHEREPIETSKIAN
	8	?		KRZANIAN	ZHIZDRINSKIAN
	9	REINSDORF		ZBÓJNIAN	CHEKALINIAN
	10	?	LIVIECIAN	MIDDLE POLISH GLACIATIONS	KALUGANIAN
	11	HOLSTEINIAN	MAZOVIAN		LIKHVINIAN
	12	ELSTERIAN 2	SANIAN 2		OKANIAN

Fig. 5. Correlation of the main climatostratigraphic units of the Late Quaternary in mid-eastern Europe

Sediments of the Schöningen Interglacial are covered by a till of the Drenthe Stage (B. Urban, 1995), and sediments of the Cherepietskian Interglacial by two tills of the Dnieperian Glaciation (N. S. Bolikhovskaya, N. G. Sudakova, 1996). This fact results in two conclusions. The first one is that the glacial episode, distinguished in Poland at the traditional position of the Odranian Glaciation, i.e. between the interglacials Zbójnian and Lubavian, is presumably older than the Drenthe Stage in Germany and the Dnieperian Glaciation in the Russian Plain. In this paper, the name of the Krznanian Glaciation is proposed for this glacial event (Figs. 4 and 5). The second conclusion is a proposal to accept the previous limit of ice sheet of the Odranian Glaciation as of the maximum stadial of the Wartanian Glaciation.

REFERENCES

BARANIECKA M. D. (1975) — The dependences of the development of Quaternary deposits upon the structure and dynamics of the basement in the central part of the Polish Lowlands (in Polish with English summary). *Biul. Inst. Geol.*, **288**: 5–97.

BOLIKHOVSKAYA N. S., SUDAKOVA N. G. (1996) — Stratigraphic and correlation significance of the Chekalin (Likhvin) key section of the Pleistocene of the Russian Plain (in Russian). *Stratigraphy, Geological Correlation* **4** (3): 88–99. Moscow.

BOWEN D. Q. (1978) — Quaternary geology: a stratigraphic framework for multidisciplinary work. Pergamon Press. Oxford.

BOWEN D. Q., RICHMOND G. M., FULLERTON D. S., ŠIBRAVA V., FULTON R. J., VELICHKO V. V. (1986) — Correlation of Quaternary glaciations in the Northern Hemisphere. In: Quaternary glaciations in the Northern Hemisphere (eds. V. Šibrava *et al.*). *Quatern. Sc. Rev.*, **5**: 509–510.

HARASIMIUK M., HENKIEL A. (1981) — Fossil valley forms in the vicinities of Łęczna and their importance for the palaeogeography of the Wieprz river drainage system (in Polish with English summary). *Kwart. Geol.*, **25** (1): 147–162.

HARASIMIUK M., MARUSZCZAK H., WOJTANOWICZ J. (1988) — Quaternary stratigraphy of the Lublin region, southeastern Poland. *Quatern. Stud.*, **8**: 15–25.

JANCZYK-KOPIKOWA Z. (1981) — Pollen analysis of the Pleistocene sediments at Kaznów and Krępiec (in Polish with English summary). *Biul. Inst. Geol.*, **321**: 249–258.

JANCZYK-KOPIKOWA Z. (1991) — Problems of the palynostratigraphy of the Pleistocene in Poland and the palynological analysis of interglacial deposits from Biesiekierz, Central Poland (in Polish with English summary). *Ann. UMCS, Sec. B*, **46**, suppl. 1.

JANCZYK-KOPIKOWA Z. (1998) — Sukcesja pyłkowa typu Zbójnia w dotychczasowych publikacjach. *Mat. V Konf. „Stratygrafia plejstocenu Polski”* (eds. S. Lisicki *et al.*): 30. Iznota.

JAŚKOWSKI B., KOWALSKI B. J. (1997) — Problem zasięgu lądolodu zlodowacenia Warty w profilu Grójec–Szydłowiec. In: *Problemy zlodowacenia*

- waceń środkowopolskich w Polsce południowo-zachodniej (eds. D. Krzyszkowski, B. Przybylski). IV Konf. „Stratygrafia plejstocenu Polski”: 211–212. Wrocław.
- JĄSKOWSKI B., KOWALSKI B. J. (1998) — Extent of Warta glacier in the Grójec–Szydłowiec area in the light of TL examination of the glacial tills, Central Poland (in Polish with English summary). *Prz. Geol.*, **46** (4): 355–358.
- KONDRATIENE O., GUDELIS V. (1983) — Pleistocene marine sediments in the Peribaltica area (in Polish). *Prz. Geol.*, **31** (8–9): 497–502.
- KRUPIŃSKI K. M. (1995) — Pollen stratigraphy and succession of vegetation during the Mazovian Interglacial (in Polish with English summary). *Acta Geogr. Lodz.*, **70**.
- KRUPIŃSKI K. M., MARKS L. (1985) — Interglacial site at Losy near Lubawa in the Mazury Lakeland, preliminary report (in Polish with English summary). *Kwart. Geol.*, **29** (3–4): 767–779.
- KRUPIŃSKI K. M., MARKS L. (1986) — Interglacial sediments at Losy, Mazury Lakeland. *Bull. Pol. Acad. Sc., Earth Sc.*, **34** (4): 375–386.
- KRZYSZKOWSKI D. (1991) — The middle Pleistocene polyinterglacial Czyżów Formation in the Kleszczów Graben (Central Poland): stratigraphy and palaeogeography. *Folia Quatern.*, **61–62**: 5–58.
- KUSZELL T. (1997) — Nowe stanowiska osadów interglacjalnych i wistulianu w południowo-zachodniej Polsce. In: *Problemy zlodowaceń środkowopolskich w Polsce południowo-zachodniej* (eds. D. Krzyszkowski, B. Przybylski). *Mat. IV Konf. „Stratygrafia plejstocenu Polski”*: 181–182. Wrocław.
- LINDNER L. (1971) — Pleistocene stratigraphy and palaeogeomorphology of the north-western margin of the Holy Cross Mountains, Poland (in Polish with English summary). *Studia Geol. Pol.*, **35**.
- LINDNER L. (1984) — An outline of Pleistocene chronostratigraphy in Poland. *Acta Geol. Pol.*, **34** (1–2): 27–49.
- LINDNER L. (1988) — Stratigraphy and extents of Pleistocene continental glaciations in Europe. *Acta Geol. Pol.*, **38** (1–4): 63–83.
- LINDNER L. (1992) — Stratygrafia (klimatostratygrafia) czwartorzędu. In: *Czwartorzęd: osady, metody badań, stratygrafia* (ed. L. Lindner): 441–633. Wyd. PAE. Warszawa.
- LINDNER L. (1995) — Till sequences and local moraines in the Holy Cross Mountains area in Central Poland. In: *Glacial deposits in North-East Europe* (eds. J. Ehlers *et al.*): 329–337. A. A. Balkema. Rotterdam.
- LINDNER L. (1996) — Post-Mazovian glacial and glacialfluvial sediments in the Biała Podlaska region and their age in light of discussion on the Warta Glaciation in Poland (in Polish with English summary). *Biul. Państw. Inst. Geol.*, **373**: 87–96.
- LINDNER L. (1998) — Main relief components of Giełniów Height and their influence on the ice extent during the Odra (Warta?) Glaciation in the NW peripheral zone of Holy Cross Mountains (in Polish with English summary). In: *Główne kierunki badań geomorfologicznych w Polsce — stan aktualny i perspektywy* (ed. K. Pękała): 343–347. *Referaty i Komunikaty IV Zjazdu Geomorfologów Polskich*. Lublin.
- LINDNER L., BRYKCYŃSKA E. (1980) — Organogenic deposits at Zbójno by Przedbórz, western slopes of the Holy Cross Mts., and their bearing on stratigraphy of the Pleistocene of Poland. *Acta Geol. Pol.*, **30** (2): 155–163.
- LINDNER L., FEDOROWICZ S. (1996) — TL age of Pleistocene deposits in Janów on the Radomka River (Central Poland) and the problem of extent of continental glaciers of Middle Polish Glaciations NW of the Holy Cross Mts., Central Poland (in Polish). *Prz. Geol.*, **44** (9): 935–937.
- LINDNER L., GRZYBOWSKI K. (1982) — Middle Polish glaciations (Odranian, Wartanian) in southern Central Poland. *Acta Geol. Pol.*, **32** (3–4): p. 191–208.
- LINDNER L., MARCINIĄK B. (1998) — The occurrence of four interglacials younger than the Sanian 2 (Elsterian 2) Glaciation in the Pleistocene of Europe. *Acta Geol. Pol.*, **48** (3): 247–264.
- LINDNER L., MARKS L. (1995) — Outline of palaeogeomorphology of the Polish territory during the Scandinavian glaciation (in Polish with English summary). *Prz. Geol.*, **43** (7): 591–594.
- LINDNER L., SEMIL J. (in print) — Morphology and Quaternary sediments within the area of Giełniów Elevation, NW margin of the Holy Cross Mts. (in Polish with English summary). *Pr. Inst. Geogr. WSP w Kielcach*, **4**.
- LINDNER L., LAMPARSKI Z., DĄBROWSKI S. (1982) — River valleys of the Mazovian Interglacial in eastern Europe. *Acta Geol. Pol.*, **32** (3–4): 179–190.
- LINDNER L., KRUPIŃSKI K. M., MARCINIĄK B., NITYCHORUK J. (1990) — Geological and floristic situation of Pleistocene organogenic deposits in the Ossówka region, Southern Podlasie (in Polish with English summary). *Prz. Geol.*, **38** (11): 476–483.
- LISICKI S. (1997) — Pleistocene of Mragowo Lakeland. *Geol. Quart.*, **41** (3): 327–346.
- LISICKI S. (1998a) — Interpretation results of petrographic analysis of gravel fraction in glacial till with reference to its genesis (in Polish with English summary). *Prz. Geol.*, **46** (5): 410–416.
- LISICKI S. (1998b) — Nowa interpretacja stratygrafii plejstocenu w wybranych profilach Polski północno-wschodniej. *Mat. V Konf. „Stratygrafia plejstocenu Polski”* (eds. S. Lisicki *et al.*): 5–6. *Iznota*.
- MAKOWSKA A. (1977) — Interglacial horizon in sections of the Middle Polish Glaciation from the Lower Vistula River (in Polish with English summary). *Kwart. Geol.*, **21** (4): 769–787.
- MAKOWSKA A. (1979) — Eemian Interglacial in valley of Lower Vistula River (in Polish with English summary). *Studia Geol. Pol.*, **63**.
- MAKOWSKA A. (1986) — Pleistocene seas in Poland — sediments, age and palaeogeography (in Polish with English summary). *Pr. Inst. Geol.*, **120**.
- MAKOWSKA A. (1994) — Climatic variation in the pre-glacial part of the Toruń Glaciation in the Lower Vistula Region and the Elbląg Elevation (preliminary report). *Geol. Quart.*, **38** (1): 133–154.
- MARCINIĄK B. (1980) — Middle Pleistocene diatoms from lacustrine deposits from Krępiec (in Polish with English summary). *Kwart. Geol.*, **24** (2): 349–360.
- MARCINIĄK B. (1983) — Diatoms in the Mazovian Interglacial of the Lublin Upland. *Bull. Pol. Acad. Sc. Ser. Terre*, **30** (1): 77–85.
- MARCINIĄK B., KOWALSKI W. W. (1978) — Dominant diatoms, pollen, chemistry and mineralogy of the Eemian lacustrine sediments from Nidzica (North Poland): a preliminary report. *Pol. Arch. Hydrobiol.*, **25** (2): 269–281.
- MARKS L. (1988) — Relation of substrate to the Quaternary paleorelief and sediments, western Mazury and Warmia (Northern Poland). *Zesz. Nauk AGH, 1165, Geol. Kwart.*, **14** (1).
- MARKS L. (1991) — Attempt of correlation of the Saale Glaciation in Central European Lowland. *Bull. Pol. Acad. Sc. Earth Sc.*, **39** (2): 187–198.
- MARKS L. (1994) — Mid-Pleistocene ice-dam sediments in the Lower Vistula Region (in Polish with English summary). *Acta Univ. Nicolai Copernici, Geografia*, **27**: 225–232.
- MARKS L. (1995) — Correlation of the middle Pleistocene ice-dam lacustrine sediments in the Lower Vistula and the Lower Elbe regions. *Acta Geol. Pol.*, **45** (1–2): 143–152.
- MARKS L., LINDNER L., NITYCHORUK J. (1995) — New approach to a stratigraphic position of the Warta Stage in Poland. *Acta Geogr. Lodz.*, **68**: 135–147.
- MOJSKI J. E. (1993) — Europa w plejstocenic; ewolucja środowiska przyrodniczego. *Wyd. PAE. Warszawa*.
- MOJSKI J. E. (1995) — Pleistocene glacial events in Poland. In: *Glacial deposits in North-East Europe* (eds. J. Ehlers *et al.*): 287–292. A. A. Balkema. Rotterdam.
- NITYCHORUK J. (1994) — Stratygrafia plejstocenu i paleogeomorfologia południowego Podlasia. *Roczn. Międzyrzecki*, **26**: 23–107.
- POŻARYSKI W., MARUSZCZAK H., LINDNER L. (1994) — Chronostratigraphy of Pleistocene deposits and evolution of the middle Vistula River valley with particular attention to the gap through the South Polish Uplands (in Polish with English summary). *Pr. Państw. Inst. Geol.*, **47**.
- RÓŻYCKI S. Z. (1964) — Les oscillations climatiques pendant le Grand Interglaciaire. *Report of the Vth INQUA Congress*, **2**: 211–225. Łódź.
- RÓŻYCKI S. Z. (1965) — Traits principaux de la stratigraphie et de la paléomorphologie de la Pologne pendant le Quaternaire. *Report of the Vth INQUA Congress*, **1**: 123–142. Łódź.
- RÓŻYCKI S. Z. (1972) — Plejstocen Polski Środkowej na tle przeszłości w górnym trzeciorzędzie. *PWN. Warszawa*.
- RÜHLE E. (1955) — Review of the data concerning the substratum of the Quaternary in the north-eastern part of the Polish Lowland (in Polish with English summary). *Biul. Inst. Geol.*, **70**: 159–173.
- RÜHLE E. (1967) — Podłoże czwartorzędu i jego wpływ na rozmieszczenie osadów czwartorzędowych w Polsce. In: *Czwartorzęd Polski* (eds. R. Galon, J. Dylík): 9–17. *PWN. Warszawa*.

- RÜHLE E. (1970) — Les nouvelles unités stratigraphiques de la glaciation de la Pologne Centrale (Riss) sur le territoire entre la moyenne Vistule et le bas Bug (in Polish with French summary). *Acta Geogr. Lodz.*, **24**: 389–412.
- RÜHLE E. ed. (1973) — *Metodyka badań osadów czwartorzędowych*. Wyd. Geol. Warszawa.
- ŚRODOŃ A., GOŁĄBOWA M. (1956) — Pleistocene flora at Bedno, Central Poland (in Polish with English summary). *Biul. Inst. Geol.*, **100**: 7–44.
- URBAN B. (1995) — Palynological evidence of younger Middle Pleistocene Interglacials (Holsteinian, Reinsdorf, and Schöningen) in the Schöningen open cast lignite mine (eastern Lower Saxony, Germany). *Meded. Rijks Geol. Dienst.*, **52**: 175–186.
- URBAN B. (1997) — Grundzüge der eiszeitlichen Klima- und Vegetationsgeschichte in Mitteleuropa. In: *Homo heidelbergensis* von Mauer. Das Auftreten des Menschen in Europa (eds. G. A. Wagner, K. W. Beinhauer): 241–263. Heidelberg.
- WINNICKI J. (1997) — Geological structure of the Trzebnica Hills in the light of new investigation. *Geol. Quart.*, **41** (3): 365–380.
- WINTER H. (1998) — Nowe spojrzenie na stratyografię plejstocenu środkowego na obszarze Polski północno-wschodniej jako wynik badań palinologicznych. *Mat. V Konf. "Stratygrafia plejstocenu Polski"* (eds. S. Lisicki *et al.*): 7–9. Iznota.
- WINTER H., LISICKI S. (1998) — New palyno- and lithostratigraphic interpretation of the Cenozoic lake sediments the section Goleń, Mazury Lakeland. *Geol. Quart.*, **42** (1): 87–98.
- WOŹNIAK P. (1989) — Zbójno Interglacial in the Suwałki Region (in Polish with English summary). *Kwart. Geol.*, **33** (3–4): 561–572.
- ŻARSKI M. (1994) — Extent of the Liwiec (Wieprz) Glaciation of the Middle Wisła River Valley (Central Poland) in light of age determination with the use of TL method (in Polish). *Prz. Geol.*, **42** (4): 285–286.

NOWE SPOJRZENIE NA STRATYGRAFIĘ OSADÓW JEZIORNICH I LODOWCOWYCH MŁODSZEJ CZĘŚCI ŚRODKOWEGO PLEJSTOCENU W POLSCE

Streszczenie

Przedstawiono nowy pogląd na stratyografię młodszej części środkowego plejstocenu Polski, a zwłaszcza na liczbę i wiek zlodowaceń środkowopolskich. Charakterystyka tego okresu została oparta między innymi na analizie 10 profili osadów czwartorzędowych (Grabówka, Nowiny, Goleń, Raczki Wielkie, Losy, Nidzica, Krępiec, Ossówka, Zbójno i Bedno) ze środkowo-wschodniej i północno-wschodniej Polski (fig. 1 i 2). Podstawą rozważań było wyłączenie z interglacjalu eemskiego osadów tzw. morza sztumskiego, coraz liczniejsze doniesienia o bardziej południowym zasięgu gliny zwalowej o parametrach geochronologicznych i litologicznych właściwych dla zlodowacenia warty, odkrycie nowych stanowisk interglacjalu zbójna w Polsce oraz przeprowadzona rekonstrukcja sieci rzecznej interglacjalu mazowieckiego w Polsce środkowej (fig. 3). Nawiązano do wcześniejszych prób korelacji poszczególnych jednostek zlodowacenia solawy w środkowo-zachodniej Europie (L. Marks, 1991) oraz opinii o występowaniu 4 interglacjalów młodszych od zlodowacenia sanu 2 (elstery 2) na kontynencie europejskim (L. Lindner, B. Marciniak, 1998). Umożliwiło to nieco inne spojrzenie na zagadnienie zlodowaceń środkowopolskich (fig. 2 i 4) oraz na bardziej uzasadnioną korelację głównych jednostek podziału klimatostratigraficznego młodszej części środkowego plejstocenu Polski z analogicznymi jednostkami na obszarze Niemiec i Równiny Rosyjskiej (fig. 5).

Wyróżnione ostatnio na obszarze Niemiec w profilu Schöningen (por. B. Urban, 1995, 1997) dwa nowe interglacjalny: reinsdorf i schöningen (fig. 5), mimo że nie są oddzielone od siebie i starszego od nich interglacjalny holsztyńskiego pokładami glin zwalowych, to jednak należy uznać za odpowiedniki interglacjalów: zbójna i lubawy (por. L. Lindner, B. Marciniak, 1998). Osady jeziorne interglacjalny holsztyńskiego w tym profilu, wraz z poprzedzającymi i następującymi po nim stadiami i interstadiami, leżą na glinie zwalowej zlodowacenia elstery 2, a osady najmłodszego z wymienionych interglacjalów (schöningen) przykryte są gliną zwalową stadium drenthe i wyżej leżącymi osadami ekstraglacialnymi(?) stadium warthe oraz osadami cemskimi (B. Urban, 1997).

Na obszarze Równiny Rosyjskiej w profilu Czekalin (por. N. S. Bolikhovskaya, N. G. Sudakova, 1996) wyróżniono ostatnio dwa nowe interglacjalny: czekaliński i czerepiecki, jako młodsze od interglacjalny lichwińskiego

i również nie rozdzielone osadami lodowcowymi. Leżą one na glinie zwalowej zlodowacenia oki i na niżej położonych osadach interglacjalny białowieckiego. Są one w tym profilu rozdzielone osadami odpowiadającymi ochłodzeniom klimatycznym: kaluskiemu i żidzińskiemu (fig. 5), a osady najmłodszego z tych interglacjalów (czerepieckiego) przykryte są dwiema glinami zwalowymi zlodowacenia dniewrowskiego.

Stwierdzono daleko idące podobieństwo w liczbie i randze klimatostratigraficznej jednostek młodszej części środkowego plejstocenu na obszarze Niemiec, Polski i Równiny Rosyjskiej. Zarówno w Niemczech, jak i na Równinie Rosyjskiej, wykształcenie tych jednostek w dużym stopniu przypomina rytm ociepleń i ochłodzeń klimatycznych identyfikowanych na obszarze Polski w interglacjalny wielkim (w rozumieniu S. Z. Różyckiego, 1964, 1972) czy też interglacjalny mazowieckiego *s.l.* (w rozumieniu E. Rühlego, 1973). Nowsze dane potwierdzają istnienie tego rytmu w owym czasie na obszarze Polski, ale tylko w obrębie kopalnych systemów obniżen, których rozwój był uwarunkowany procesami neotektonicznymi i działalnością rzeczną (por. fig. 3). W strefie kopalnych wysoczyzn polodowcowych przetrwały na obszarze Polski pokłady glin zwalowych oddzielające interglacjalny mazowiecki, zbójna i lubawy, wskazujące na co najmniej dwukrotną transgresję lądolodu skandynawskiego.

Z dotychczas zebranych materiałów dotyczących plejstocenu Polski oraz przykrycia osadów interglacjalny schöningen przez glinę zwalową stadium drenthe (B. Urban, 1995), a osadów interglacjalny czerepieckiego przez dwie gliny zwalowe zlodowacenia dniewrowskiego (N. S. Bolikhovskaya, N. G. Sudakova, 1996), mogą wynikać ponadto dwa kolejne wnioski. Pierwszym z nich jest to, że wyróżniany na obszarze Polski środkowo-wschodniej epizod glacialny w dotychczasowej pozycji zlodowacenia odry, oddzielającego interglacjalny zbójna i lubawy, jest prawdopodobnie starszy od stadium drenthe w Niemczech i zlodowacenia dniewrowskiego na Równinie Rosyjskiej. W niniejszym artykule proponuje się dla niego nazwę zlodowacenia krzny. Drugim wnioskiem jest propozycja uznania dotychczasowego rozprzestrzenienia lądolodu skandynawskiego o zasięgu zlodowacenia odry za stadią maksymalny zlodowacenia warty (fig. 4).