

Impact of the projected climate change on soybean water needs in the Kuyavia region in Poland

Wiesława Kasperska-Wołowicz¹ ✉ , Stanisław Rolbiecki² , Hicran A. Sadan² ,
Roman Rolbiecki² , Barbara Jagosz³ , Piotr Stachowski⁴ , Daniel Liberacki⁴ ,
Tymoteusz Bolewski¹ , Piotr Prus⁵ , Ferenc Pal-Fam⁶ 

¹) Institute of Technology and Life Sciences – National Research Institute, Hrabska Av. 3, Falenty, 05-090 Raszyn, Poland

²) Bydgoszcz University of Science and Technology, Faculty of Agriculture and Biotechnology,
Department of Agrometeorology, Plant Irrigation and Horticulture, Bydgoszcz, Poland

³) University of Agriculture in Krakow, Faculty of Biotechnology and Horticulture,
Department of Plant Biology and Biotechnology, Krakow, Poland

⁴) Poznan University of Life Sciences, Faculty of Environmental Engineering and Mechanical Engineering,
Department of Land Improvement, Environmental Development and Spatial Management, Poznań, Poland

⁵) Bydgoszcz University of Science and Technology, Faculty of Agriculture and Biotechnology,
Laboratory of Economics and Agribusiness Advisory, Bydgoszcz, Poland

⁶) Hungarian University of Agriculture and Life Sciences (MATE), Kaposvár, Hungary

RECEIVED 19.10.2021

REVIEWED 10.11.2021

ACCEPTED 15.12.2021

Abstract: According to the SRES A1B climate change scenario, by the end of the 21st century temperature in Poland will increase by 2–4°C, no increase in precipitation totals is predicted. This will rise crop irrigation needs and necessity to develop irrigation systems. Due to increase in temperature and needs of sustainable agriculture development some changes in crop growing structure will occur. An increase interest in high protein crops cultivation has been noted last years and further extension of these acreage is foreseen. Identifying the future water needs of these plants is crucial for planning and implementing sustainable agricultural production. In the study, the impact of projected air temperature changes on soybean water needs, one of the most valuable high-protein crops, in 2021–2050 in the Kuyavia region in Poland was analysed. The calculations based on meteorological data collected in 1981–2010 were considered as the reference period. Potential evapotranspiration was adopted as a measure of crop water requirements. The potential evapotranspiration was estimated using the Penman–Monteith method and crop coefficient. Based on these estimations, it was found that in the forecast years the soybean water needs will increase by 5% in the growing period (from 21 April to 10 September), and by 8% in June–August. The highest monthly soybean water needs increase (by 15%) may occur in August. The predicted climate changes and the increase in the arable crops water requirements, may contribute to an increase in the irrigated area in the Kuyavia region and necessity of rational management of water resources.

Keywords: climate change scenario, crop water needs, evapotranspiration, *Glycine max* (L.) Merr., irrigation, precipitation, protein crops

INTRODUCTION

In the light of the climate change scenario considered to be the most probable (A2), it is predicted that at the end of the 21st century the increase in global temperature will amount to about 4°C [ALCAMO *et al.* 2007; IPCC 2007; ŁABĘDZKI *et al.* 2013; RANDALL *et al.* 2007]. In Poland, an increase in temperature can be expected in the range of 2–4°C [ŁABĘDZKI *et al.* 2013]. On the basis of over 80 years of observations carried out in Bydgoszcz, situated in Kuyavian-Pomeranian Province, it has been noticed a significant increasing trend in the mean annual air temperature in this region, raising by 0.19°C per 10 years and consequently by about 2.0°C per 100 years [KASPERSKA-WOŁOWICZ, BOLEWSKI 2015]. In this region also unfavorable distribution of precipitation, with frequent and long-lasting dry periods is recorded. Such periods, lasting even longer than 21 days, are also recorded in wet years [KASPERSKA-WOŁOWICZ *et al.* 2003]. It is important that, most of the scenarios for Poland do not predict an increase in total annual precipitation. However, one can expect an increase in winter precipitation and a decrease in summer precipitation [ALCAMO *et al.* 2007; EEA 2008; IPCC 2007; KUNDZEWICZ 2003; 2007; PARRY 2000; RANDALL *et al.* 2007]. Similar climate change scenarios also apply to the Bydgoszcz region, both in terms of temperature [BAK, ŁABĘDZKI 2014b] and precipitation [BAK, ŁABĘDZKI 2014a].

Kuyavian-Pomeranian Province is an important area of agricultural crop production in Poland. Especially, its central and southern part, called Kuyavia (*pl Kujawy*), faces a serious challenge of increasing risk of production caused by aforementioned climatic changes, manifested among others by changes in temperature and precipitation distribution [ŁABĘDZKI 2009a; 2009b]. In Kuyavian-Pomeranian Province, similarly to Greater Poland and Mazovian provinces, there is the greatest natural need for the development of irrigation in Poland. This region, especially Kuyavia, is also characterized by very high irrigation water needs due to climate conditions (large rainfall deficits resulting from negative values of the climatic water balance, that is the difference between precipitation and reference evapotranspiration) [KACA 2017; KACA, REK-KACA 2019; ŁABĘDZKI 2017].

The global warming being observed for years may, due to the high climatic requirements of soybeans [ŚLIWA *et al.* 2015], contribute to increasing both the area and efficiency of soybean production in Poland. Soybean (*Glycine max* (L.) Merr.) is one of the most valuable species among legumes. Soybean, which contains 40–45% protein, 18–24% fat and 26% carbohydrate, is important for both animal and human nutrition [KARACA *et al.* 2018]. Due to the fact that it comes from the continental climate, it is genetically adapted to periodical water deficits. Soybean tolerates drought quite well, what is favored on the one hand by strongly developed pile root system, and on the other hand by hair on the stems, leaves and pods reducing transpiration. However, in the conditions of prolonged drought, soybean reacts very strongly to the lack of water: it grows poorly, has fewer pods and seeds in the pods, has small seeds and matures unevenly. Prematurely dried seeds from the upper parts of the stem do not develop viable embryos. Under such conditions, the germination capacity of soybean can drop even to 40% [BASF 2017; JANECZKO *et al.* 2011]. In the research by ECK *et al.* [1987], water stress throughout the last three weeks of the soybean growing period reduced yields in the two consecutive years of the study by 21% and 65%, respectively. Water stress imposed at the beginning seed

formation and extending to the end of the growing period (5 weeks) reduced yields by 45% and 88%, respectively, in two consecutive years of the study.

The area of soybean cultivation in Poland is not large, in the years 2014–2019 ranged between 12000 and 24000 ha and has been increasing [Polska... 2021]. The low cultivation area of soybean is mainly due to the high climatic requirements of this species, i.e. sowing in soil at temperature at least of 12°C and the average air temperature during crop growth not lower than 15°C [FLOREK *et al.* 2012; JERZAK *et al.* 2012]. According to the most recent source, the optimal date for sowing soybean in Poland is from the last decade of April till the first decade of May, when the top layer of the soil reaches temperature of 10°C [COBORU 2021; DORSZEWSKI *et al.* 2019; Polska... 2021]. The reason for the increase in the acreage of soybean cultivation in Poland is primarily the growing number of cultivars adapted to cultivation in the climatic conditions of Poland, as well as compared to other high-protein crops, higher content of nutrients and lower soil requirements [GAWĘDA *et al.* 2014].

Nowadays, the cultivation of soybean in Poland is subsidized due to the fact that it is a high-protein plant. As a result, the profitability of soybean cultivation is higher than that of cereals (including maize) and similar to the profitability of rapeseed. In addition, this plant cultivation is beneficial for soil biodiversity [DORSZEWSKI *et al.* 2019; Polska... 2021].

The costs of establishing a soybean plantation can be balanced by crop yield amount of about 2 Mg·ha⁻¹. In Poland the yield of soybean varies from about 1.5 to 2.5 Mg·ha⁻¹ but even more than 3 Mg·ha⁻¹ have been already achieved. In addition to the profit resulting directly from grain production, the profitability of soybean cultivation is also influenced by additional economic benefits, as nitrogen bound by nodular bacteria, improved soil structure and an increase in yield of succeeding crops. The value of these additional benefits can be estimated at approximately PLN800·ha⁻¹ [Polska... 2021].

Summing up, among the economic and environmental/agronomy opportunities of growing soybean in climate conditions of Poland are: growing demand for soybean in food industry, fodder and biofuel; increasing number of soybean purchase points in the country; currently subsidies for high-protein crops; non-genetically modified varieties; crop rotation enriched with soybean makes production more sustainable and diversified; due to symbiosis with nodule bacteria less doses of nitrogen fertilizers are required for soybeans and crops grown after; increasing tendency in air temperature. Among the risks are: lack of agricultural subsidies; reduction in crop yield and production due to unfavorable weather and not maintained irrigation and drainage systems.

In this context, the question arises as to how the expected climate change will affect the water needs in crop production in Poland. This study is an attempt to analyze this impact in relation to soybean cultivation in 2021–2050 based on the climate change scenario for Poland SRES: A1B. The A1 scenarios presents three alternative directions in terms of technological changes in the energy system, which differ in technological emphasis: A1FI – fossil intensive, A1T – non-fossil energy sources, and A1B – a balance between all sources, where “balanced” is not relying too heavily on one particular energy source [IPCC 2007].

The aim of our study was to estimate the water needs of soybeans during the growing season in the period 2021–2050 in

the Kuyavia region. The calculations based on the current climate conditions and expected changes in air temperatures. The Kuyavia region is located in central Poland, the zone with the greatest needs for supplementary irrigation during the growing season [JAGOSZ *et al.* 2021; KACA, REK-KACA 2019; ŁABĘDZKI 2017; MARKIEWICZ 2011; STACHOWSKI, ROLBIECKI *et al.* 2021].

STUDY MATERIALS AND METHODS

The study is based on the projected in the years 2021–2050 values of average monthly air temperatures for the agricultural area, called Kuyavia, situated in the southern part of Kuyavian-Pomeranian Province, in central Poland (Fig. 1). The predicted values were used according to the climate change scenario for Poland SRES (Special Report on Emissions Scenarios): A1B [BĄK, ŁABĘDZKI 2014b; IPCC 2007]. As the reference, thirty years period of 1981–2010 was adopted, using the average monthly air temperatures at Bydgoszcz measured at the agro-meteorological station located in the experimental field of the Institute of Technology and Life Sciences. The station, due to the long history of measurements, can be considered as representative for the Kuyavia agro-climatic conditions. The calculations were made for the growing season of soybeans, that is from 21 April to 10 September.

The water needs of soybeans (*Glycine max* (L.) Merr.) were calculated using the crop coefficient method based on the reference evapotranspiration according to Penman–Monteith formula [ŁABĘDZKI *et al.* 1996]. In this method, water needs are identified with crop evapotranspiration of a given plant species. The crop evapotranspiration of soybeans was calculated from the Equation (1):

$$ET_p = ET_o \cdot kc \quad (1)$$

where: ET_p = crop potential evapotranspiration (mm); ET_o = reference evapotranspiration (mm); kc = crop coefficient calculated as the relation of evapotranspiration measured in conditions of sufficient soil moisture to the reference evapotranspiration [ŁABĘDZKI 2006].

The values of the crop coefficient for the individual months of the soybean growing season are presented in Table 1.

In the reference period (1981–2010) the reference evapotranspiration was determined according to the Penman–Monteith Equation (2) [ALLEN *et al.* 1998; DOORENBOS, KASSAM 1979; DOORENBOS, PRUITT 1977; ŁABĘDZKI *et al.* 2011]:

$$ET_o = n \frac{0.408 \Delta R_n + \gamma \frac{900}{T+273} u (e_s - e_a)}{\Delta + \gamma(1 + 0.34u)} \quad (2)$$

where: ET_o = reference evapotranspiration (mm·d⁻¹); n = number of days; Δ = slope vapour pressure curve (kPa·°C⁻¹); R_n = net solar radiation at the crop surface (MJ·m⁻²·d⁻¹); γ = psychrometric constant (kPa·°C⁻¹); T = air temperature at 2 m height (°C); u = wind speed at 2 m height (m·s⁻¹); e_s = saturation vapour pressure (kPa); e_a = actual vapour pressure (kPa).

In the years 2021–2050 the reference evapotranspiration was calculated using simple linear regression between the Penman–Monteith reference evapotranspiration and air temperature. These equations were determined using the data of the reference

Table 1. Crop coefficient for the Penman–Monteith equation for soybean in the Kuyavia region

Month	Decade	Crop coefficient
April	3	0.13
May	1	0.67
	2	
	3	
June	1	1.00
	2	
	3	
July	1	1.10
	2	
	3	
August	1	0.90
	2	
	3	
September	1	0.17

Source: own elaboration on the basis of ALLEN *et al.* [1998], DOORENBOS and PRUITT [1977], DOORENBOS and KASSAM [1979].

period (2021–2050). The same methodological assumption was previously adopted by ŁABĘDZKI *et al.* [2013] estimating reference evapotranspiration according to the Penman–Monteith to determine water needs of late potato crop evapotranspiration in the long term period of 2021–2050 and 2071–2100.

The results were statistically processed by determining the following values: mean, normal (median), maximum and minimum, as well as standard deviation and coefficient of variation. An attempt was also made to specify the possible tendencies of changes in the analyzed indicator of soybean water needs in two compared periods with the use of linear regression analysis and identifying correlation and determination coefficients. The significance of the correlation coefficients, with the sample size $n = 30$, was determined for $p = 0.05$. The value of the correlation coefficient, according to the confidence interval, was significant for $r_p \geq 0.362$.

RESULTS

The highest differences in monthly sums of soybean water needs in the reference period (1981–2010), expressed as standard deviation, were calculated in the months from May to August and ranged from 8.0 to 18.7 mm (Tab. 2). The highest monthly standard deviation values, 13.4 and 18.7 mm, occurred in June and July, respectively. The strongest diversification of monthly sums of soybean water needs in the forecast years (2021–2050) also was noted in the period from May to August and varied from 9.9 to 16.8 mm. Also in June and July, the highest monthly standard deviation values, amounting to 16.8 and 14.9 mm, respectively, were predicted. Relative differentiation of soybean water needs during the period from April to September, expressed by the coefficient of variation, amounted to 8.3 and 9.3% for the reference and forecast period, respectively. The highest values of coefficient of variation were found in September (15.0%) in the reference period and May (17.5%) in the forecast one.




Fig. 1. Location of Kuyavian-Pomeranian Province in Poland (a) and Europe (b), location of the agro-climatic station in Bydgoszcz and the potential irrigation needs in Kuyavian-Pomeranian Province [ŁABĘDZKI 2017]: dark grey – high, grey – moderate, white – low (c). The highest needs concern south and south-western part of the province, in which a large part is the Kuyavia region; source: own elaboration and literature

Table 2. Statistical characteristics of soybean water needs during the growing season (from 21 April to 10 September) in the reference (1981–2010) and forecast (2021–2050) years

Characteristic	Months						
	21–30 Apr	1–31 May	1–30 Jun	1–31 Jul	1–31 Aug	1–10 Sept	Apr–Sep
1981–2010							
Mean (mm)	2.8	66.0	106.9	127.4	84.6	2.8	390.5
Minimum (mm)	2.3	46.3	80.4	93.9	66.5	2.1	311.8
Maximum (mm)	3.9	81.6	139.4	167.2	105.3	3.7	465.5
Median (mm)	2.7	67.5	106.0	120.1	83.9	2.7	384.2
Standard deviation (mm)	0.4	8.0	13.4	18.7	9.9	0.4	32.6
Coefficient of variation (%)	13.0	12.5	12.5	14.7	11.7	15.0	8.3
2021–2050							
Mean (mm)	2.6	61.3	109.2	137.9	97.9	3.5	412.4
Minimum (mm)	2.0	43.1	78.2	104.2	74.8	2.8	332.5
Maximum (mm)	3.5	85.6	140.9	164.6	118.3	4.0	483.3
Median (mm)	2.6	60.3	109.1	137.8	97.9	3.4	414.8
Standard deviation (mm)	0.3	10.7	16.8	14.9	9.9	0.3	38.4
Coefficient of variation (%)	12.1	17.5	15.4	10.8	10.1	9.5	9.3

Source: own study.

Table 3 presents the linear trend equations of the temporal variability of soybean water needs, which were determined for its growing periods (from 21 April to 10 September) in two periods 1981–2010 and 2021–2050. The results show that in the forecast period (2021–2050), with the exception of the first decade of September, the monthly water needs of soybean presented an upward tendency.

In each decade (ten years) of the forecast thirty years (2021–2050), the seasonal water needs of soybean (in the period from the 3rd decade of April to the 1st decade of September) increased by 11 mm. Among the analysed months, soybean water needs increased the most in June and August, by 4.0 mm and 4.4 mm, respectively, but the significant trend was found only in August (Fig. 2, Tab. 3).

Table 3. Trend equations of the sums of soybean water needs (mm) during the growing season (from 21 April to 10 September) in the reference (1981–2010) and forecast (2021–2050) years

Period	Trend equation	Coefficient of determination R^2	Tendency of water needs (mm-decade ⁻¹)
1981–2010			
21–30 April	$y = 0.0095x + 2.6231$	0.0535	0.1
1–31 May	$y = -0.1297x + 68.028$	0.0205	-1.3
1–30 June	$y = 0.2764x + 102.66$	0.0331	2.8
1–31 July	$y = -0.0704x + 128.5$	0.0011	-0.7
1–31 August	$y = -0.1297x + 86.58$	0.0133	-1.3
1–10 September	$y = 0.0007x + 2.8024$	0.0002	0.0
June–August	$y = 0.0762x + 317.74$	0.0005	0.8
April–September	$y = -0.0434x + 391.2$	0.0001	-0.4
2021–2050			
21–30 April	$y = 0.0081x + 2.4897$	0.0505	0.1
1–31 May	$y = 0.1836x + 58.459$	0.0228	1.8
1–30 June	$y = 0.395x + 103.05$	0.0429	4.0
1–31 July	$y = 0.0797x + 136.7$	0.0022	0.8
1–31 August	$y = 0.4393x + 91.115^*$	0.1540	4.4
1–10 September	$y = -0.0038x + 3.5238$	0.0101	0.0
June–August	$y = 0.914x + 330.86$	0.0678	9.1
April–September	$y = 1.1019x + 395.34$	0.0639	11.0

* Significant at $p = 0.05$.
 Source: own study.

The average daily water needs of soybean varied from month to month, the highest amount was found in July (4.4 mm in the forecast period and 4.1 mm in the reference period) (Fig. 3). In June, both in the forecast period and in the reference period, the average daily water needs of soybean was 3.6 mm. In May, daily water needs were estimated by 2.0 mm and 2.1 mm in the forecast and reference periods respectively, and in August by 3.2 mm and 2.7 mm in the forecast and reference periods respectively. Soybean needed the least amount of water (about




Fig. 2. Predicted amount of soybean water needs in August in the forecast years (2021–2050); source: own study




Fig. 3. Average daily water needs of soybean during the growing season in the reference (1981–2010) and forecast (2021–2050) years; source: own study

0.3 mm per day) in the beginning and the end of development stages, i.e. third decade of April and the first decade of September.

The calculations carried out for the thermal conditions forecast in the Kuyavia region in the thirty-year period 2021–2050 show that the water needs for soybean cultivation, expressed by the values of potential evapotranspiration, will increase (Fig. 4, Tab. 4). In the growing season of soybean (from the 3rd decade of April to the 1st decade of September) an increase in water needs by 5% (21 mm) can be expected. In the period June–August, the value of water needs will increase by 26 mm (8%) in view of the expected temperature change. The highest monthly water needs increase for soybean (by 13 mm, i.e. 15%) may occur in August.




Fig. 4. Cumulative sums of soybean water needs from the third decade of April till the first decade of September in the reference (1981–2010) and forecast (2021–2050) years; source: own study

Table 4. The soybean water needs (mm) during the growing season (from 21 April to 10 September) in the reference (1981–2010) and forecast (2021–2050) years

Years	Period							
	21–20 Apr	1–31 May	1–30Jun	1–31 Jul	1–31 Aug	1–10 Sep	Jun– Aug	Apr– Sep
1981–2010 (A)	3	66	107	127	85	3	319	391
2021–2050 (B)	3	61	109	138	98	3	345	412
Difference (mm) A–B (%)	0	–5	2	11	13	0	26	21
	0	8	2	9	15	0	8	5

Source: own study.

DISCUSSION

Taking into account the importance of soybean cultivation and problem of decreasing water resources due to climate change, the discussion is focused on main topic of the study the soybean water needs.

When deciding to grow a given plant species, it is important to estimate the size of the yield that allows achieving a level of direct surplus competitive with other crops. The profitability of crop cultivation can be measured by the value of the gross margin per 1 ha, that is (in absolute term) calculated as the difference between the production value (revenue) and direct costs. Currently there is large differentiation of the soybean production value in Poland, influenced mainly by agronomy practices/cultivation technology and weather. The variation in direct costs is small, the dominant share in these there is that of the seed material. The analysis of growing non-genetically modified soybean in Kuyavian-Pomeranian Province in Poland show, to be relatively profitable, the soybean yield should be at least 2.50 Mg·ha⁻¹. Currently at this yield the revenues are well above expenses (acc. to authors of current study, the gross margin rate is above 50%). Achieving at least this yield ensures a competitive gross margin with other crops [DORSZEWSKI *et al.* 2019].

The water needs of soybean in the Kuyavia region (central Poland), calculated using the crop coefficient method and expressed as potential evapotranspiration, amounted to 391 mm in the reference period (1981–2010) and 412 mm in the forecast period (2021–2050). For comparison, under the climate conditions of eastern Nebraska (quite similar to these in central Poland but more humid climate), the water needs of soybean (expressed as evapotranspiration) on average in the first five years of cultivation were 452 mm under irrigation and 431 mm without irrigation (rainfed soybean) [SUYKER, VERMA 2009]. In the study by SPECHT *et al.* [2001] high temperatures and no rainfall in 1995 led to such severe drought that the 100% of soybean evapotranspiration (equalled 410 mm) had to be supplemented by irrigation treatments. In turn, in the field experiment carried out in the climatic conditions of Vojvodina region (Novi Sad, Serbia) in 1993–2004 evapotranspiration rate in irrigation conditions ranged from 432 to 501 mm, and in rainfed conditions in the interval from 170 to 450 mm [PEJIĆ *et al.* 2011].

This study is a pioneering one dealing with the impact of the projected climate change on the water needs of soybean in the years 2021–2050 in the Kuyavia region. The results obtained in this study, indicating a systematic increase in the water needs of plants, are consistent with many previous reports on the water needs of other plant species [COSTA *et al.* 2007; FRAGA *et al.* 2018;

KENNY, HARRISON 1992; KUCHAR *et al.* 2015; 2017; KUCHAR, IWANICKI 2011; 2013; ŁABĘDZKI *et al.* 2013; SERRA *et al.* 2014]. The reports from such Institutions as the European Environment Agency [EFA 2008] or the Intergovernmental Panel on Climate Change [IPCC 2007] also predict an increase in water needs of arable crops in central Europe in the 21st century.

However, it should be clearly emphasized that in the present study, the increase in water needs estimated for soybean grown in the Kuyavia region is the effect only of an increase in air temperature, which results mainly from the adopted methodological assumptions [BAK, ŁABĘDZKI 2014b; ŁABĘDZKI *et al.* 2013]. In the current research, the forecast changes in other meteorological elements (solar radiation, air humidity, wind speed), which have undoubtedly big impact on evapotranspiration, were not taken into account. It was also assumed that the nature of the relationship between air temperature and reference evapotranspiration will not change. The relationships derived for individual months of the growing season in the 1981–2010 period were extrapolated and used for calculations of the reference evapotranspiration in the forecast thirty-year period 2021–2050. Similar methodological assumptions were apply in the study by ŁABĘDZKI *et al.* [2013], who extrapolated the relationships derived for the 1970–2004 period and then used them to calculate the reference evapotranspiration in two predicted long term periods (2021–2050 and 2071–2100). In the present study, it was also assumed that the crop coefficient would not change. However, it should be clearly emphasized that in the light of the current knowledge, there are no credible premises and no research indicating that these relationships and parameters, under the conditions of future climate change, will be the same [ŁABĘDZKI 2006; ŁABĘDZKI *et al.* 1996; 2013].

It seems very likely that the predicted climate change and the associated increase in plants water needs, as well as decrease in available water resources in soils, should result in an increase in irrigated area and an rise in irrigation water requirements. Such scenarios are considered in the other European countries, as well in other parts of the World [ABEDINPOUR *et al.* 2014; DEWEDAR 2021]. For example, net irrigation water needs is expected to increase by 40% in northern Germany, and by as much as 70% in south-eastern England [DÖLL 2002]. A similar increase can also be expected in Poland [ŁABĘDZKI 2009a; 2009b; ŁABĘDZKI *et al.* 2013]. Climate change also forces the need to modify the technology of water use and management of water resources in the field (improving the irrigation systems used) towards saving water and increasing the efficiency of water use [ALTOBELLI *et al.* 2021; BARKER *et al.* 2019; BELLALLOUI, MENGISTU 2008; FALLOON, BETTS 2010; ŁABĘDZKI 2009a; 2009b; OLESEN, BINDI 2002; SCHLEGEL *et al.* 2016].

Due to limited water resources, one of the strategies for optimum utilization of water resources in agriculture is deficit irrigation. This irrigation strategy has been investigated as a valuable and sustainable approach to production in dry areas [ABDELRAOUF *et al.* 2021] and it is appreciated in temperate climate as well. Based on the study of the effects of different irrigation regimes on soybean yield, water use efficiency, harvest index, oil and protein, it was found that any increase in the irrigation water amount resulted in the rise of the dry matter and seed yield of different soybean cultivars. Increasing the amount of irrigation water also had a significant influence on the percentage of protein and oil in seed [SHAMS *et al.* 2014]. In the study carried out by CRABTREE *et al.* [1985] soybean were irrigated using furrow systems. It was found that although the yield of alternate-furrow-irrigated soybean was significantly lower, an acceptable yield was still achieved with the use of 40–50% less irrigation water; i.e. the water savings in the alternate-furrow irrigation and the resulting lower yields are an acceptable compromise. To demonstrate the usefulness of supplemental irrigation as an alternative to full-season and non-irrigation to achieve high yield and seed quality of soybean, in the mid-south USA studies were conducted, which concluded that supplemental irrigation at the reproductive stage may be a possible alternative for full season irrigation of this species [BELLALOU, MENGISTU 2008]. Also, ECK *et al.* [1987] found that in soybean cultivation in the stressful climate of the Southern High Plains of the USA, limited irrigation is appropriate, however, the susceptibility of plants to drought stress during seed development may cause some limitations in the use of this irrigation method. In this study, soybean has been found to be a plant for which limited irrigation is more suitable than maize, but less than cotton, grain sorghum, or wheat. Also in the study of MARKOVIĆ *et al.* [2016] deficit irrigation of soybean was applied. The experiment was carried out in three different years in terms of meteorological conditions in sub-humid area. Averaged across the years at full irrigated plots, grain yield of soybean was reduced in comparison to deficit irrigated plots. These studies confirm that the management of deficit irrigation is an effective means of protecting water resources and increase soybean yields.

CONCLUSIONS

Based on calculations estimating the impact of climate change on soybean water needs, it was found that in the projected thirty-year season of 2021–2050, the water needs of soybean during its growing period (from the 3rd decade of April to the 1st decade of September) will increase and the expected growth will be 5% (21 mm). In the period from 1 June to 31 August, the water needs of soybean, in view of the expected temperature increase, will rise by 26 mm (8%). The highest monthly increase in soybean water needs, by 13 mm (i.e. 15%), may occur in August.

Growing soybean as a valuable nutrient and rotation crop can be a part of sustainable and profitable crop production in the near future. It will be possible in fields with an efficiently operating and maintained irrigation and drainage system.

The expected climate changes, and consequently the forecast increase in the arable crops water demands, may contribute to an increase in the irrigated area in the Kuyavia region and necessity of rational management of water resource.

ACKNOWLEDGMENTS

Andrzej Klimek, our colleague, created the Figure 1. Andrzej died while this publication was being prepared. Scientific cooperation with Andrzej was an honour for us. Andrzej will always remain in our memory.

REFERENCES

- ABEDINPOUR M., SARANGI A., RAJPUT T.B.S., SINGH M. 2014. Prediction of maize yield under future water availability scenarios using the AquaCrop model. *The Journal of Agricultural Science*. Vol. 152 (4) p. 558–574. DOI 10.1017/S0021859614000094.
- ABDELRAOUF R.E., EL-SHAWADFY M.A., DEWEDAR O.M., HOZAYN M. 2021. Field and modelling study for deficit irrigation strategy on roots volume and water productivity of wheat. *Journal of Water and Land Development*. No. 49 p. 129–138. DOI 10.24425/jwld.2021.137105.
- ALCAMO J., MORENO J.M., NOVÁKY B., HINDI M., COROBOV R., DEVOY R.J. N., GIANNAKOPOULOS C., MARTIN E., OLESN J.E., SHVIDENKO A. 2007. Europe. Climate Change 2007. Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Eds. M.L. Parry, O.F. Canziani, J.P. Palutikof, P.J. van der Linden, C.E. Hanson. Cambridge, UK. Cambridge University Press p. 541–580.
- ALLEN R.G., PEREIRA L.S., RAES D., SMITH M. 1998. Crop evapotranspiration. Guidelines for computing crop water requirements. FAO Irrigation and Drainage Paper. No. 56. Rome, Italy. FAO. ISSN 0254-5284 pp. 300.
- ALTOBELLI F., MARTA A.D., HEINEN M., JACOBS C., GIAMPIETRI E., MANCINI M., ..., DEL GIUDICE T. 2021. Irrigation Advisory Services: Farmers preferences and willingness to pay for innovation. *Outlook on Agriculture*. Vol. 48(2) p. 136–142. DOI 10.1177/00307270211002848.
- BARKER J.B., BHATTI S., HEEREN D.M., NEALE C.M., RUDNICK D.R. 2019. Variable rate irrigation of maize and soybean in West-Central Nebraska under full and deficit irrigation. *Frontiers in Big Data*. Vol. 2, 34. DOI 10.3389/fdata.2019.00034.
- BASF 2017. Soja. Poradnik agrotechniczny [Soybean. Agrotechnical guideline] 2017. [Access 23.06.2021]. Available at: <https://www.agro.basf.pl/Documents/broszury/poradnik-agrotechniczny-soja-basf-2017.pdf?1570453654459>
- BĄK B., ŁABĘDZKI L. 2014a. Thermal conditions in Bydgoszcz region in growing seasons 2011–2050 in view of expected climate change. *Journal of Water and Land Development*. No. 23 p. 21–29. DOI 10.1515/jwld-2014-0026.
- BĄK B., ŁABĘDZKI L. 2014b. Prediction of precipitation deficit and excess in Bydgoszcz region in view of predicted climate change. *Journal of Water and Land Development*. No. 23 p. 11–19. DOI 10.1515/jwld-2014-0025.
- BELLALOU N., MENGISTU A. 2008. Seed composition is influenced by irrigation regimes and cultivar differences in soybean. *Irrigation Science*. Vol. 26 p. 261–268. DOI 10.1007/s00271-007-0091-y.
- COBORU 2021. Wyniki plonowania odmian roślin rolniczych w doświadczeniach porejestrowych w województwie łódzkim. Soja 2020 [Yield results of agricultural plant varieties in post-registration experiments in the Łódź Province. Soybean 2020]. Sulejów. Centralny Ośrodek Badania Odmian Roślin Uprawnych SDOO pp. 14.

- COSTA J.M., ORTUÑO M.F., CHAVES M.M. 2007. Deficit irrigation as a strategy to save water: Physiology and potential application to horticulture. *Journal of Integrative Plant Biology*. Vol. 49(10) p. 1421–1434. DOI 10.1111/J.1672-9072.2007.00556.X.
- CRABTREE R.J., YASSIN A.A., KARGOUGOU I., McNEW R.W. 1985. Effects of alternate-furrow irrigation: Water conservation on the yields of two soybean cultivars. *Agricultural Water Management*. Vol. 10 (3) p. 253–264. DOI 10.1016/0378-3774(85)90015-0.
- DEWEDAR O., PLAUBORG F., EL-SHAFFIE A., MARWA A. 2021. Response of potato biomass and tuber yield under future climate change scenarios in Egypt. *Journal of Water and Land Development*. No. 49 p. 139–150. DOI 10.24425/jwld.2021.137106.
- DOORENBOS J., PRUITT W.O. 1977. Guidelines for predicting crop water requirements. *FAO Irrigation and Drainage Paper*. No. 24. Rome. FAO pp. 192.
- DOORENBOS J., KASSAM A. 1979. Yield response to water. *FAO Irrigation and Drainage Paper*. No. 33. Rome. FAO. ISBN 9251007446 pp. 192.
- DORSZEWSKI P., SOBCZYŃSKI T., WENDA-PIESIK A. 2019. Soja w województwach kujawsko-pomorskim i wielkopolskim – innowacyjne rozwiązania w uprawie i skarmianiu dla gospodarstw rolnych. Raport z badań dla Konsorcjum „Moja Soja” [Soybeans in Kuyavian-Pomeranian and Greater Poland provinces – innovative solutions in cultivation and feeding for farms. Research report for the “Moja Soja” Consortium] [online]. Minikowo. Kujawsko-Pomorski Ośrodek Doradztwa Rolniczego. [Access 20.08.2021]. Available at: <http://www.mojasoja.eu/>
- DÖLL P. 2002. Impact of climate change and variability on irrigation requirements: A global perspective. *Climatic Change*. Vol. p. 269–293. DOI 10.1023/A:1016124032231.
- ECK H.V., MATHERS A.C., MUSICK J.T. 1987. Plant water stress at various growth stages and growth and yield of soybeans. *Field Crops Research*. Vol. 17(1) p. 1–16. DOI 10.1016/0378-4290(87)90077-3.
- EEA 2008. Impacts of Europe’s changing climate – 2008 indicator-based assessment. Report No 4/2008. European Environment Agency. ISBN 978-92-9167-372-8 pp. 246.
- FALLOON P., BETTS R. 2010. Climate impacts on European agriculture and water management in the context of adaptation and mitigation – the importance of an integrated approach. *Science of The Total Environment*. Vol. 408(23) p. 5667–5687. DOI 10.1016/j.scitotenv.2009.05.002.
- FLOREK J., CZERWIŃSKA-KAYZER D., JERZAK M.A. 2012. Aktualny stan i wykorzystanie produkcji upraw roślin strączkowych [Current status and use of legume crops production]. *Fragmenta Agronomica*. Vol. 29(4) p. 45–55.
- FRAGA H., GARCÍA DE CORTÁZAR ATAURI I., SANTOS J.A. 2018. Viticultural irrigation demands under climate change scenarios in Portugal. *Agricultural Water Management*. Vol. 196 p. 66–74. DOI 10.1016/j.agwat.2017.10.023.
- GAWĘDA D., CIERPIAŁA R., BUJAK K., WESOŁOWSKI K. 2014. Soybean yield under different tillage systems. *Acta Scientiarum Polonorum Hortorum Cultus*. Vol. 13(1) p. 43–54.
- IPCC 2007. AR4 Climate change 2007. Fourth Assessment Report [online]. Intergovernmental Panel on Climate Change. [Access 23.06.2021]. Available at: <https://www.ipcc.ch/assessment-report/ar4/>
- JAGOSZ B., ROLBIECKI S., ROLBIECKI R., ŁANGOWSKI A., SADAN H.A., PTACH W., STACHOWSKI P., KASPERSKA-WOŁOWICZ W., PAL-FAM F., LIBERACKI D. 2021. The water needs of grapevines in central Poland. *Agronomy*. Vol. 11(3), 416. DOI 10.3390/agronomy11030416
- JANECKO A., BIESAGA-KOŚCIELNIAK J., DZIURKA M., OKLEŠTKOVÁ J., KOCUREK M., SZAREK-ŁUKASZEWSKA G., JANECKO Z. 2011. Response of Polish cultivars of soybean (*Glycine max* (L.) Merr.) to brassinosteroid application. *Acta Scientiarum Polonorum. Agricultura*. Vol. 10(2) p. 33–50.
- JERZAK M.J., CZERWIŃSKA-KAYZER D., FLOREK J., ŚMIGLAK-KRAJEWSKA M. 2012. Determinanty produkcji roślin strączkowych jako alternatywnego źródła białka w ramach nowego obszaru polityki rolnej w Polsce [Determinants of legume production as an alternative source of protein in a new area of agricultural policy in Poland]. *Roczniki Nauk Rolniczych*. Nr 99(1) p. 113–120.
- KACA E. 2017. Methodology of assessing the relative environmental validity of developing drainage and irrigation on a regional scale. *Journal of Water and Land Development*. No. 35 p. 101–112. DOI 10.1515/jwld-2017-0073.
- KACA E., REK-KACA G. 2019. Względna przyrodnicza zasadność rozwoju nawodnień w skali województw w Polsce [Relative nature of the development of irrigation in the scale of provinces in Poland]. In: *Proceedings of the Symposium on Plant Irrigation “Irrigation of plants in the light of sustainable rural development – natural, production and technical and infrastructural aspects”*. [11–14 June 2019 Bydgoszcz-Fojutowo, Poland] p. 33–35.
- KARACA C., TEKELIOĞLU B., BUYUKTAS D., BASTUG R. 2018. Relations between crop water stress index and stomatal conductance of soybean depending on cultivars. *Fresenius Environmental Bulletin*. Vol. 27(6) p. 4212–4219.
- KASPERSKA-WOŁOWICZ W., BOLEWSKI T. 2015. Zmienność temperatury powietrza w Bydgoszczy w latach 1931–2013 [Variability of air temperature in Bydgoszcz in the years 1931–2013]. *Woda-Środowisko-Obszary Wiejskie*. T. 15. Z. 3(51) p. 25–43.
- KASPERSKA-WOŁOWICZ W., ŁABĘDZKI L., BĄK B. 2003. Okresy posuszne w rejonie Bydgoszczy [Dry periods in the Bydgoszcz region]. *Woda-Środowisko-Obszary Wiejskie*. T. 3. Z. 9 p. 39–56.
- KENNY G.J., HARRISON P.A. 1992. The effects of climate variability and change on grape suitability in Europe. *Journal of Wine Research*. Vol. 3(3) p. 163–183. DOI 10.1080/09571269208717931.
- KUCHAR L., IWAŃSKI S. 2011. Symulacja opadów atmosferycznych dla oceny potrzeb nawodnień roślin w perspektywie oczekiwanych zmian klimatycznych [Rainfall simulation for the prediction of crop irrigation in future climate]. *Infrastruktura i Ekologia Terenów Wiejskich*. Vol. 5 p. 7–18.
- KUCHAR L., IWAŃSKI S. 2013. Ocena opadów atmosferycznych dla potrzeb produkcji roślinnej w perspektywie lat 2050–2060 i wybranych scenariuszy zmian klimatu w północno-centralnej Polsce [Rainfall evaluation for crop production until 2050–2060 and selected climate change scenarios for North Central Poland]. *Infrastruktura i Ekologia Terenów Wiejskich*. Vol. 2 p. 187–200.
- KUCHAR L., IWAŃSKI S., DIAKOWSKA E., GAŚIOREK E. 2015. Symulacja warunków hydrotermicznych w północnej części centralnej Polski w perspektywie lat 2050–2060 dla potrzeb produkcji roślinnej i wybranych scenariuszy klimatycznych [Simulation of hydrothermal conditions for crop production purpose until 2050–2060 and selected climate change scenarios for North Central Poland]. *Infrastruktura i Ekologia Terenów Wiejskich*. Vol. 2 p. 319–334. DOI 10.14597/infraeco.2015.2.1.026.
- KUCHAR L., IWAŃSKI S., DIAKOWSKA E., GAŚIOREK E. 2017. Ocena suszy meteorologicznej w 2015 roku w północnej części centralnej Polski z wykorzystaniem wskaźnika hydrotermicznego (HTC) w kontekście zmian klimatycznych [Assessment of meteorological drought in 2015 for North Central part of Poland using hydrothermal coefficient (HTC) in the context of climate change]. *Infrastruktura i Ekologia Obszarów Wiejskich*. Vol. 1 p. 257–273. DOI 10.14597/infraeco.2017.1.2.019.
- KUNDZEWICZ Z. 2003. Scenariusze zmian klimatu. W: *Czy Polsce grożą katastrofy klimatyczne?* [Climate change scenarios. In: *Is Poland*

- at risk of climate disasters?]. Warszawa. Komitet Prognoz "Polska 2000 Plus". Polski Komitet Międzynarodowego Programu „Zmiany globalne geosfery i biosfery” przy Prezydium PAN p. 14–31.
- KUNDZEWICZ Z. 2007. Projekcje zmian klimatu – ekstrema hydro-meteorologiczne [Climate change projections – hydrometeorological extremes]. Proceedings of the I Polish Conference ADAGIO. Poznań, Poland. 24 April 2007.
- ŁABĘDZKI L. 2006. Susze rolnicze. Zarys problematyki oraz metody monitorowania i klasyfikacji [Agricultural droughts. Outline of the issues and methods of monitoring and classification]. Woda-Środowisko-Obszary Wiejskie. Rozprawy naukowe i monografie. Nr 17. ISSN 1644-1095 pp. 107.
- ŁABĘDZKI L. 2009a. Przewidywane zmiany klimatyczne a rozwój nawodnień w Polsce [Foreseen climate changes and irrigation development in Poland]. Infrastruktura i Ekologia Terenów Wiejskich. Vol. 3 p. 7–18.
- ŁABĘDZKI L. 2009b. Expected development of irrigation in Poland in the context of climate change. Journal of Water and Land Development. No. 13 p. 17–29.
- ŁABĘDZKI L. 2017. Potrzeby i stan nawodnień w województwie kujawsko-pomorskim [Irrigation needs and condition in Kuyavian-Pomeranian Province]. Warsztaty dla interesariuszy projektu OPERA [Workshops for the stakeholders of the OPERA project]. [27 November 2017 Minikowo, Poland].
- ŁABĘDZKI L., BĄK B., LISZEWSKA M. 2013. Wpływ przewidywanej zmiany klimatu na zapotrzebowanie ziemniaka późnego na wodę [Impact of climate change on water needs of late potato]. Infrastruktura i Ekologia Terenów Wiejskich. Vol. 2(1) p. 155–165.
- ŁABĘDZKI L., KANECKA-GESZKE E., BĄK B., SŁOWIŃSKA S. 2011. Estimation of reference evapotranspiration using the FAO Penman-Monteith method for climatic conditions of Poland. In: Evapotranspiration. Ed. L. Łabędzki. Rijeka, Croatia. InTech. ISBN 98-953-307-251-7 pp. 446. DOI 10.5772/14081.
- ŁABĘDZKI L., SZAJDA J., SZUNIEWICZ J. 1996. Ewapotranspiracja upraw rolniczych – terminologia, definicje, metody obliczania – Przegląd stanu wiedzy [Evapotranspiration of agricultural crops – terminology, definitions, calculation methods. Review]. Falenty. IMUZ. Materiały Informacyjne. Nr 33 pp. 15.
- MARKOVIĆ M., JOSIPOVIĆ M., RAVLIĆ M., JOSIPOVIĆ A., ZEBEC V. 2016. Deficit irrigation of soybean (*Glycine max* (L.) Merr.) based on monitoring of soil moisture, in sub-humid area of eastern Croatia [online]. Romanian Agricultural Research. Vol. 33 p. 1–8. [Access 23.06.2021]. Available at: https://www.researchgate.net/publication/305267560_Deficit_irrigation_of_soybean_-_Glycinemax_L_Merr_based_on_monitoring_of_soil_moisture_in_sub-humid_area_of_eastern_croatia
- OLESEN J.E., BINDI M. 2002. Consequences of climate change for European agricultural productivity, land use and policy. European Journal of Agronomy. Vol. 16(4) p. 239–262. DOI 10.1016/S1161-0301(02)00004-7.
- PARRY M.L. (ed.) 2000. Assessment of potential effects and adaptation for climate change in Europe: The Europe ACACIA Project. Jackson Environment Institute. Norwich. University of East Anglia pp. 324.
- PEJIĆ B., MAKSIMOVIĆ L., CIMPEANU S., BUCUR D., MILIĆ S., ČUPINA B. 2011. Response of soybean to water stress at specific growth stages. Journal of Food Agriculture and Environment. Vol. 9(1) p. 280–284.
- Polska Soja 2021. Soja w Polsce. Uprawa soi w Polsce [Soybeans in Poland. Growing soybean in Poland]. [Access 23.06.2021]. Available at: https://www.polskasoja.pl/asp/pl_start.asp?typ=14&menu=3&strona=1
- RANDALL D.A., WOOD R.A., BONY S., COLMAN R., FICHEFET T., FYFE J., ... TAYLOR K.E. 2007. Climate models and their evaluation. In: Climate Change 2007. The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change [Eds. S. Solomon, D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor, H.L. Miller]. Cambridge, UK, New York, USA. Cambridge University Press pp.74.
- ROLBIECKI R., ROLBIECKI S., FIGAS A., JAGOSZ B., STACHOWSKI P., SADAN H. A., PRUS P., PAL-FAM F. 2021. Requirements and effects of surface drip irrigation of mid-early potato cultivar Courage on a very light soil in Central Poland. Agronomy. Vol. 11(1), 33. DOI 10.3390/agronomy11010033.
- SHAMS B.M., BOROUMANDNASAB S., MALEKI A. 2014. Effect of different irrigation regimes on yield water use efficiency and harvest index of soybean cultivars in Khorramabad. Journal of Irrigation Sciences and Engineering. Vol. 37(3) p. 13–20. DOI 20.1001.1.25885952.1393.37.3.2.4.
- SCHLEGEL A.J., ASSEFA Y., O'BRIEN D., LAMM F.R., HAAG L.A., STONE L.R. 2016. Comparison of corn, grain sorghum, soybean, and sunflower under limited irrigation. Agronomy Journal. Vol. 108 (2) p. 670–679. DOI 10.2134/agronj2015.0332.
- SERRA I., STREVER A., MYBURGH P., DELOIRE A. 2014. The interaction between rootstocks and cultivars (*Vitis vinifera* L.) to enhance drought tolerance in grapevine. Australian Journal of Grape Wine Research. Vol. 20 (1) p. 1–14. DOI 10.1111/ajgw.12054.
- SPECHT J.E., CHASE K., MACRANDER M., GRAEF G.L., CHUNG J., MARKWELL J.P., GERMANN M., ORF J.H., LARK K.G. 2001. Soybean response to water: A QTL analysis of drought tolerance. Crop Science. Vol. 41(2) p. 493–509. DOI 10.2135/cropsci2001.412493x.
- STACHOWSKI P., MARKIEWICZ J. 2011. Potrzeba nawodnień w centralnej Polsce na przykładzie powiatu kutnowskiego [The need of irrigation in central Poland on the example of Kutno country]. Rocznik Ochrona Środowiska. Vol. 13 p. 1453–1472.
- SUYKER A.E., VERMA S.B. 2009. Evapotranspiration of irrigated and rainfed maize–soybean cropping systems. Agricultural and Forest Meteorology. Vol. 149(3–4) p. 443–452. DOI 10.1016/j.agrfor-met.2008.09.010.
- ŚLIWA J., ZAJĄC T., OLEKSY A., KLIMEK-KOPYRA A., LORENC-KOZIK A., KULIG B. 2015. Comparison of the development and productivity of soybean (*Glycine max* (L.) Merr.) cultivated in western Poland. Acta Scientiarum Polonorum. Agricultura. Vol. 14(4) p. 81–95.