





GARGI J. TRIVEDI  (Anand)
VISHANT SHAH  (Vadodara)
JAITA SHARMA  (Vadodara)
RAJESH SANGHVI  (Anand)

On solution of non-instantaneous impulsive Hilfer fractional integro-differential evolution system

Abstract In this article, we have considered the non-instantaneous fractional integro-differential evolution system with Hilfer fractional differential operator in the Banach space and discussed its existence results for the mild solution for the equation with local and non-local conditions. These results are obtained by applying the method of a C_0 operator generated by the linear part of the equation combined with the concept of nonlinear functional analysis and the fixed point theorems. We have discussed the examples to highlight the applicability of the results.

2010 Mathematics Subject Classification: [034A08, 47H10, 47H20].

Key words and phrases: Fractional Evolution Equation, Hilfer fractional differential operator, Operator Semi-group, Non-instantaneous Impulses, Fixed Point Theorem..

1. Introduction Fractional calculus and differential equations became an important branch of applied mathematics. This is because of many problems from the fields of physical sciences, chemical sciences, biological sciences, finance, and image processing which are modeled using fractional differential operators and give better approximations than those modeled using integer order differential operators [13, 12, 1]. Researchers generalized the fractional order differential operator in a way that coincides with the integer order differential operator and this leads to the existence of many differential operators like Riemann-Liouville, Caputo, Grownwell-Letnikov, and Conformable fractional differential operators. The qualitative properties and applications of fractional dynamical systems are found in [3, 16, 4]. Sometimes the system with non-local initial conditions gives better approximations than classical conditions. The qualitative properties and applications of the non-local systems are found in [5]. Hilfer came up with a new fractional differential operator which is a homotopy between Riemann-Liouville and Caputo fractional differential operators. The qualitative properties and applications of dynamical systems with Hilfer differential operator including classical and non-local conditions are found in [7] and monograph Hilfer (2000).

Systems having an abrupt change in the state at a fixed time moment or in a small time interval are modeled into instantaneous impulsive evolution or not-instantaneous impulsive evolution equation respectively. The qualitative properties and applications of the integer order evolution systems with instantaneous impulses are found in [14] and the same for the fractional systems are found in [9, 8]. In some of the evolutionary processes, not-instantaneous impulses give better approximations instead of instantaneous impulses. The qualitative properties and applications of systems with non-instantaneous impulsive systems are found in [15, 11, 10].

This work considered non-instantaneous impulsive integro-differential fractional order ($0 < \lambda \leq 1$ and $0 \leq \mu \leq 1$) evolution system of Hilfer type

$$D_{0+}^{\lambda, \mu} u(\zeta) = -Au(\zeta) + f\left(\zeta, u(\zeta), \int_0^\zeta a(\zeta, \tau, u(\tau))d\tau\right), \zeta \in \left[\cup [s_i, t_{i+1}] \cup [s_p, T_0]\right]$$

$$u(\zeta) = g_k(\zeta, u(\zeta)), \zeta \in [t_1, s_1) \cup [t_2, s_2) \cup \dots \cup [t_p, s_p),$$

and discussed the existence of solutions with local condition $\mathcal{I}_{0+}^{(1-\lambda)(1-\mu)}u(0) = u_0$ and non-local $\mathcal{I}_{0+}^{(1-\lambda)(1-\mu)}[u(0) - h(u)] = u_0$ initial conditions over the finite interval $[0, T_0]$ in a Banach space \mathbb{U} . $D^{\lambda, \mu}$ differential operators of Hilfer type, $A : \mathbb{U} \rightarrow \mathbb{U}$ is a linear part of the integrodifferential evolution equation, $Ku = \int_0^\zeta a(\zeta, \tau, u(\tau))d\tau$ is nonlinear Volterra integral operator on \mathbb{U} , $f : [0, T_0] \times \mathbb{U} \times \mathbb{U} \rightarrow \mathbb{U}$ is nonlinear function and $g_k : [0, T_0] \times \mathbb{U}$ are set of non-linear functions applied in the interval $[t_k, s_k)$ for all $i = 1, 2, \dots, p$.

The outline of the manuscript is as follows: Section-2 discusses the preliminaries to establish the results, section-3 established existence result for the non-instantaneous fractional order Hilfer integro-differential evolution system with classical conditions followed by a nonlocal condition in section-4. Finally, the conclusion is discussed in section-5.

2. Preliminaries This section is devoted to the definitions of integral $\mathcal{I}_{\zeta_0}^\lambda$ and Hilfer fractional differential operator $\mathcal{D}_{\zeta_0+}^{\lambda, \mu}$, Wright-type function M_λ and properties, concept operator semi-group $\mathcal{T}(\zeta)$, the operators like $\mathcal{K}_\lambda(\zeta)$, $\mathcal{S}_{\lambda, \mu}(\zeta)$.

DEFINITION 2.1 (Jaiswal and Bahuguna (2020)) For $\lambda > 0$, the fractional integral of order λ of a function $h(\zeta)$ is defined by

$$\mathcal{I}_{\zeta_0}^\lambda h(\zeta) = \frac{1}{\Gamma(\lambda)} \int_{\zeta_0}^\zeta (\zeta - \tau)^{\lambda-1} h(\tau) d\tau,$$

provided the integral on the right exists.

DEFINITION 2.2 (Jaiswal and Bahuguna (2020)) The Hilfer fractional derivative of order λ , $0 < \lambda < 1$ and type μ , $0 \leq \mu \leq 1$ is defined by

$$\mathcal{D}_{\zeta_0+}^{\lambda,\mu} h(\zeta) = \mathcal{I}_{\zeta_0+}^{\mu(1-\lambda)} \frac{d}{d\zeta} \mathcal{I}_{\zeta_0+}^{(1-\lambda)(1-\mu)} h(\zeta),$$

provided the right value exists.

DEFINITION 2.3 (Jaiswal and Bahuguna (2020)) For all $\theta \in \mathbb{C}$ and $\lambda > 0$, the Wright-type function M_λ is defined as:

$$M_\lambda(\theta) = \sum_{n \in \mathbb{N}} \frac{(-\theta)^{n-1}}{\Gamma(1-\lambda n)(n-1)!} \quad (1)$$

provided the sum on the right exists.

Wright-type function satisfies the following properties:

- (1) $M_\lambda(\theta) > 0$ for all $\lambda > 0$.
- (2) For $-1 < \eta < \infty$ the integral, $\int_0^\infty \theta^\eta M_\lambda(\theta) d\theta = \frac{\Gamma(1+\eta)}{\Gamma(1+\lambda\eta)}$.
- (3) For $r > 0$ the integral, $\int_0^\infty \frac{\lambda}{\theta^{\lambda+1}} e^{-r\theta} M_\lambda(\theta^{-\lambda}) d\theta = e^{-r^\lambda}$ for all $\lambda > 0$.

Let, $\mathcal{T}(\zeta)$ be the family of semi-group generated by the linear operator $-A$ and define two linear operators $\mathcal{S}_\lambda(\zeta)$ and $\mathcal{Q}_\lambda(\zeta)$ as:

$$\mathcal{S}_\lambda(\zeta) = \int_0^\infty M_\lambda(\theta) \mathcal{T}(\zeta^\lambda \theta) d\theta \quad (2)$$

$$\mathcal{Q}_\lambda(\zeta) = \int_0^\infty \lambda \theta M_\lambda(\theta) \mathcal{T}(\zeta^\lambda \theta) d\theta \quad (3)$$

Following properties are satisfied by $\mathcal{S}_\lambda(\zeta)$ and $\mathcal{Q}_\lambda(\zeta)$.

LEMMA 2.4 (Jaiswal and Bahuguna (2020)) If $\mathcal{T}(\zeta)$ be the family of C_0 -semigroup generated by the linear operator $-A$ for all $\zeta \in [0, T_0]$ then the families of operators $\mathcal{S}_\lambda(\zeta)$ and $\mathcal{Q}_\lambda(\zeta)$ defined by (2) and (3) are:

- (1) continuous and bounded for all $\zeta \in [0, T_0]$.
- (2) strongly continuous over the interval $\zeta \in (0, T_0]$.

The operators $\mathcal{S}_\lambda(\zeta)$ and $\mathcal{Q}_\lambda(\zeta)$ generate new linear operators $\mathcal{S}_{\lambda,\mu}(\zeta)$ and $\mathcal{K}_\lambda(\zeta)$.

$$\mathcal{S}_{\lambda,\mu}(\zeta) = \mathcal{I}_0^{\mu(1-\lambda)} \mathcal{K}_\lambda(\zeta) \quad (4)$$

$$\mathcal{K}_\lambda(\zeta) = \zeta^{\lambda-1} \mathcal{Q}_\lambda \zeta \quad (5)$$

These operators $\mathcal{S}_{\lambda,\mu}(\zeta)$ and $\mathcal{K}_\lambda(\zeta)$ satisfies following properties:

LEMMA 2.5 (*Jaiswal and Bahuguna (2020)*) If $\mathcal{T}(\zeta)$ be the family of C_0 -semigroup generated by the linear operator $-A$ for all $\zeta \in [0, T_0]$ then the families of operators $\mathcal{S}_{\lambda, \mu}(\zeta)$ and $\mathcal{K}_\lambda(\zeta)$ defined by (4) and (5) are:

- (1) continuous and bounded for all $\zeta \in [0, T_0]$.
- (2) strongly continuous over the interval $\zeta \in (0, T_0]$.
- (3) $\|\mathcal{K}_\lambda(\zeta)u\| \leq \frac{\zeta^{\lambda-1}M}{\Gamma(\lambda)}\|u\|$
- (4) $\|\mathcal{S}_{\lambda, \mu}(\zeta)u\| \leq \frac{M(\mu(1-\lambda))\zeta^{\lambda+\mu-\lambda\mu-1}}{\Gamma(\lambda+\mu-\lambda\mu-1)}\|u\|$.

Where, M is the bound of $\mathcal{T}\zeta$ over the interval $[0, T_0]$.

THEOREM 2.6 (*Banach Fixed Point Theorem v. Borah and Bora (2019)*) Let E be closed subset of a Banach Space $(\mathbb{U}, \|\cdot\|)$ and let $T : E \rightarrow E$ contraction then, T has unique fixed point in E .

THEOREM 2.7 (*Krasnoselskii's Fixed Point Theorem v. Borah and Bora (2019)*) Let E be closed convex nonempty subset of a Banach Space $(\mathbb{U}, \|\cdot\|)$ and P and Q are two operators on E satisfying:

- (1) $Pu + Qv \in E$, whenever $u, v \in E$,
- (2) P is contraction,
- (3) Q is completely continuous

then, the equation $Pu + Qu = u$ has a solution.

3. Equation with Classical Conditions This section establishes the existence results for fractional order ($0 < \lambda \leq 1$ and $0 \leq \mu \leq 1$) non-instantaneous impulsive Hilfer integro-differential evolution systems with classical conditions.

$$\begin{aligned}
 D_{0+}^{\lambda, \mu} u(\zeta) &= -Au(\zeta) + f\left(\zeta, u(\zeta), \int_0^\zeta a(\zeta, \tau, u(\tau))d\tau\right), \\
 &\zeta \in \left[\cup [s_i, t_{i+1}] \right] \cup [s_p, T_0] \\
 u(\zeta) &= g_k(\zeta, u(\zeta)), \quad \zeta \in [t_1, s_1] \cup [t_2, s_2] \cup \dots \cup [t_p, s_p] \\
 \mathcal{I}_{0+}^{(1-\lambda)(1-\mu)} u(0) &= u_0
 \end{aligned} \tag{6}$$

over the interval $[0, T_0]$ in the Banach space \mathbb{U} .

DEFINITION 3.1 The function $u(\zeta)$ is called mild solution of the impulsive fractional equation (6) over the interval $[0, T_0]$ if $u(t)$ satisfies the integral

equation

$$u(\zeta) = \begin{cases} \mathcal{S}_{\lambda,\mu}(\zeta)u_0 + \int_0^\zeta \mathcal{K}_\lambda(\zeta - \tau)f(\tau, u(\tau), Ku(\tau))d\tau, & \zeta \in [0, t_1) \\ g_k(\zeta, u(\zeta)), & \zeta \in [t_k, s_k) \\ \mathcal{S}_{\lambda,\mu}(\zeta - s_k)g_k(s_k, u(s_k)) + \int_0^\zeta \mathcal{K}_\lambda(\zeta - \tau)f(\tau, u(\tau), Ku(\tau))d\zeta, & \zeta \in [s_k, t_{k+1}) \end{cases} \quad (7)$$

for all k .

Following theorem establishes the existence result for the Hilfer fractional integro-differential evolution system with classical condition (6).

THEOREM 3.2 If,

(A1) The evolution operator $-A$ generates C_0 semigroup $S(t)$ for all $t \in [0, T_0]$.

(A2) The function $f : [0, T_0] \times \mathbb{U} \times \mathbb{U} \rightarrow \mathbb{U}$ is continuous with respect to ζ and there exist a positive constants f_1^* and f_2^* such that $\|f(\zeta, u_1, v_1) - f(\zeta, u_2, v_2)\| \leq f_1^*\|u_1 - u_2\| + f_2^*\|v_1 - v_2\|$ for $u_1, v_1, u_2, v_2 \in B_{r_0} = \{u \in \mathbb{U}; \|u\| \leq r_0\}$ for some r_0 .

(A3) The operator $K : [0, T_0] \times \mathbb{U} \rightarrow \mathbb{U}$ is continuous and there exist a constant k^* such that $\|Ku - Kv\| \leq k^*\|u - v\|$ for $u, v \in B_{r_0}$.

(A4) The functions $g_k : [t_k, s_k] \times \mathbb{U}$ are continuous and there exist a positive constants $0 < g_k^* < 1$ such that $\|g_k(\zeta, u(\zeta)) - g_k(\zeta, v(\zeta))\| \leq g_k^*\|u - v\|$.

are satisfied, then the fractional integro-differential system (6) with not-instantaneous impulses has a unique mild solution.

Proof : Define the operator \mathcal{F} on Banach space \mathbb{U} by

$$\mathcal{F}u(t) = \begin{cases} \mathcal{F}_1u(\zeta), & \zeta \in [0, t_1) \\ \mathcal{F}_{2k}u(\zeta), & \zeta \in [t_k, s_k) \\ \mathcal{F}_{3k}u(\zeta), & \zeta \in [s_k, t_{k+1}) \end{cases}$$

where, the operators \mathcal{F}_1 , \mathcal{F}_{2k} and \mathcal{F}_{3k} are defined as

$$\mathcal{F}_1u(\zeta) = \mathcal{S}_{\lambda,\mu}(\zeta)u_0 + \int_0^\zeta \mathcal{K}_\lambda(\zeta - \tau)f(\zeta, u(\tau), Ku(\tau))d\tau, \quad \zeta \in [0, t_1)$$

$$\mathcal{F}_{2k}u(\zeta) = g_k(\zeta, u(\zeta)), \quad \zeta \in [t_k, s_k)$$

$$\mathcal{F}_{3k}u(\zeta) = \mathcal{S}_{\lambda,\mu}(\zeta - s_k)g_k(s_k, u(s_k)) + \int_0^\zeta \mathcal{K}_\lambda(\zeta - \tau)f(\tau, u(\tau), Ku(\tau))ds, \\ \zeta \in [s_k, t_{k+1})$$

for all $k = 1, 2, \dots, p$.

In view of this definition of the operator \mathcal{F} , the equation (7) has a unique solution if and only if the operator equation $u(\zeta) = \mathcal{F}u(\zeta)$ has a unique solution. This is possible if and only if each of $u(\zeta) = \mathcal{F}_1u(\zeta)$, $u(\zeta) = \mathcal{F}_{2k}u(\zeta)$ and $u(\zeta) = \mathcal{F}_{3k}u(\zeta)$ has unique solution over the interval $[0, t_1)$, $[t_k, s_k)$ and $[s_k, t_{k+1})$ for all $k = 1, 2, \dots, p$ respectively.

For all $\zeta \in [0, t_1)$ and $u, v \in B_{r_0}$,

$$\begin{aligned} \|\mathcal{F}_1^{(n)}u(\zeta) - \mathcal{F}_1^{(n)}v(\zeta)\| &\leq \int_0^\zeta \int_0^{\tau_1} \cdots \int_0^{\tau_{n-1}} \|\mathcal{K}_\lambda(\zeta - \tau_1)\| \|\mathcal{K}_\lambda(\tau_1 - \tau_2)\| \cdots \\ &\quad \|\mathcal{K}_\lambda(\tau_{n-1} - s)\| (f_1^* + k^* f_2^*)^n \|u - v\| \\ &\quad ds d\tau_{n-1} \cdots d\tau_1 \end{aligned}$$

Assuming (A1), (A2) and (A3) and using lemma 2.5 and simplifying we get,

$$\begin{aligned} \|\mathcal{F}_1^{(n)}u(t) - \mathcal{F}_1^{(n)}v(t)\| &\leq \int_0^{t_1} \int_0^{t_1} \cdots \int_0^{t_1} \frac{t_1^{n(\lambda-1)} M^n(f_1^* + k^* f_2^*)^n}{(\Gamma(\lambda))^n} \\ &\quad \times \|u - v\| ds d\tau_{n-1} \cdots d\tau_1 \\ &\leq \frac{t_1^{n(\lambda-1)} M^n(f_1^* + k^* f_2^*)^n}{(\Gamma(\lambda))^n (n-1)!} \int_0^{t_1} (t_1 - \tau)^{n-1} \|u - v\| d\tau \\ &\leq \frac{t_1^{n\alpha} M^n(f_1^* + f_2^* k^*)^n}{n! (\Gamma(\alpha))^n} \|u - v\| \\ &\leq c^* \|u - v\|. \end{aligned}$$

Taking limit n tending to ∞ over interval $[0, t_1)$, $\|\mathcal{F}_1^{(n)}u - \mathcal{F}_1^{(n)}v\| \leq c^* \|u - v\| \rightarrow 0$ for fixed t_1 . Therefore, there exist at least one m such that $\mathcal{F}_1^{(m)}$ is contraction on B_{r_0} . Thus, by general Banach contraction theorem the operator equation $u(\zeta) = \mathcal{F}_1u(\zeta)$ has unique solution over the interval $[0, t_1)$.

For all $k = 1, 2, \dots, p$, $\zeta \in [t_k, s_k)$ and $u, v \in \mathbb{U}$ and assuming (A4)

$$\|\mathcal{F}_{2k}u(\zeta) - \mathcal{F}_{2k}v(\zeta)\| = \|g_k(\zeta, u(\zeta)) - g_k(\zeta, v(\zeta))\| \leq g_k^* \|u - v\|.$$

Therefore, \mathcal{F}_{2k} is contraction, and by Banach fixed point theorem the operator equation $u(\zeta) = \mathcal{F}_{2k}u(\zeta)$ has a unique solution for the interval $[t_k, s_k)$ for all $k = 1, 2, \dots, p$. This means for all $k = 1, 2, \dots, p$, $u(\zeta) = g_k(\zeta, u(\zeta))$ has unique solution for all $t \in [t_k, s_k)$. Using Lipschitz continuity of g_k , the solution u is unique at s_k also.

For all $k = 1, 2, \dots, p$, $t \in [s_k, t_{k+1})$ and $u, v \in B_{r_0}$,

$$\begin{aligned} \|\mathcal{F}_{3k}^{(n)}u(\zeta) - \mathcal{F}_{3k}^{(n)}v(\zeta)\| &\leq \int_{s_k}^t \int_{s_k}^{\tau_1} \cdots \int_{s_k}^{\tau_{n-1}} \|\mathcal{K}_\lambda(\zeta - \tau_1)\| \|\mathcal{K}_\lambda(\tau_1 - \tau_2)\| \cdots \\ &\quad \|\mathcal{K}_\lambda(\tau_{n-1} - s)\| (f_1^* + k^* f_2^*)^n ds d\tau_{n-1} \cdots d\tau_1 \end{aligned}$$

Assuming (A1), (A2) and (A3) and using lemma 2.5 and simplifying we get,

$$\begin{aligned} \|\mathcal{F}_{3k}^{(n)} u(\zeta) - \mathcal{F}_{3k}^{(n)} v(\zeta)\| &\leq \int_{s_k}^{t_{k+1}} \int_{s_k}^{t_{k+1}} \dots \int_{s_k}^{t_{k+1}} \frac{(t_{k+1} - s_k)^{n(\lambda-1)} M^n(f_1^* + k^* f_2^*)^n}{(\Gamma(\lambda))^n} \\ &\quad \|u - v\| ds d\tau_{n-1} \dots d\tau_1 \\ \|\mathcal{F}_{3k}^{(n)} u(\zeta) - \mathcal{F}_{3k}^{(n)} v(\zeta)\| &\leq \frac{(t_{k+1} - s_k)^{n(\lambda-1)} M^n(f_1^* + f_2^* k^*)^n}{(n-1)! (\Gamma(\lambda))^n} \\ &\quad \int_{s_k}^{t_{k+1}} (t_{k+1} - s)^{n-1} ds \|u - v\| \\ &\leq \frac{(t_{k+1} - s_k)^{n\lambda} M^n(f_1^* + f_2^* k^*)^n}{n! (\Gamma(\lambda))^n} \|u - v\| \\ &\leq c^* \|u - v\|. \end{aligned}$$

Over interval $[s_k, t_{k+1})$ and taking $n \rightarrow \infty$, $\|\mathcal{F}_{3k}^{(n)} u - \mathcal{F}_{3k}^{(n)} v\| \leq c^* \|u - v\| \rightarrow 0$ for fixed sub-interval $[s_k, t_{k+1})$ for all $k = 1, 2, \dots, p$. Thus, there exist at least one m such that $\mathcal{F}_{3k}^{(m)}$ is contraction on B_{r_0} . Thus by general Banach contraction theorem the operator equation $u(t) = \mathcal{F}_{3k} u(t)$ has unique solution over the interval $[s_k, t_{k+1})$ for all $k = 1, 2, \dots, p$.

Hence, the operator equation $u(\zeta) = \mathcal{F}u(\zeta)$ has a unique solution over the interval $[0, T_0]$ which is nothing but the mild solution of the equation (6). **QED**

EXAMPLE 3.3 Consider the integro-differential Hilfer system of order $(0 < \lambda \leq 1$ and $0 \leq \mu \leq 1)$:

$$\begin{aligned} D_{\zeta}^{\lambda, \mu} u(\zeta, \psi) &= u_{\psi, \psi}(\zeta, \psi) + u(\zeta, \psi) u_{\psi}(\zeta, \psi) + \int_0^{\zeta} e^{-u(\tau, \psi)} d\tau, \\ &\quad \zeta \in [0, 1/3] \cup [2/3, 1] \end{aligned} \tag{8}$$

$$u(\zeta, \psi) = \frac{u(\zeta, \psi)}{2(1 + u^2(\zeta, \psi))}, \quad \zeta \in [1/3, 2/3]$$

over the interval $[0, 1]$ with initial condition $u(0, \psi) = u_0(\psi)$ and boundary condition $u(\zeta, 0) = u(\zeta, 1) = 0$.

The equation (8) can be reformulated as fractional order abstract equation in $\mathbb{U} = L^2([0, 1], \mathbb{R})$ as:

$$\begin{aligned} D^{\lambda, \mu} z(\zeta) &= Az(\zeta) + f(\zeta, z(\zeta), Kz(\zeta)), \quad \zeta \in [0, 1/3] \cup [2/3, 1] \\ z(\zeta) &= g(\zeta, z(\zeta)) \quad \zeta \in [1/3, 2/3] \end{aligned} \tag{9}$$

over the interval $[0, 1]$ by defining $z(\zeta) = u(\zeta, \cdot)$, operator $Au = u''$ (second order derivative with respect to ψ). The functions f and g over respected domains are as $f(\zeta, z(\zeta), Kz(\zeta)) = (z^2(t\zeta))'/2 + \int_0^{\zeta} e^{-z(\tau)} d\tau$ and $g(\zeta, z(\zeta)) = \frac{z(\zeta)}{2(1+z^2(\zeta))}$ respectively.

- (1) The linear operator A over the domain $D(A) = \{u \in \mathbb{U}; u'' \text{ exist and continuous with } u(0) = u(1) = 0\}$ is self-adjoint, compact, and resolvent. Therefore A is the infinitesimal generator of C_0 semi- group $\mathcal{T}(\zeta)$ over the interval $[0, 1]$ given by

$$\mathcal{T}(\zeta)u = \sum_{n=1}^{\infty} \exp(-n^2\pi^2\zeta) \langle u, \phi_n \rangle \phi_n, \quad (10)$$

where $\phi_n(\psi) = \sqrt{2}\sin(n\pi\psi)$ for all $n = 1, 2, \dots$ is the orthogonal basis for the space \mathbb{U} .

- (2) The function $K : [0, 1] \times [0, 1] \times \mathbb{U} \rightarrow \mathbb{U}$ is continuous with respect to ζ and differentiable with respect to z for all z and hence K is Lipschitz continuous with respect to z . This means there exist positive constant k^* such that $\|K(\zeta, z_1) - K(\zeta, z_2)\| \leq k^* \|z_1 - z_2\|$.
- (3) The function $f : [0, 1] \times \mathbb{U} \times \mathbb{U} \rightarrow \mathbb{U}$ is continuous with respect to ζ and is differential with respect to argument z and Kz . Therefore there exist positive constants f_1^* and f_2^* such that $\|f(\zeta, z_1, Kz_1) - f(\zeta, z_2, Kz_2)\| \leq f_1^* \|z_1 - z_2\| + f_2^* \|Kz_1 - Kz_2\|$, $z_1, z_2 \in B_{r_0}$ for some r_0 .
- (4) The impulse g is continuous with respect to ζ and Lipschitz continuous with respect to z with Lipschitz constant $g^* = 1/2 < 1$.

Thus, by theorem-3.2 the equation (9) has unique solution over $[0, 1]$. Hence, the equation (8) has a unique solution over the interval $[0, 1]$.

4. Equation with Nonlocal Conditions This section establishes the existence results for fractional order ($0 < \lambda \leq 1$ and $0 \leq \mu \leq 1$) non-instantaneous impulsive Hilfer integro-differential evolution systems with non-local initial conditions.

$$D_{0+}^{\lambda, \mu} u(\zeta) = -Au(\zeta) + f\left(\zeta, u(\zeta), \int_0^{\zeta} a(\zeta, \tau, u(\tau))d\tau\right),$$

$$\zeta \in \left[\cup [s_i, t_{i+1}] \right] \cup [s_p, T_0] \quad (11)$$

$$u(\zeta) = g_k(\zeta, u(\zeta)), \quad \zeta \in [t_1, s_1) \cup [t_2, s_2) \cup \dots [t_p, s_p)$$

$$u(0) = u_0 + h(u)$$

in the Banach space \mathbb{U} .

DEFINITION 4.1 The function $u(t)$ is called mild solution of the impulsive

fractional equation (11) over the interval if $u(t)$ satisfies the integral equation

$$u(\zeta) = \begin{cases} \mathcal{S}_{\lambda,\mu}(\zeta)[u_0 + h(u)] + \int_0^\zeta \mathcal{K}_\lambda(\zeta - \tau)f(\tau, u(\tau), Ku(\tau))d\tau, & \zeta \in [0, t_1) \\ g_k(\zeta, u(\zeta)), & \zeta \in [t_k, s_k) \\ \mathcal{S}_{\lambda,\mu}(\zeta - s_k)g_k(s_k, u(s_k)) + \int_0^\zeta \mathcal{K}_\lambda(\zeta - \tau)f(\tau, u(\tau), Ku(\tau))d\zeta, & \zeta \in [s_k, t_{k+1}) \end{cases} \quad (12)$$

The following theorem establishes the existence of the solution for the non-local non-instantaneous fractional integro-differential evolution system (11) of Hilfer type over the interval $[0, T_0]$.

THEOREM 4.2 If,

- (B1) *The evolution operator $-A$ generates C_0 semigroup $S(t)$ for all $t \in [0, T_0]$.*
- (B2) *The function $f(\zeta, \cdot, \cdot)$ is continuous and $f(\cdot, u, v)$ is measurable on $[0, T] \times \mathbb{U} \times \mathbb{U}$. Also, there exist $\gamma \in (0, \lambda)$ with $m_f \in L^{\frac{1}{\gamma}}([0, T_0], \mathbb{R})$ such that $|f(\zeta, u, v)| \leq m_f(\zeta)$ for all $u, v \in \mathbb{U}$.*
- (B3) *The operator $K : [0, T_0] \times \mathbb{U} \rightarrow \mathbb{U}$ is continuous and there exist a constant k^* such that $\|Ku - Kv\| \leq k^*\|u - v\|$.*
- (B4) *The operator $h : \mathbb{U} \rightarrow \mathbb{U}$ is Lipschitz continuous with respect to u with Lipschitz constant $0 < h^* \leq 1$.*
- (B5) *The functions $g_k : [t_k, s_k] \times \mathbb{U}$ are continuous and there exist a positive constants $0 < g_k^* < 1$ such that $\|g_k(t, u(t)) - g_k(t, v(t))\| \leq g_k^*\|u - v\|$.*

*are satisfied then the non-local non-instantaneous fractional order integro-differential evolution system (12) has a mild solution provided $M_0^*h^* < 1$ and $M_0^*g^* < 1$.*

Proof: Using the lemma-2.5 and (B4),

$$|\mathcal{S}_{\lambda,\mu}(\zeta)(u_0 + h(u))| \leq \frac{M(\mu(1 - \lambda))\zeta^{\lambda+\mu-\lambda\mu-1}}{\Gamma(\lambda + \mu - \lambda\mu - 1)}(|u_0| + h^*\|u\| + |h(0)|). \quad (13)$$

for all $u \in B_k = \{u \in \mathbb{U} : \|u\| \leq k\}$ for any positive constant k and $\zeta \in [0, T_0]$. Using (B2), $(t - s)^{\lambda-1} \in L^{\frac{1}{1-\gamma}}[0, \zeta]$ for all $\zeta \in [0, T_0]$ and $\gamma \in (0, \lambda)$. Taking $M_1 = \|m_f\|_{L^{\frac{1}{\gamma}}}$ and using Holder's inequality and assuming (B2), for

$\zeta \in [0, T_0]$

$$\begin{aligned} \int_0^\zeta \mathcal{K}_\lambda(\zeta - \tau) f(\tau, u(\tau), Ku(\tau)) ds &\leq \frac{M}{\Gamma(\lambda)} \left(\int_0^\zeta (\zeta - \tau)^{\frac{\lambda-1}{1-\gamma}} ds \right)^{1-\gamma} M_1 \\ &\leq \frac{MM_1(\gamma-1)}{\Gamma(\lambda)(\lambda-\gamma)} \zeta^{\lambda-\gamma}. \end{aligned} \quad (14)$$

For $\zeta \in [0, t_1)$ and for positive r consider F_1 and F_2 on B_r as,

$$\begin{aligned} F_1 u(\zeta) &= \mathcal{S}_{\lambda, \mu}(\zeta)(u_0 + h(u)) \\ F_2 u(\zeta) &= \int_0^\zeta \mathcal{K}_\lambda(\zeta - \tau) f(\tau, u(\tau), Ku(\tau)) d\tau \end{aligned}$$

The function $u(\zeta)$ is a mild solution of the semi-linear fractional integro-differential equation if and only if the operator equation $u = F_1 u + F_2 u$ has a solution for $u \in B_r$ for some r . Therefore the existence of a mild solution of (6) over the interval $[0, t_1)$ is equivalent to determining a positive constant r_0 , such that $F_1 + F_2$ has a fixed point on B_{r_0} .

Step:1 $\|F_1 u + F_2 v\| \leq r_0$ for some positive r_0 .

Let $u, v \in B_{r_0}$ where,

$$\begin{aligned} r_0 &= M_0^* \frac{|u_0| + |h(z)|}{1 - M_0^* h^*} + \frac{M_1^*}{(1 - M_0^* h^*)} t_1^{(\lambda-\gamma)}, \\ M_0^* &= \frac{M(\mu(1-\lambda))t_1^{\lambda+\mu-\lambda\mu-1}}{\Gamma(\lambda+\mu-\lambda\mu-1)}, \end{aligned}$$

and

$$M_1^* = \frac{MM_1(\gamma-1)}{\Gamma(\lambda)(\lambda-\mu)}$$

and considering

$$\begin{aligned} &|F_1 u(\zeta) + F_2 v(\zeta)| \\ &\leq \left| \mathcal{S}_{\lambda, \mu}(\zeta)(u_0 + h(u)) \right| + \left| \int_0^\zeta \mathcal{K}_\lambda(\zeta - \tau) f(\tau, u(\tau), Ku(\tau)) d\tau \right| \\ &\leq \frac{M(\mu(1-\lambda))\zeta^{\lambda+\mu-\lambda\mu-1}}{\Gamma(\lambda+\mu-\lambda\mu-1)} (|u_0| + h^* \|u\| + |h(0)|) \\ &\quad + \frac{MM_1(\gamma-1)}{\Gamma(\lambda)(\lambda-\gamma)} \zeta^{\lambda-\gamma} \\ &\quad \text{(using inequalities (13) and (14))} \\ &\leq r_0 \quad \text{(since, } M_0^* h^* < 1\text{)}. \end{aligned}$$

Therefore, $\|F_1u + F_2v\| \leq r_0$ for every pair $u, v \in B_{r_0}$.

Step: 2 To show F_1 is contraction on B_{r_0} , consider $u, v \in B_{r_0}$ and $\zeta \in [0, t_1)$,

$$\begin{aligned} |F_1u(t) - F_1v(t)| &\leq \left| \mathcal{S}_{\lambda, \mu}(\zeta)(u_0 + h(u)) - \mathcal{S}_{\lambda, \mu}(\zeta)(u_0 + h(v)) \right| \\ &\leq \frac{M(\mu(1-\lambda))\zeta^{\lambda+\mu-\lambda\mu-1}}{\Gamma(\lambda+\mu-\lambda\mu-1)} h^* \|u - v\| \leq M_0^* h^* \|u - v\| \end{aligned}$$

and $M_0^* h^* < 1$ which implies F_1 is contraction.

Step: 3 To show F_2 is completely continuous operator on B_{r_0} , consider the sequence $\{u_n\}$ in B_{r_0} converging to $u \in B_{r_0}$ then,

$$\begin{aligned} |F_2u_n(\zeta) - F_2u(\zeta)| &\leq \int_0^\zeta |\mathcal{K}_\lambda(\zeta - \tau)| |f(\tau, u_n(\tau), Ku_n(\tau)) - f(\tau, u(\tau), Ku(\tau))| d\tau \\ &\leq C^* \int_0^\zeta \|f(\tau, u_n(\tau), Ku_n(\tau)) - f(\tau, u(\tau), Ku(\tau))\| d\tau, \end{aligned}$$

where, $c^* = \frac{M(\mu(1-\lambda))\zeta^{\lambda+\mu-\lambda\mu-1}}{\Gamma(\lambda+\mu-\lambda\mu-1)}$ and using continuity of f with respect to the second and third argument $\|F_2u_n - F_2u\| \rightarrow 0$ as $n \rightarrow \infty$. So, F_2 is continuous. Now to show $\{F_2u, u \in B_{r_0}\}$ is relatively compact it is sufficient to show

- (1) The family of functions $\{F_2u, u \in B_{r_0}\}$ is uniformly bounded and equicontinuous.
- (2) For any $t \in [0, t_1)$, $\{F_2u(t), u \in B_{r_0}\}$ is relatively compact in \mathbb{U} .

Clearly, for any $u \in B_{r_0}$, $\|F_2u\| \leq r_0$, this means that the family $\{F_2u, u \in B_{r_0}\}$ is uniformly bounded in \mathbb{U} .

For any $u \in B_{r_0}$ and $0 \leq \zeta_1 < \zeta_2 < t_1$,

$$\begin{aligned} |F_2u(\zeta_2) - F_2u(\zeta_1)| &= \left| \int_0^{\zeta_2} \mathcal{K}_\lambda(\zeta_2 - \tau) f(\tau, u(\tau), Ku(\tau)) d\tau - \int_0^{\zeta_1} \mathcal{K}_\lambda(\zeta_1 - \tau) \right. \\ &\quad \left. f(\tau, u(\tau), Ku(\tau)) d\tau \right| \\ &= \left| \int_{\zeta_1}^{\zeta_2} \mathcal{K}_\lambda(\zeta_2 - \tau) f(\tau, u(\tau), Ku(\tau)) d\tau + \int_0^{\zeta_1} \mathcal{K}_\lambda(\zeta_2 - \tau) \right. \\ &\quad \left. f(\tau, u(\tau), Ku(\tau)) d\tau - \int_0^{\zeta_1} \mathcal{K}_\lambda(\zeta_1 - \tau) f(\tau, u(\tau), Ku(\tau)) d\tau \right| \\ &\leq \left| \int_{\zeta_1}^{\zeta_2} \mathcal{K}_\lambda(\zeta_2 - \tau) f(\tau, u(\tau), Ku(\tau)) d\tau \right| + \left| \int_0^{\zeta_1} [\mathcal{K}_\lambda(\zeta_2 - \tau) - \right. \\ &\quad \left. \mathcal{K}_\lambda(\zeta_1 - \tau)] f(\tau, u(\tau), Ku(\tau)) d\tau \right| \\ &\leq I_1 + I_2, \end{aligned}$$

where,

$$I_1 = \left| \int_{\zeta_1}^{\zeta_2} \mathcal{K}_\lambda(\zeta_2 - \tau) f(\tau, u(\tau), Ku(\tau)) d\tau \right|$$

$$\leq \frac{MM_1(\gamma - 1)}{\Gamma(\lambda)(\lambda - \gamma)} (\zeta_2 - \zeta_1)^{\lambda - \gamma}$$

(Applying inequality (14) over interval $[\zeta_1, \zeta_2]$),

this implies the integral $I_1 \rightarrow 0$ as $\zeta_1 \rightarrow \zeta_2$. Similarly,

$$I_2 = \left| \int_0^{\zeta_1} [\mathcal{K}_\lambda(\zeta_2 - \tau) - \mathcal{K}_\lambda(\zeta_1 - \tau)] f(\tau, u(\tau), Ku(\tau)) d\tau \right|$$

$$= \left| \int_0^{\zeta_1} [(\zeta_2 - \tau)^{\lambda - 1} \mathcal{Q}_\lambda(\zeta_2 - \tau) - (\zeta_1 - \tau)^{\lambda - 1} \mathcal{Q}_\lambda(\zeta_1 - \tau)] \right.$$

$$\left. f(\tau, u(\tau), Ku(\tau)) d\tau \right|$$

$$\leq \left| \int_0^{\zeta_1} (\zeta_2 - \tau)^{\lambda - 1} [\mathcal{Q}_\lambda(\zeta_2 - \tau) - \mathcal{Q}_\lambda(\zeta_1 - \tau)] f(\tau, u(\tau), Ku(\tau)) d\tau \right|$$

$$+ \left| \int_0^{\zeta_1} [(\zeta_2 - \tau)^{\lambda - 1} - (\zeta_1 - \tau)^{\lambda - 1}] \mathcal{Q}_\lambda(\zeta_1 - \tau) f(\tau, u(\tau), Ku(\tau)) d\tau \right|$$

Assuming the (B1),(B2), (B3) and Holder inequality the integral

$$\left| \int_0^{\zeta_1} (\zeta_2 - \tau)^{\lambda - 1} [\mathcal{Q}_\lambda(\zeta_2 - \tau) - \mathcal{Q}_\lambda(\zeta_1 - \tau)] f(\tau, u(\tau), Ku(\tau)) d\tau \right|$$

and the integral

$$\left| \int_0^{\zeta_1} [(\zeta_2 - \tau)^{\lambda - 1} - (\zeta_1 - \tau)^{\lambda - 1}] \mathcal{Q}_\lambda(\zeta_1 - \tau) f(\tau, u(\tau), Ku(\tau)) d\tau \right|$$

also vanishes as $\zeta_1 \rightarrow \zeta_2$. The vanishing of both the integral lead to the vanishing of I_2 . Thus, $|F_2u(\tau_2) - F_2(\tau_1)|$ tends to zero as $\zeta_1 \rightarrow \zeta_2$ for independent choice of $u \in B_{r_0}$. Hence, the family $\{F_2u, u \in B_{r_0}\}$ is equicontinuous.

Define the family $X(\zeta) = \{F_2u(\zeta), u \in B_{r_0}\}$ for all $\zeta \in [0, t_1]$. Clearly, $X(0)$ is relatively compact. Let, $\zeta_0 \in [0, t_1]$ be fixed and for each $\epsilon \in [0, t_1]$, define an operator F_ϵ on B_{r_0} by formula

$$F_\epsilon u(t) = \int_0^{\zeta - \epsilon} \mathcal{K}_\lambda(\zeta - \tau) f(\tau, u(\tau), Ku(\tau)) d\tau.$$

To show that the family $X(\zeta)$ for all $\zeta \in [0, t_1]$ is relatively compact consider,

$$\begin{aligned} & |F_2u(\zeta) - F_\epsilon u(\zeta)| \\ &= \left| \int_0^\zeta \mathcal{K}_\lambda(\zeta - \tau)f(\tau, u(\tau), Ku(\tau))d\tau \right. \\ &\quad \left. - \int_0^{\zeta-\epsilon} \mathcal{K}_\lambda(\zeta - \tau)f(\tau, u(\tau), Ku(\tau))d\tau \right| \\ &\leq \int_\epsilon^\zeta |\mathcal{K}_\lambda(\zeta - \tau)f(\tau, u(\tau), Ku(\tau))d\tau| \\ &\leq \frac{MM_1(\gamma - 1)}{\Gamma(\lambda)(\lambda - \gamma)}(\zeta - \epsilon)^{\lambda-\gamma} \quad (\text{Applying inequality (14)}). \end{aligned}$$

Thus, $X(\zeta)$ is relatively compact and hence, by Ascoli-Arzela theorem the operator F_2 is completely continuous on B_{r_0} . Using Krasnoselskii's fixed point theorem $F_1 + F_2$ has a fixed point on B_{r_0} which is a mild solution of the equation (11) over the interval $[0, t_1]$.

On the interval $[t_k, s_k)$ for all $k = 1, 2, \dots, p$ and fixed positive r_0 define the operators F_1 and F_2 on B_{r_0} as,

$$\begin{aligned} F_1u(\zeta) &= g_k(\zeta, u(\zeta)) \\ F_2u(\zeta) &= 0 \end{aligned}$$

assuming (B5) using Krasnoselskii's fixed point theorem, $u(\zeta)$ is the mild solution of the non-instantaneous Hilfer integro-differential fractional evolution system over the interval $[t_k, s_k)$.

On the interval $[s_k, t_{k+1})$ for all $k = 1, 2, \dots, p$ and for positive r we define F_1 and F_2 on B_r as,

$$\begin{aligned} F_1u(\zeta) &= \mathcal{S}_{\lambda, \mu}(\zeta - s_k)g_k(s_k, u(s_k)) \\ F_2u(\zeta) &= \int_0^\zeta \mathcal{K}_\lambda(\zeta - \tau)f(\tau, u(\tau), Ku(\tau))d\zeta \end{aligned}$$

then, the function $u(\zeta)$ is the mild solution of Hilfer fractional integro-differential evolution system over the interval $[s_k, t_{k+1})$ if and only if the operator equation $u = F_1u + F_2u$ has a solution for $u \in B_r$ for some r . This is equivalent to the a mild solution of (11) over the interval $[s_k, t_{k+1})$.

Select,

$$\begin{aligned} r_0 &= M_0^* \frac{|u_0| + |g(\cdot, z)|}{1 - M_0^*g^*} + \frac{M_1^*}{(1 - M_0^*g^*)} (t_{k+1} - s_k)_1^{(\lambda-\gamma)}, \\ M_0^* &= \frac{M(\mu(1 - \lambda))(t_{k+1} - s_k)_1^{\lambda+\mu-\lambda\mu-1}}{\Gamma(\lambda + \mu - \lambda\mu - 1)}, \end{aligned}$$

, and

$$M_1^* = \frac{MM_1(\gamma - 1)}{\Gamma(\lambda)(\lambda - \mu)}$$

and applying similar arguments as applied for interval $[0, t_1)$ and using Krasnoselskii's fixed point theorem $F_1 + F_2$ has a fixed point on B_{r_0} which is nothing but the mild solution of the Hilfer fractional integro-differential evolution system (11) over the interval $[s_k, t_{k+1})$. This completes the proof of the theorem. **QED**

EXAMPLE 4.3 Consider, the Hilfer fractional partial integro-differential evolution system with nonlocal conditions:

$$D_{\zeta}^{1/2,1} u(\zeta, \psi) = u_{\psi, \psi}(\zeta, \psi) + u(\zeta, \psi)u_{\psi}(\zeta, \psi) + \frac{1}{50} \int_0^{\zeta} e^{-u(\tau, \psi)} d\tau, \quad \zeta \in [0, 1/3) \cup [2/3, 1] \quad (15)$$

$$u(\zeta, \psi) = \frac{u(\zeta, \psi)}{10(1 + u^2(\zeta, \psi))}, \quad \zeta \in [1/3, 2/3].$$

over the interval $[0, 1]$ with initial condition $u(0, \psi) = u_0(\psi) + \sum_{i=1}^2 \frac{1}{3^i} u(1/i, \psi)$ and boundary condition $u(\zeta, 0) = u(\zeta, 1) = 0$.

The equation (15) can be reformulated as fractional order abstract equation in $\mathbb{U} = L^2([0, 1], \mathbb{R})$ as:

$$D^{1/2,1} z(\zeta) = Az(\zeta) + f(\zeta, z(\zeta), Kz(\zeta)), \quad \zeta \in [0, 1/3) \cup [2/3, 1] \quad (16)$$

$$z(\zeta) = g(\zeta, z(\zeta)) \quad \zeta \in [1/3, 2/3]$$

over the interval $[0, 1]$ by defining $z(\zeta) = u(\zeta, \cdot)$, operator $Au = u''$ (second order derivative with respect to ζ). The functions f and g over respected domains are defined as $f(\zeta, z(\zeta), Kz(\zeta)) = \frac{1}{50} \int_0^{\zeta} e^{-z(\tau)} d\tau$ and $g(\zeta, z(\zeta)) = \frac{z(\zeta)}{10(1+z^2(\zeta))}$ respectively.

The equation (16) satisfies the conditions (B1-B5) of the hypothesis with $M_0^* h^* < 1$ and $M_0^* g^* < 1$. Hence the equation (16) has a mild solution over the interval $[0, 1]$.

5. Conclusion This manuscript established the results concerning the mild solutions of non-instantaneous impulsive fractional integro-differential evolution system on the Banach space \mathbb{U} by considering classical as well as non-local conditions. These results are obtained using the concept of non-linear functional analysis and fixed point theorems. Using these results one can obtain a mild solution for the non-instantaneous impulsive Hilfer fractional integro-differential system.

REFERENCES

- [1] S. Ahmad, A. Ullah, Q. M. Al-Mdallal, H. Khan, K. Shah, and A. Khan. Fractional order mathematical modeling of covid-19 transmission. *Chaos, Solitons & Fractals*, 139:110256, 10 2020. ISSN 09600779. doi:

- [10.1016/j.chaos.2020.110256](https://doi.org/10.1016/j.chaos.2020.110256). URL <https://linkinghub.elsevier.com/retrieve/pii/S0960077920306524>. Cited on p. 3.
- [2] J. Borah and S. N. Bora. Existence of mild solution of a class of nonlocal fractional order differential equation with not instantaneous impulses. *Fractional Calculus and Applied Analysis*, 22:495–508, 4 2019. ISSN 1311-0454. doi: [10.1515/fca-2019-0029](https://doi.org/10.1515/fca-2019-0029). URL <https://link.springer.com/10.1515/fca-2019-0029>. Cited on p. 6.
- [3] M. M. El-Borai. Some probability densities and fundamental solutions of fractional evolution equations. *Chaos, Solitons & Fractals*, 14:433–440, 8 2002. ISSN 09600779. doi: [10.1016/S0960-0779\(01\)00208-9](https://doi.org/10.1016/S0960-0779(01)00208-9). URL <https://linkinghub.elsevier.com/retrieve/pii/S0960077901002089>. Cited on p. 3.
- [4] M. M. El-Borai. Semigroups and some nonlinear fractional differential equations. *Applied Mathematics and Computation*, 149:823–831, 2 2004. ISSN 00963003. doi: [10.1016/S0096-3003\(03\)00188-7](https://doi.org/10.1016/S0096-3003(03)00188-7). URL <https://linkinghub.elsevier.com/retrieve/pii/S0096300303001887>. Cited on p. 3.
- [5] M. Fečkan, J.-R. Wang, and Y. Zhou. On the new concept of solutions and existence results for impulsive fractional evolution equations. *Dynamics of Partial Differential Equations*, 8:345–361, 2011. ISSN 1548159X. doi: [10.4310/DPDE.2011.v8.n4.a3](https://doi.org/10.4310/DPDE.2011.v8.n4.a3). Cited on p. 3.
- [6] R. Hilfer, editor. *Applications of fractional calculus in physics*. Singapore: World Scientific, 2000. ISBN 981-02-3457-0. Zbl 0998.26002. Cited on p. 3.
- [7] A. Jaiswal and D. Bahuguna. Hilfer fractional differential equations with almost sectorial operators. *Differential Equations and Dynamical Systems*, 1 2020. ISSN 0971-3514. doi: [10.1007/s12591-020-00514-y](https://doi.org/10.1007/s12591-020-00514-y). URL <http://link.springer.com/10.1007/s12591-020-00514-y>. Cited on pp. 3, 4, 5, and 6.
- [8] H. Kataria and P. Patel. Existence and uniqueness of non-local cauchy problem for fractional deferential equation on banach space. *International Journal of Pure and Applied Mathematics*, 120:10237, 07 2018. ISSN 10237-10252. Cited on p. 4.
- [9] H. R. Kataria and P. H. Patel. Congruency between classical and mild solutions of caputo fractional impulsive evolution equation on banach space. *International Journal of Advanced Science and Technology*, 29: 1923–1936, 2020. ISSN 2005-4238. Cited on p. 4.

- [10] H. R. Kataria, P. H. Patel, and V. Shah. Existence results of noninstantaneous impulsive fractional integro-differential equation. *Demonstratio Mathematica*, 53:373–384, 12 2020. ISSN 2391-4661. doi: [10.1515/dema-2020-0029](https://doi.org/10.1515/dema-2020-0029). Cited on p. 4.
- [11] A. Meraj and D. N. Pandey. Existence of mild solutions for fractional non-instantaneous impulsive integro-differential equations with nonlocal conditions. *Arab Journal of Mathematical Sciences*, 26:3–13, 8 2020. ISSN 1319-5166. doi: [10.1016/j.ajmsc.2018.11.002](https://doi.org/10.1016/j.ajmsc.2018.11.002). Cited on p. 4.
- [12] K. S. Miller and B. Ross. *An introduction to the fractional calculus and fractional differential equations*. A Wiley-Interscience Publication. John Wiley & Sons, Inc., New York, 1993. ISBN 0-471-58884-9. Cited on p. 3.
- [13] I. Podlubny. *Fractional differential equations*, volume 198 of *Mathematics in Science and Engineering*. Academic Press, Inc., San Diego, CA, 1999. ISBN 0-12-558840-2. An introduction to fractional derivatives, fractional differential equations, to methods of their solution and some of their applications. MR [1658022](https://www.ams.org/mathscinet/item?id=1658022). Cited on p. 3.
- [14] V. Shah, R. K. George, J. Sharma, and P. Muthukumar. Existence and uniqueness of classical and mild solutions of generalized impulsive evolution equation. *International Journal of Nonlinear Sciences and Numerical Simulation*, 19:775–780, 12 2018. Cited on p. 4.
- [15] C.-J. Xu and P.-L. Li. Mild solution of fractional order differential equations with not instantaneous impulses. *Open Mathematics*, 13:null, 2015. URL <http://eudml.org/doc/271049>. Cited on p. 4.
- [16] C. Yu and G. Gao. On the solution of nonlinear fractional order differential equation. *Nonlinear Analysis: Theory, Methods & Applications*, 63:e971–e976, 11 2005. ISSN 0362546X. doi: [10.1016/j.na.2005.01.008](https://doi.org/10.1016/j.na.2005.01.008). Cited on p. 3.

O rozwiązaniu impulsowego ułamkowego układu całkowo-różnicowego Hilfera z opóźnieniem.

Gargi J. Trivedi, Vishant Shah, Jaita Sharma, Rajesh Sanghvi

Streszczenie Artykuł poświęcony jest ułamkowym, z opóźnieniem, systemom ewolucji całkowo-różniczkowej opisanym ułamkowym operatorem różniczkowym Hilfera w przestrzeni Banacha. Analizowane jest istnienie gładkiego rozwiązania równania z warunkami lokalnymi i nielokalnymi. Wyniki uzyskano stosując do operatora C_0 generowanego przez liniową część równania metody nieliniowej analizy funkcjonalnej z twierdzeniami o punkcie stałym. Zamieszczone przykłady podkreślają znaczenie otrzymanych wyników.

Klasyfikacja tematyczna AMS (2010): [034A08, 47H10, 47H20].

Słowa kluczowe: Ułamkowe równanie ewolucji, ułamkowy operator różniczkowy Hilfera, półgrupa operatorów, impulsy opóźnione, twierdzenie o punkcie stałym..



Gargi J. Trivedi is pursuing her Ph.D.(Mathematics) from CVM University V. V. Nagar, Gujarat, India, She is working as a Teaching Assistant in the Department of Applied Mathematics, Faculty of Technology and Engineering, The Maharaja Sayajirao University of Baroda, Vadodara, Her main academic interests are Dynamical Systems, Image processing, Machine learning, and medical image analysis.



Vishant Shah is an Assistant Professor, in the Department of Applied Mathematics, Faculty of Technology and Engineering, The Maharaja Sayajirao University of Baroda, Vadodara, Gujarat, India. His research area is Mathematical Control Theory, Impulsive Dynamical Systems, Functional Analysis, and Orbital Mechanics.



Jaita Sharma is an Assistant Professor, in the Department of Applied Mathematics, Faculty of Technology and Engineering, The Maharaja Sayajirao University of Baroda, Vadodara, Gujarat, India. She is also a Research supervisor in the same Department. Her research area is Mathematical Control Theory, Functional Analysis, and Dynamical Systems.




Rajesh Sanghvi is an Assistant Professor, in G. H. Patel College of Engineering and Technology, CVM University, V V Nagar, Gujarat, India. He is also a Research supervisor in the same Department. His scientific interests include image processing, machine learning, fuzzy logic, optimization, and evolutionary algorithms.

E-MAIL: gargi1488@gmail.com

VISHANT SHAH 
THE M. S. UNIVERSITY OF BARODA, VADODARA
THE M. S. UNIVERSITY OF BARODA, VADODARA
PRATAPGUNJ, VADODARA, GUJARAT 390002, INDIE
E-MAIL: vishantmsu83@gmail.com

JAITA SHARMA 
THE MAHARAJA SAYAJIRAO UNIVERSITY OF BARODA
THE M. S. UNIVERSITY OF BARODA
PRATAPGUNJ, VADODARA, GUJARAT 390002, INDIE
E-MAIL: jaita.sharma@gmail.com

RAJESH SANGHVI 
CHARUTAR VIDYA MANDAL UNIVERSITY
DEPARTMENT OF MATHEMATICS
G. H. PATEL COLLEGE OF ENGINEERING AND TECHNOLOGY
CVM UNIVERSITY, VALLABH VIDHYANAGAR, GUJARAT 388120, INDIA
E-MAIL: rajeshsanghvi@gcet.ac.in

COMMUNICATED BY: Vishant Shah

(Received: 11th of October 2022; revised: 17th of January 2023)
