

ENERGY EXPENDITURE OF SWIMMERS AND SCUBA DIVERS

Przemysław Michniewski

Department of Maritime and Tropical Medicine, Military Medical Institute, Gdynia, Poland

ABSTRACT

This article discusses the energy input and effort of swimmers during long-distance swimming, classifying them according to the definitions of work intensity. It also refers to the diver's effort while performing tasks underwater.

Keywords: effort, swimming, scuba diver, energy value.

ARTICLE INFO

PolHypRes 2019 Vol. 69 Issue 4 pp. 91 – 96

ISSN: 1734-7009 **eISSN:** 2084-0535

DOI: 10.2478/phr-2019-0023

Pages: 6, figures: 0, tables: 5

page www of the periodical: www.phr.net.pl

Publisher

Polish Hyperbaric Medicine and Technology Society

Review article

Article from 1974, unpublished

Acceptance for print in PHR: 17.03.2018 r.

INTRODUCTION

Most authors analysing the burden of work and physical effort in various occupations and sports classify swimming and diving as heavy or very heavy work [1,2,5,6,7,8,9]. One of the elements of the assessment of the workload is to determine the amount of energy expenditure, expressed as the amount of calorie consumption in connection with work performed [3,4].

In the body of a person doing physical work, considerable amounts of heat are generated, just like in a running combustion engine. The energy source of the engine is fuel in the form of petrol or diesel oil, while the energy source for working muscles is the "burning" of carbohydrates, fats and proteins contained in food. The energy released as a result of chemical changes in the body is expressed in units of thermal energy, i.e. in calories or kilocalories (cal, kcal). During the burning of 1g of carbohydrates into carbon dioxide and water, 4.1 kcal are produced (similarly to 1g of protein) and of 1g of fats - 9.3 kcal. Each form of energy can be transformed into other forms, e.g. chemical into mechanical and the latter into thermal, so when considering the amount of energy spent, different units of work, energy and power can be used. For example:

1 cal = 4.186 joules

1 kilocalory (kcal)= 1000 cal kcal/h = 1.163W

1 watt (W) = joule/s = 2.389×10^{-1} cal/s

1 kGm/min = 2.34×10^{-3} kcal/min = 2.19×10^{-4} KM/min

1 KM/min = 1.07×10 kcal/min

A 1 kilowatt bulb, for example, produces heat in the amount of 14.4 kcal/min, which corresponds to the amount of heat produced by 10 adult people who are sitting and busy talking [1]. In physiology, the energy value of food is expressed in kcal. A person at rest uses the energy corresponding to 1 kcal/min. As a source of energy, the human body is an inefficient instrument. Its power per kilogram of body weight is 0.005-0.007 HP/1 HP = 735.5 W. The same size for a combustion engine is 1.4 hp x kg⁻¹. The body's efficiency, understood as the ratio of useful energy to total energy consumed, is 20%, which means that for each kilocalorie spent in the form of work there are 4 kcal spent as heat [4]. The efficiency of the steam engine is 25% and that of the combustion engine 30-40%.

The attached Table I provides data on the energy obtained from the oxidation of nutrients as well as on the RQ ratio, i.e. the ratio of carbon dioxide produced by the organism to oxygen absorbed.

Tab. 1

Energy obtained by oxidation of nutrients [1].

1g	Required quantity of O ₂ per ml	Quantity of produced CO ₂ per ml	RQ	Energy kcal	Calorific equivalent of oxygen
Starch	828,8	828,8	1,000	4,183	5,047
Animal fat	2019,2	1427,3	0,707	9,461	4,686
Protein	966,1	782,7	0,809	4,442	5,600

The overall energy expenditure of the body consists of three components:

- Basal metabolic rate (BMR) defined for a person during physical and mental rest. BMR is about 1 kcal/h/kg body weight. For example, in a man weighing 70 kg BMR = 1680 kcal/24h.
- Functional metabolism (washing, shaving, eating, etc.). This form of metabolism amounts on average to 400 kcal for men and 300 kcal/24h for women.
- Working metabolism (working energy expenditure) closely related to work or sport (swimming, diving) and usually expressed in kcal/min.

In practice, the so-called indirect calorimetry method is used to measure the working output of energy, consisting of measuring the volume of exhaled air and chemical analysis of its composition (O₂, CO₂). If the respiratory quotient RQ is, for example, 0.85, then 1 litre of oxygen consumed corresponds to the production of 4.85 kcal of energy. By multiplying the amount of oxygen consumed in litres by 4.85, the amount of energy released can be calculated. Industrial physicians use the ready-made data in the tables and expressed in kcal/min for various professional activities and sports. The Spitzer-Hettinger energy expenditure tables are in common use [8].

Tables II, III, IV and V below present, for illustration purposes, data defining energy expenditure in the course of particular types of work, data on swimmers crossing the English Channel, energy expenditure and oxygen consumption by divers and scuba divers, and finally energy expenditure depending on style and speed of swimming.

Classification of work in industry [1].

Work	Energy expenditure	
	Men kcal/min/65 kg	Women kcal/min/55 kg
Light	2.0-4.9	1.5-3.4
Medium heavy	5.0-7.4	3.5-5.4
Heavy	7.5-9.9	5.5-7.4
Very heavy	10.0-12.4	7.5-9.4
Extremely heavy	12.5-...	9.5-...

Tab. 3

Contestants, participants in the English channel swimming competition (1955 r.).

Contestants	Age in years	Height in cm	Weight in kg	Body surface in m ²	Speed km/h	Moves per minute	Lung ventilation 1/min	O ₂ consumption 1/min	Energy expenditure kcal/min	Vital capacity of lungs 1
Men										
1.N.B.	58	164.0	91.2	1.98	2.65	42.4	58.3	2.39	11.7	4.2
2.G.G.	25	172.0	82.3	1.96	3.22	50.4	54.6	3.10	15.0	5.8
3.B.P.	34	@160.2	76.5	1.80	3.43	58.2	65.4	3.09	14.4	3.3
4.M.S	25	157.8	63.0	1.62	3.65	66.9	61.4	2.37	11.6	3.2

Width of English Channel 34.5 km Acc. to Pugh et al. [6,7].

Time of swimming 12-17h.

Assuming an average swimming time over the above distance of 14 h, the energy expenditure of the competitors will be between 10,000 and 13,000 kcal.

On 6.08.1970 Kevin Murphy crossed the English Channel back and forth in 35h 10 minutes.

The energy expenditure connected with covering almost 70 km of the route is estimated at 17-21,000 kcal.

Tab. 4

Energy expenditure and oxygen consumption by divers and scuba divers [1].

Experimental conditions	Average oxygen consumption 1/min	Range	Average energy expenditure kcal/min	Scope
Diving in a swimming pool with equipment and a suit				
Sitting, standing	0.26	0.20-0.32	1.3	1.0-1.6
Minimum movement	0.57	0.40-0.73	2.9	2.0-3.7
Maximum movement	1.53	1.24-1.96	7.7	6.2-9.8
Diving in open waters				
Minimum movement	1.08	0.75-1.61	5.4	3.8-8.1
Maximum movement	1.96	1.41-2.35	9.8	7.1-11.8
Underwater swimming in a pool, flippers, without a suit				
Moderate movement	2.32	1.60-2.68	11.6	8.0-13.4
Considerable mobility	3.09	2.65-3.60	15.5	13.3-18.8

Energy expenditure in the function of swimming style and speed.

Style	Speed		Energy expenditure Kcal/h/70 kg
	Km/h	m/min	
Classic	1.85	30.8	410
Freestyle (Crawl)	1.85	30.8	420
On the back	1.85	30.8	500
Freestyle (Crawl)	2.96	49.3	700
Freestyle (Crawl)	4.07	67.9	1600
Classic	4.07	67.9	1850
Classic	4.44	74.0	2530
Classic	4.99	83.2	3690

Comparison of the data contained in Tables I-V allows for a correct assessment of the physical effort of swimmers and scuba divers, which is of great importance in the process of their training.

Knowing the energy expenditure per minute allows for objective assessment of the physical burden of the working system and operation using generally known units such as "cal", "kcal", "kcal/min". Nowadays, when most of the work has been mechanised, the working energy expenditure is in the range of 500-3000 kcal/day and will probably decrease due to further technological progress. The situation is different in sports. It is observed from year to year and from one Olympics to another that the existing records associated with long-term physical effort such as swimming, marathon running, cycling marathons, cross-country skiing etc. are constantly being broken. Achieving by swimmers within a dozen or so hours of energy expenditure of 12-17000 kcal or even more, shows the existence of hitherto unknown possibilities of energy expenditure by the human organism.

Work with an energy output of more than 10 kcal/min, described as very heavy according to the data from Table No. II, is performed by the manual worker in the course of their normal working day over short periods of time. The load on a long-distance swimmer with an energy output of 15 kcal/min lasts several hours without interruption. Unlike a combustion engine, a human being can work without fuel (food), burning his or her own energy resources in the form of glycogen and adipose tissue, naturally over a limited time. However, a healthy and fully functioning body requires a constant supply of food, equivalent to the calorific value of the energy spent.

What should be the daily calorific demand of a swimmer at an energy output of e.g. 12000 kcal/day?

$$\text{Daily quantity of calories in food} = \frac{\text{working energy expenditure} + \text{functional energy expenditure} + \text{BMR}}{0,88}$$

kcal

A value of 0.88 in the denominator indicates that only 88% of the consumed food is assimilated and can provide energy.

$$\text{Daily calorific demand of a swimmer} = \frac{12000 + 400 + 1700}{0,88} \text{ kcal} = 16022 \text{ kcal.}$$

In the extremely hard-working body of a swimmer, the universally applicable law on energy conservation has been violated. The energy expenditure

has far exceeded the supply, because despite taking food while swimming, it is impossible to assimilate in the same time 16,000 kcal (1.8 kg fat or more than 4 kg protein in meat). The energy balance of such a highly stressed body is balanced out in a few or more days. Hence the conclusion that a repetition of such an effort can take place after a sufficiently long break.

CONCLUSIONS

- Determination of energy expenditure in commonly known physical units such as cal, kcal, kcal/min allows for objective, comparative assessment of physical effort in professional work, military service and sports.
- The physical effort of swimmers and scuba divers, assessed on the basis of their energy expenditure, is considered to constitute heavy and extremely heavy work.
- The knowledge of the energy expenditure ranges for different jobs and activities allows military commanders and coaches in sport to efficiently gauge the effort of competitors.
- The knowledge of energy expenditure per minute makes it possible to compare the physical effort of a swimmer, runner or mountaineer and to estimate their calorific demand.

REFERENCES

1. Durnin J., Passmore R.: Energy, work and leisure. PWN, Warsaw 1969;
2. Falls H.: Exercise physiology. Acad. Press, N. York 1968;
3. Kozłowski S.: Physical effort physiology. PZWL, Warsaw 1970;
4. Krause M.: An overview of ergonomics. PZWL, Warszawa 1970;
5. Michniewski P.: Lekarz Wojsk. 1971, 6; 554;
6. Morehouse L., Miller A.: Physiology of exercise. Mosby Co, S. Louis 1967;
7. Pugh L. et al.: Clin. Sci. 1960, 19; 257;
8. Pugh L., Edholm O.: Lancet, 1955, 8; 733;
9. Spitzer-Hettinger: Tafeln für den Kalorienumsatz bei körperlicher Arbeit. Beuth Vertrieb GmbH, Berlin 1964.