

A reassessment of the Pennsylvanian lycophyte cone *Triplosporite* Brown

BARRY A. THOMAS¹ AND JIŘÍ BEK²

¹*Institute of Biological, Environmental and Rural Sciences, Aberystwyth University, Aberystwyth, Ceredigion SY231NP, UK.*

E-mail: bat@aber.ac.uk

²*Laboratory of Palaeobiology and Palaeocology, Institute of Geology, Academy of Sciences v.v.i., Rozvojová 269, 165 00 Prague 6, Czech Republic.*

E-mail: mrbean@gli.cas.cz

ABSTRACT:

Thomas, B.A. and Bek, J. 2014. A reassessment of the Pennsylvanian lycophyte cone *Triplosporite* Brown. *Acta Geologica Polonica*, **64** (2), 139–145. Warszawa.

The type collection of the lycopsid cone species *Triplosporite brownii* Unger was re-examined to assess its *in situ* spores. The cones are monosporangiate with only microspores that possess both cingulum and zona. They equate to the dispersed microspore genus *Lycospora* and would be identified as *Lycospora* cf. *pseudoannulata*. Therefore, the genus *Triplosporite* Brown is shown to be a junior synonym to *Lepidostrobus* and a species emendation is given. A comparison is given with the other *Lepidostrobus* cones which yielded similar *in situ* microspores of the *Lycospora pellucida* Group.

Key words: *Lepidostrobus*; *Lycospora*; *in situ* spores; *Triplosporite*; Cones; Carboniferous.

INTRODUCTION

For over 159 years the lycophyte cone genus *Lepidostrobus* Brongniart was accepted to contain species that were wholly microsporangiate and others that were bisporangiate containing both megaspores and microspores. Thomas (1970, 1978) and Thomas and Dytko (1980) suggested that the microspores, referred to the genus *Lycospora* (Schopf, Wilson and Bentall) Potonié and Kremp, extracted from microsporangiate species of *Lepidostrobus* have both cingulum and zona, differing from those produced by bisporangiate cones.

The lectotype of *Lepidostrobus*, *L. ornatus* Brongniart, was shown to have only microspores that had a cingulum and a developed zona (both 2.5–3 µm wide, see

Bek 2012) with ornamented proximal and distal surfaces (Brack-Hanes and Thomas 1983). From this evidence *Lepidostrobus* was restricted to microsporangiate cones that yielded cingulate and cingulizionate microspores of the *Lycospora*-type, whereas the genus *Flemingites* Caruthers was redefined to include those bisporangiate cones that were originally referred to *Lepidostrobus*. The *Lycospora* microspores in *Flemingites* do not have a zona and possess prominent densely microspinulate distal surface. The proximal sculpture is usually laevigate, sometimes microverrucate to microgranulate (Bek 2012).

Within the systematic descriptions in Brack-Hanes and Thomas (1983) the microsporangiate *Triplosporite brownii* Unger was referred to the genus *Flemingites* on

the basis of the structure of its microspores figured by Brown (1848). By examining the drawings of the microspores in Brown's paper Brack-Hanes and Thomas (1983) came to the conclusion that the spores had only a narrow cingulum and therefore included *Triplosporites brownii* in *Flemingites* naming it *Flemingites brownii* (Unger) comb. nov. They did not see Brown's preparations.

Brown (1848) first used the name *Triplosporite* in describing the upper part of a petrified cone from an unknown locality in the Carboniferous of France. Later Brown (1851) described the cone as having bractae (sporophylls) with sporangia on their upper surfaces "filled with innumerable microscopic sporules, originally connected in threes (very rarely in fours) but ultimately separating...". It is because he thought the majority of the sporules (microspores) to be in threes that Brown called it *Triplosporite* to distinguish the cone from those genera having spores in "constant quadruple union" i.e. in tetrads. Brown gave no diagnosis for *Triplosporite* but did remark that it "approaches most nearly, among recent tribes, to *Lycopodiaceae* and *Ophioglosseae*; and among fossils, no doubt, to *Lepidostrobus*, and consequently to *Lepidodendron*." In an addendum to the paper, Brown accepted Joseph Hooker's observation (Hooker 1843) that some microspores in a specimen of *Lepidostrobus* appeared to be in threes and also noted the general acceptance of Brongniart's (1828–1838) view of *Lepidostrobus* being the fructification of *Lepidodendron* Sternberg. Nevertheless, Brown hesitated from referring *Triplosporite* to *Lepidostrobus* on the grounds that he believed the structure of the latter to be imperfectly known.

Although Brown gave no diagnosis for *Triplosporite* we regard it as being validly published. A few years later, Unger (1850) gave a Latin diagnosis of what he now called *Triplosporites* Rob. Brown in his *Genera et Species* (p. 270) and a single sentence diagnosis for his solitary species *Triplosporites brownii*. The generic diagnosis included Unger's details of the sporophylls and sporangia and that the spores were in threes and occasionally fours. The species diagnosis was based on the structure of the sporangia "Capsules lenticulares compressae, obcordatae, v. reniformes, acuminatae." The species was later referred to *Lepidostrobus brownii* by Schimper (1870).

This produced a taxonomic problem that was missed at the time by Brack-Hanes and Thomas (1983), because, if the two genera were synonyms as they suggested, *Triplosporite* Brown (1845, 1850) predated *Flemingites* Carruthers (erected in 1865) it has priority. To resolve this problem, the slides of *Triplosporite brownii*, which are in the Natural History Museum,

London (numbers V10980, V10980a-b, V1326, V1326 a-s), were examined.

The sporangia were full of microspores [Text-fig. 1a, 2] approximately 28–36 µm in diameter. However, the slide sections were thick enough to make focusing on the spores difficult and the preservation of many spores was not perfect showing very indistinct outlines and ornamentation. Many were still in tetrads that in some views could be mistaken for clusters of three as Brown had assumed [Text-figs 1c, d]. Other spores were better preserved and clearly showed equatorial structures 3–5 µm wide, consisting of a cingulum and probably a perforated zona [Text-figs 2d]. It was difficult to be certain about the ornamentation of the exine, but the distal surface is microspinulate to microgranulate and proximal sculpture seems to be laevigate. The structure of these spores shows that *Triplosporite* is part of a microsporangiate cone and not bisporangiate as in *Flemingites*. Therefore, *Triplosporite* is a junior synonym of *Lepidostrobus*.

MATERIAL AND METHODS

There is no hand specimen of the cone because it has been completely cut into sections. Most sections were transverse although the more apical part of the cone was cut longitudinally. Measurements of the sporangia, sporophyll pedicels and laminae were made by combining information from transverse and longitudinal sections. The spores had to be examined in the slides because there was no opportunity to extract them for closer study. Specimen is stored in the Natural History Museum, London, UK (No. V.10980c).

The spores are classified according to the system of dispersed spores suggested by Potonié and Kremp (1954, 1955) and improved by Dettmann (1963) and Smith and Butterworth (1967). The terms used for the description of the morphology, including the sculptural elements follows the Punt *et al.* (2007) classification. The species determination is based only on these original diagnoses, and not on the interpretations of subsequent authors. Measurements of the holotypes of dispersed species of this group, *in situ* microspores of this type and their parent cones are given in Table 1.

In situ spores are generally accepted to be as an integral part of the diagnoses of any fructification and as part of this the spores are compared with known dispersed species of spores; preferably with the types of the dispersed spore species (Thomas 1987). The details of the *in situ* spores in *Triplosporite brownii* must, therefore, be included in the emended diagnosis for the cone.

PENNSYLVANIAN LYCOPHYTE CONE

Dispersed spores (holotypes)					
Dispersed species	Diameter (µm)	Width of cingulum (µm)	Width of zona (µm)	Stratigraphic level of holotype	Reference
<i>Lycospora pellucida</i>	39 × 46	2	2-4	Bolsovia	Somers <i>et al.</i> 1972
<i>Lycospora pseudoannulata</i>	32 × 35	2	3-4	Langsettian	Somers <i>et al.</i> 1972
<i>Lycospora intermedia</i>	47 × 49	2	3-4	Bolsovia	Somers <i>et al.</i> 1972
<i>Lycospora loganii</i>	32 × 34	2	2	Duckmantian	Somers <i>et al.</i> 1972
In situ spores					
Parent plant	Diameter (µm)	Width of cingulum (µm)	Width of zona (µm)	Stratigraphic level	Reference
<i>Lepidostrobus fayettevillense</i>	42-50	Not measured	Not measured	Pendleian	Taylor and Eggert, 1968
<i>Lepidostrobus barnsleyensis</i>	20-35	1.5-2.5	2-4.5	Duckmantian	Thomas, 1965
<i>Lepidostrobus spinosus</i>	21-39	1.5	5	Duckmantian	Thomas, 1965
<i>Lepidostrobus binneyanus</i>	20-35	2.5	4	Duckmantian	Thomas, 1970
<i>Azaniadendron fertile</i>	21-36	Not measured	Not measured	Early Permian	Rayner, 1986
<i>Lepidostrobus oldhamius</i> (associated with <i>Lepidophloios harcourtii</i>)	29-41.6	2-3	4-11	Langsettian-Duckmantian	Willard, 1989
<i>Lepidostrobus</i> sp. C	33-37	2	3-5	Duckmantian	Hagemann, 1966
<i>Lepidostrobus brownii</i>	28-36	2-3.3	2.5-4	Mississippian	Herein

Table 1. Measurements and stratigraphic ranges of holotypes of dispersed and *in situ* spores of the *Lycospora pellucida* Group (from Bek 2012) and their parent cones

SYSTEMATIC PALAEOLOGY

Class Lycopsidea Scott, 1909

Order Lepidocarpaceae Thomas and Brack-Hanes, 1984

Genus *Lepidostrobus* Brongniart, 1828

TYPE SPECIES: *Lepidostrobus ornatus* Brongniart, 1828

Lepidostrobus brownii (Unger, 1850) Schimper 1870
(Text-figs 1–2)

1848. *Triplosporite* sp.; Brown 1848, p. 344.

1850. *Triplosporite brownii* Unger; Unger, p. 270.

1851. *Triplosporite* sp.; Brown, pls 23, 24.

1870. *Lepidostrobus brownii* Unger; Schimper, pl. 62: figs 13, 14, 16–19, 21–22, (V13236), 15 (V1098c), 17 (V13236a), 19 and 20 (V10980a), 23–26.


1894 *Lepidostrobus brownii* Unger; Bower, pl. 47, p. 103.

HOLOTYPE: Brown, 1848, pl. 23, fig. A.


TYPE LOCALITY: Unknown locality in France.

STRATIGRAPHY: Mississippian (see Brown 1848).

EMENDED DIAGNOSIS: Microsporangiata cone, c. 54 mm in diameter, central cone axis 3.1 mm in diameter with vascular traces 80 µm across. Sporangia 12–14 in each whorl. Sporangia 13 mm long and 7 mm broad and 3 mm high at the distal end. Sporangial wall c. 11 µm thick. Sporophyll pedicel 900 µm thick, dipping slightly at end of sporangium with lamina slightly divergent from the vertical. Lamina c. 7 mm broad, c. 5 mm thick, with a single vascular bundle c. 60 µm in diameter. Trilete circular to subcircular microspores, 28–36 µm in diameter. Cingulum 2 (2.6) 3 µm wide, developed as a dark ring on the outer margin of the central body. Zona is 2.5 (3.2) 4 µm in width in form of lighter ring on the outer margin of cin-



Text-fig. 1. Sections of *Triplosporite browni* (Unger, 1850). A – Transverse section of the cone (No. V.10980e), scale bar 10 mm; B – Tangential longitudinal section of the apical part of the cone (No. V.10980k), scale bar 10 mm; C – Close to a radial longitudinal section of the apical part of the cone (No. V.10980r), scale bar 10 mm; D – Enlargement of b showing sporangia and sporophylls in transverse section, scale bar 5 mm; E – Central stele of the cone (No. V.10980e), scale bar 3 mm



Text-fig. 2. A – Section of sporangial wall and microspore tetrads. B, C – Microspore tetrads; D – Spore showing dark cingulum and outer zona. Scale bars 20 μm

gulum. Proximal surface laevigate, distal surface microspinate to microgranulate. Zona pitted or perforated. Rays of trilete mark extend to the inner margin of cingulum.

COMPARISON

Zeiller (1911) gave a detailed account of a specimen in the Paris Natural History Museum, France that he named *Lepidostrobus brownii* (Unger) Schimper and suggested that it might even have been a part of the same cone described by Brown. However the specimen described by Zeiller is bisporangiate with microsporangia in the upper part and megasporangia in the lower part. The microspores (his pl. 11, figs 15, 17, 18) show no sign of a zona, therefore being of the type found in the genus *Flemingites*. For these reasons we cannot accept that Zeiller's specimen belongs to *Lepidostrobus*

brownii as we have redefined it here. At this stage we do not give Zeiller's specimen a new name other than to suggest that it does not equate to any described species of *Flemingites*. Chaloner (1967) included several other species as synonyms of *L. brownii* to: *L. dabadianus* Schimper, *L. rouville* Renault and Saporta and *L. laurentia* Zeiller (1907). On the available evidence we cannot accept these as synonyms because nothing is known about their *in situ* spores.

In comparing the *in situ* spores of *Lepidostrobus brownii* with dispersed and other known *in situ* microspores of the genus *Lycospora* we determine that the microspores are of the cingulizionate type and belong to the *Lycospora pellucida* Group (Bek 2012). These microspores are typified by having a relatively broad cingulum and zona that can be pitted or perforated. Measurements of holotypes of dispersed species of this group, *in situ* microspores of this type and their parent cones are in Table 1. The microspores can be

equated to the dispersed spore *Lycospora* cf. *pseudoanulata* Kosanke. They differ from the original diagnosis (Kosanke 1950, p. 45) in having a wider and more prominent cingulum.

Microspores macerated from *Lepidostrobus binneyanus* Arber by Thomas (1970) are similar in dimensions and morphology to those described here, but the sculpture of their surfaces is different, i.e. distal surfaces of *L. binneyanus* microspores are laevigate and not microgranulate. *In situ* microspores isolated from *Lepidostrobus fayettevillense* Taylor and Eggert by Taylor and Eggert (1968) and those macerated from *Lepidostrobus barnsleyensis* Thomas by Thomas (1965) differ in prominent perforations of the zona and larger diameter. Microspores described by Hagemann (1966) from *Lepidostrobus* sp. C do not possess pitted or perforated zona. Willard (1989) macerated microspores of this type from cones identified as *Lepidostrobus oldhamius* Williamson (associated with *Lepidophloios harcourtii* Witham), which differ in having a much wider cingulum and zona (10 µm on average). *In situ* microspores from *Lepidostrobus spinosus* Kidston described by Thomas (1965) possess different sculpture, narrow cingulum and wider zona. Microspores from the Permian cone *Azaniodendron fertile* Rayner possesses different sculpture, lack perforations of the zona and were isolated not from mono- but a bisporangiate cone (Rayner 1986).

Acknowledgement

J.B. acknowledges financial support from the Grant Agency of the Academy of Sciences of the Czech Republic (IAA210/12/2053), the Research Program of the Institute of Geology AS CR v.v.i. (AVOZ30130516 and RVO67985831). We also thank Dr. Peta Hayes (Natural History Museum, London, UK) for supplying the photographs of the cone sections.

REFERENCES

- Bek, J. 2012. A review of the genus *Lycospora*. *Review of Palaeobotany and Palynology*, **174**, 122–135.
- Bower, F.O. 1894. Studies in the Morphology of spore producing members. Part 1: Equisetineae and Lycopodineae. *Philosophical Transactions of the Royal Society*, **185B**, 473–572.
- Brack-Hanes, S.D. and Thomas, B.A. 1983. A re-examination of *Lepidostrobus*. *Botanical Journal of the Linnean Society*, **86**, 125–133.
- Brongniart, A. 1828–1838. Histoire des végétaux fossiles, ou recherches botaniques et géologiques sur les végétaux renfermés dans les diverses couches du globe. 1, 1–488; 2: 1–72. Dufour et d' Ocagne; Paris.
- Brown, R. 1848. Some Account of *Triplosporite*, an undescribed Fossil Fruit. *Proceedings of the Linnean Society*, London **1**, 344–345.
- Brown, R. 1851. Some account of *Triplosporite*, an undescribed fossil fruit. *Transactions of the Linnean Society*, London **20**, 469–475.
- Carruthers, W. 1865. On an undescribed cone from the Carboniferous beds of Airdrie, Lanarkshire. *Geological Magazine*, **2**, 433–440.
- Chaloner, W.G. 1967. *Lycophyta*. In: Boureau E. (Ed.), *Traité de Paléobotanique II*. 243–497. Masson et Cie; Paris.
- Dettmann, M.E. 1963. Upper Mesozoic microfloras from south-eastern Australia. *Proceeding of the Royal Society of Victoria*, **77**, 1–148.
- Hagemann, H.W. 1966. Sporen aus köhlig erhaltenen Lepidophytenzapfen des Westfals. *Fortschritte der Geologische von Rheinland und Westphalen*, **13**, 317–388.
- Hooker, J.B. 1843. Remarks on the structure and affinities of some *Lepidostrobus*. *Memoires of the Geological Survey of Great Britain*, **2**, 440–456.
- Kosanke, R.M. 1950. Pennsylvanian spores of Illinois and their use for correlation. *Bulletin of the Illinois State Geological Survey*, **74**, 1–128.
- Potonié, R. and Kremp, G. 1954. Die Gattungen der Paläozoischen *Sporae dispersae* und ihre Stratigraphie. *Geologisches Jahrbuch*, **69**, 111–193.
- Potonié, R. and Kremp, G. 1955. Die *Sporae dispersae* des Ruhrkarbons, ihre Morphographie und Stratigraphie mit Ausblicken auf Arten anderer Gebiete und Zeitabschnitte. Teil I. *Palaeontographica*, **B98**, 1–136.
- Punt, W. Hoen, P.P. Blackmore, S. Nilsson, S. and LeThomas, A. 2007. Glossary of pollen and spore terminology. *Review of Palaeobotany and Palynology*, **143**, 1–81.
- Rayner, R.J. 1986. A new genus of lycopod from South Africa. *Review of Palaeobotany and Palynology*, **47**, 129–143.
- Schimper, W.P. 1870. *Traité de Paléontologie végétale, La flore du monde primitif dans ses rapports avec les formations géologiques de la flore du monde actuel*. Ballière. Paris. vol. 2.
- Schimper, W.P. 1874. *Traité de Paléontologie Végétale, la Flore du Monde Primitif*. Vol. III and Atlas. Paris.
- Scott, D.H. 1909. *Studies in Fossil Botany*, 2nd edition. Adam and Charles Black, London.
- Somers, Y., Alpern, B., Doubinger, J., Grebe, H., 1972. Revision du genre *Lycospora* Schopf, Wilson & Bentall. In: Alpern, B., Streel, M. (Eds.), *Les Spores*, **5**, 9–110.
- Smith, A.H.V. and Butterworth, M.A. 1967. Miospores in the coal seams of the Carboniferous of Great Britain. *Palaeontological Special Papers*, **1**, 1–324.

- Taylor, T.N. and Eggert, D. 1968. Petrified plants from the Upper Mississippian of North America II. *Lepidostrobus fayettevillense* sp. nov. *American Journal of Botany*, **55**, 306–313.
- Thomas, B.A. 1965. Some studies on Carboniferous lycopods. Ph.D. thesis, Reading University. [unpublished]
- Thomas, B.A. 1970. A new specimen of *Lepidostrobus binneyanus* from the Westphalian B of Yorkshire. *Pollen et Spores*, **12**, 217–234.
- Thomas, B.A. 1978. The use of in-situ spores for defining species of dispersed spores. *Review of Palaeobotany and Palynology*, **51**, 227–233.
- Thomas, B.A. and Dytko, A. 1980. *Lepidostrobus haslingdenensis*: a new species from the Lancashire Millstone Grit. *Geological Journal*, **15**, 137–142.
- Unger, F. 1850. *Genera et Species Plantarum Fossilien*. Wilhelm Baumüller, Wien.
- Willard, D.A. 1989. Source plants for Carboniferous microspores: *Lycospora* from permineralized *Lepidostrobus*. *American Journal of Botany*, **76**, 820–826.
- Zeiller, M.R. 1907. Sur quelques *Lepidostrobus* de la region Pyrénéenne. *Compte Rendu de l'Academie des Sciences, Paris*, **145**, 1122–1126.
- Zeiller, M.R. 1911. Étude sur le *Lepidostrobus brownii* (Unger) Schimper. Bureau des Longitud, de l'École Polytechnique; Paris.

Manuscript submitted: 18th November 2013

Revised version accepted: 10th April 2014