

Karolina Kafarska, Wojciech M. Wolf

Lodz University of Technology, Faculty of Chemistry, Institute of General and Ecological Chemistry
116 Żeromskiego Street, 90-924 Lodz, Poland, karolina.kafarska@p.lodz.pl

NOVEL SILVER COMPLEXES WITH POPULAR NON-STEROIDAL ANTI-INFLAMMATORY DRUGS

Abstract

Non-steroidal anti-inflammatory drugs (NSAID) are class of drugs with antipyretic, analgesic and anti-inflammatory properties. They have also exhibited anti-tumor activity. Even though the mode of their anti-inflammatory activity action is well understood, they exhibit significant adverse effects. Metal complexation with NSAID may be a promising option for side effects reduction. The novel silver complexes with commonly used non-steroidal anti-inflammatory drugs: ibuprofen, naproxen, mefenamic acid and ketoprofen, were synthesized and characterized by elemental analysis, IR- spectroscopy and thermal decomposition techniques. Coordination of ligands to the silver ions was confirmed by IR spectroscopy. IR data clearly indicate that NSAID anions are bonded in a monodentate mode. The thermal behavior of complexes was studied by TG, DTG and DTA methods in air. Upon heating all compounds decomposed progressively to silver oxide, which was the final product of pyrolysis.

Key words

Non-steroidal anti-inflammatory drugs, metal complexes, FTIR, thermal investigations





Introduction

Non-steroidal anti-inflammatory drugs are among the oldest [1] and most popular drugs in contemporary medicine [2, 3]. The precursors of non-steroidal anti-inflammatory drugs were salicylic acid and salicylates, isolated from natural sources, have long been used as medicines. The acid was chemically synthesized in 1860 and was further used as an antiseptic, an antipyretic, and an antirheumatic agent. Almost 40 years later, aspirin was developed as a more palatable form of salicylate. Soon after, several drugs having similar action to aspirin were discovered, and the whole group was named the 'aspirin-like drugs'. It was growing quite rapidly and more recently got a new name i.e. the nonsteroidal anti-inflammatory drugs (NSAID)[4]. In 1971 Vane showed that NSAID hampers biosynthesis of prostaglandins (PG) by inhibiting activity of an enzyme – cyclooxygenase (COX) and suggested mechanism of their action [4-5]. Further investigations lead to the discovery of two isoforms of COX. COX-1 is a constitutive enzyme which prompts PG synthesis further protecting stomach and kidney. On the other hand, COX-2 is induced by inflammatory factors, like cytokines, and produces PG that contribute to the pain and inflammation swelling [6-7]. Therefore, drugs which are selective COX-2 inhibitors should show pronounced anti-inflammatory action without side effects on the kidney and stomach. Additionally, selective inhibitors have much wider therapeutic potential including ovulation and premature labors. Several NSAID delay progress of the Alzheimer's disease [8-9]. The quite recently discovered protective action of NSAID on cancer follows from their action on COX-2 [10-15]. Unfortunately, the widely prescribed NSAID to control pain and inflammation are gastrointestinal toxic and may lead to skin allergies as well as cardiovascular and renal diseases in particular when administrated over a long time and without the usual care [16-20].

It is quite well recognized, that several transition metal complexes with non-steroidal drugs are more effective and show lower toxicity, than their parent drugs [1,21-22]. Recently Banti and Hadjikakou documented anti-cancer activity of metal complexes with non-steroidal anti-inflammatory drugs [23]. Silver ion is also known as antibacterial and antifungal agent with low toxicity as compared to the other transition metal [24]. Moreover, silver complexes containing various ligands exhibit selectivity against variety of cancer cells [25]. In this respect, coordination chemistry is an useful method to improve the overall efficiency of NSAID and in the same time reduce their well-recognized side effects .

This paper describes synthesis, spectroscopic and thermal properties of silver complexes with four commonly used in medicine NSAID [26-33], namely ibuprofen (ibup), naproxen (napx), mefenamic acid (mef) and ketoprofen(ket) (Table 1).

Table 1. The structural formulae and popular application of chosen NSAID

| NSAID | Chemical formula | Typical use |
|----------------|---|--|
| Ibuprofen |  |) pain) inflammation) headache) toothache |
| Naproxen |  |) arthritis) ankylosing) spondylitis) tendinitis |
| Mefenamic acid |  |) menstrual pain |
| Ketoprofen |  |) rheumatoid) osteoarthritis) acute, chronic pain |

Source: own compilation on the basis of [26-33]

Materials and measurements

Pure ligands were received as gift from Polish pharmaceutical companies: Pabianickie Zakłady Farmaceutyczne Polfa S.A., Medana Pharma S.A. and Emo-farm Sp. z o.o.; EtOH p.a. was purchased from Aldrich, MeOH from Lab-Scan, all other chemicals were from POCh- Gliwice.

All complexes were obtained according to similar procedures. The first step of synthesis was preparation sodium salt of particular ligand by dissolution of ibuprofen, naproxen, ketoprofen or mefenamic acid (1 mmol) in 50 ml freshly precipitated NaOH aqueous- ethanol solution (1:1) ($0,02 \text{ mol} \cdot \text{L}^{-1}$). The mixture was heated up to 60°C and added to aqueous solution of silver nitrate (1 mmol in 25 mL). The reaction mixture was kept in 60°C for 2 hours. After several days the solid precipitates were isolated by filtration, washed with hot water and dried on air.

The composition of complexes was determined by chemical and elemental analyses. The contents of N, C and H were established using automatic Carbo-Erba analyser in a pure oxygen over V_2O_5 catalyst. The amount of metal was determined by complexometric titration of a mineralized sample. An open system with nitric(V) acid was applied. IR spectra were recorded on FTIR-8501 Shimadzu spectrophotometer over $4000\text{Z}400 \text{ cm}^{-1}$ range using KBr pellets. Thermal stabilities were studied by TG and DTG techniques with the Netzsch TG 209 apparatus. Samples (10mg) were heated in ceramic crucibles up to 1000°C , at a heating rate $10^\circ\text{C min}^{-1}$ in air atmosphere. The solid decomposition products were identified on TG and DTG curves and further confirmed by the X-ray powder diffraction of sinters (Siemens D-5000 diffractometer, graphite monochromatised CuK_α radiation).

Results and discussion

Empirical formulae of complexes augmented by relevant analytical data are summarized in Table 2. All synthesized complexes were obtained as highly crystalline powders. The latter was unequivocally confirmed by X-ray diffraction. They were stable in air and practically insoluble in water but quite well soluble in popular polar organic solvents, like EtOH, MeOH and acetone.

Table 2. Empirical formulae of complexes augmented by relevant analytical data

| Complex | Analysis: found (calculated) /% | | | |
|----------|---------------------------------|----------------|------------------|----------------|
| | Ag | N | C | H |
| Ag(ibup) | 34.36 (34.45) | - | 49.77 (49.86) | 5.50 (5.48) |
| Ag(napx) | 32.00 (31.99) | - | 49.58 (49.88) | 3.78 (3.87) |
| Ag(mef) | 31.00 (30.98) | 4.15 (4.02) | 52.08 (51.75) | 4.05 (3.99) |
| Ag(ket) | 29.44 (29.87) | - | 53.06 (53.21) | 3.70 (3.64) |

Source: Author's

Coordination of ligands to the silver ions was confirmed by IR spectroscopy. In particular, characteristic bands from the valence vibration of carboxyl group were not observed. On the contrary, there appeared bands from asymmetric ($1567\text{-}1585\text{ cm}^{-1}$) and symmetric ($1375\text{-}1384\text{ cm}^{-1}$) vibrations of dissociated COO^- group. These bands are affected by the coordination of ligands to silver ions. According to Nakamoto criteria [34], the separation $\zeta \rightarrow \rightarrow (\text{OCO}) \rightarrow (\text{OCO})$ and the direction of the band shifts in comparison to corresponding values in parent sodium salt characterize the nature of metal – carboxylate bond. The bathochromic shifts of asymmetric (\rightarrow) and hypsochromic of symmetric (\Rightarrow) frequencies were observed. All relevant data are summarized in Table 3. They clearly indicate, that carboxylate groups in all synthesized compounds coordinated as monodentate. The IR spectra of synthesized complexes are collected on Fig. 1-4.

Table 3. Principal IR bands (cm^{-1}) for carboxylate groups in synthesized complexes and sodium salts of ligands

| Compound | ν_{asym} | ν_{sym} | $\Delta\nu = \nu_{\text{asym}} - \nu_{\text{sym}}$ |
|----------|---------------------|--------------------|--|
| Na(ibup) | 1548,7 | 1411,8 | 136,9 |
| Ag(ibup) | 1567,9 | 1384,7 | 183,2 |
| Na(napx) | 1547,0 | 1407,1 | 139,9 |
| Ag(napx) | 1564,7 | 1394,5 | 170,2 |
| Na(mef) | 1580,0 | 1380,0 | 200,0 |
| Ag(mef) | 1584,7 | 1375,4 | 209,3 |
| Na(ket) | 1567,0 | 1394,0 | 173,0 |
| Ag(ket) | 1577,5 | 1384,7 | 192,8 |

Source: Author's




Fig.1. IR spectra of Ag(ibup)

Source: Author's




Fig.2. IR spectra of Ag(napx)
Source: Author's

Additionally, mefenamic and ketoprofen anions have amine and carbonyl group, respectively. Potentially they can be available for coordination. The characteristic IR bands of these groups in free ligands are very close to those observed in silver complexes. That indicates, that neither the NH group in Ag(mef) nor CO in Ag(ket) participate directly in coordination.




Fig.3. IR spectra of Ag(mef)
Source: Author's




Fig.4. IR spectra of Ag(ket)
Source: Author's

The data obtained from TG, DTG curves supported by chemical and X-ray diffraction analysis are collected in the Table 3. The thermal decomposition curves of complexes are shown on Fig. 5-8.

The thermal decomposition of synthesized complexes is quite similar. Pyrolysis is a two-stages process for all complexes. Almost all compounds are thermal stable up to 180°C, only for Ag(ket) higher thermal resistance was detected (220°C). Further increase of temperature (in ranges 180-550°C) results in high mass loss (*ca.* 50%) caused by organic ligand decomposition. Further heating in the terminal step of pyrolysis leads to metallic silver formation. The whole mass loss 65,5% for Ag(ibup), 68% for Ag(napx), 69% for Ag(mef) and Ag(ket) is in good agreement with calculated amounting to 65,55%, 68,01%, 69,02% and 70,13% respectively for ibuprofenato, naproxenato, mefenamato and ketoprofenato complex.

Table 3. Thermal decomposition data of complexes in air

| Compound | Ranges of decomposition /°C | Mass loss /% | | Intermediate and final product |
|----------|-----------------------------|--------------|------------|--------------------------------|
| | | found | calculated | |
| Ag(ibup) | 180Z250 | 47.5 | 46.70 | AgOAc |
| | 250Z400 | 18.5 | 18.85 | Ag |
| Ag(napx) | 180Z480 | 51,0 | 50.50 | AgOAc |
| | 480Z700 | 17.0 | 17.51 | Ag |
| Ag(mef) | 180Z550 | 52.0 | 52.06 | AgOAc |
| | 550Z710 | 17.0 | 16.96 | Ag |
| Ag(ket) | 220Z480 | 53.0 | 53.78 | AgOAc |
| | 480Z640 | 16.0 | 16.35 | Ag |

Source: Author's




Fig. 5 Thermoanalytical profiles for Ag(ibup)

Source: Author's




Fig. 6. Thermoanalytical profiles for Ag(napx)
Source: Author's




Fig. 7. Thermoanalytical profiles for Ag(mef)
Source: Author's




Fig. 8. Thermoanalytical profiles for Ag(ket)
Source: Author's

Presence of silver, as a final product of thermolysis, was confirmed by powder X-ray diffraction of sinters. They were prepared by heating complexes up to relevant temperature (from TG curve). The prominent XRD reflections (Fig. 9 and 10) compared with CCDC model corroborated, that the product is metallic silver [35].




Fig. 9. XRD patterns of final product of pyrolysis of Ag(ibup) (1) and Ag(napx) (2)

Source: Author's




Fig. 10. XRD patterns of final product of pyrolysis of Ag(mef) (1) and Ag(ket) (2)

Source: Author's

Conclusion

The silver complexes with ibuprofen, naproxen, mefenamic acid and ketoprofen with formulae: Ag(ibup), Ag(napx), Ag(mef), Ag(ket) (Scheme 1) were synthesized and characterized.



Fig. 11. Formula of complexes proposed according to elemental, IR and TG/DTG data

Source: Author's

IR vibrational spectra confirmed coordination bond formation between NSAID carboxylate groups, and silver ions and indicated that ibuprofenato, naproxenato, mefenamato and ketoprofenato anions act as monodentate ligands (on the contrary to Ag(I) complex with ibuprofen presented by Pereira E Silva [36]). The thermal investigations demonstrated that all synthesized complexes are highly thermal- proof, very closely to free ligands. Additionally they were stable over time.

References

- [1] Vonkeman H.E., van de Laar M.A.F.J., Nonsteroidal Anti-Inflammatory Drugs: Adverse Effects and Their Prevention, *Semin. Arthritis Rheu.* 39 (4) (2010) 294-312
- [2] Wolfe M., Lichtenstein D., Gastrointestinal toxicity of nonsteroidal antiinflammatory drugs, *New Eng. J. Med.*, 340 (1999) 1888-1899
- [3] Lee M., Feldman M. The aging stomach: implications for NSAID gastropathy, *Gut* 41 (1997) 425–426
- [4] J.R.Vane, Mechanism of action of non-steroidal anti-inflammatory drugs, *Am.J.Med.* 104A (1998) 2S-8S
- [5] J.R. Vane, R.M. Botting, The mechanism of action of aspirin, *Thromb. Res.*, 110 (2003) 255-258
- [6] V.S.Moses, A.L.Bertone, Nonsteroidal, anti-inflammatory drugs, *Vet. Clin.Equine* 18(1) (2002) 21-37
- [7] Claria J., Cyclooxygenase-2 Biology, *Curr. Pharm. Design*, 9 (2003) 2177-2190
- [8] Etminan M., Gill S., Samii A., Effect of non-steroidal antiinflammatory drugs on risk of Alzheimer's disease: systematic review and meta-analysis of observational studies, *BMJ.*, 327, 128-31, (2003)
- [9] Szekely C.A., Town T., Zandi P.P., NSAIDs for the chemoprevention of Alzheimer's disease, *Subcell Biochem*, 42, 229-48, (2007)
- [10] A. Ianaro, G. Cirino, J.L. Wallance, Hydrogen sulfide-releasing anti-inflammatory drugs for chemoprevention and treatment of cancer, *Pharm. Res.* 111 (2016) 652-658
- [11] R. Kumar, P.F. Sirlil, F. Javid, Unusual anti-leucemia activity of naproxen and other non-steroidal anti-inflammatory drugs, *Mat.Sci.Eng. C-Matter*, 69 (2016) 1335-1344
- [12] K.W. Cheng, T. Nie, N. Ouyang, L.Huang, B. Rigas, A novel ibuprofen derivative with anti-lung cancer properties: Synthesis, formulation, pharmacokinetic and efficacy studies, *Int. J. Pharm.* 477 (2014) 236-243
- [13] M. S. M. Al-Nimer, H.G. Hamed, M.M. Mahmood, Antiproliferative effect of aspirin and other NSAIDs against the grow of cancer and fibroblast cells, *Saudi. Pharm. J.* 23 (5) (2015) 483-486
- [14] M. Vasooghi, M. Amini, The discovery and development of cyclooxygenase – 2 inhibitors as potential anticancer therapies, *Expert Opin. Drug Dis.* 9(3) (2014) 255-267
- [15] I.H. Sahin, M.M. Hassan, C.R. Garrett, Impact of non-steroidal anti-inflammatory drugs on gastrointestinal cancers, current state- of- the science, *Cancer Lett.* 345(2) (2014) 249-257
- [16] J. Benmoussa, M. Chevenon, M. Nandi, M. Forlenza, J. Nfonoyim, *Am. J. Emerg. Med.* 34(5) (2016) 942e5-942e7
- [17] I.L.Meek, M.A.F.J van de Laar, H.E. Vonkeman, Non-steroidal, anti-inflammatory drugs: An overview of cardiovascular risc, *Pharmaceuticals* 3(7) (2010) 2146-2162
- [18] G.H. Gislason, NSAIDs and cardiovascular risc, *Am. Fam. Psysician*, 80(12) (2009) 1366-1368
- [19] A.Riser, D. Donovan, J Haintzman, T. Page, NSAID Prescribing Precautions, *Am. Fam. Psysician*, 80(12) (2009) 1371-1378
- [20] A. Husjan, P. Ahuja, A. Ahmad, S.A. Khan, Synthesis, biological evaluation and pharmacocinetic studies of mefenamic acid-N-hydroxymethylsuccimide ester prodrug as safer NSAID, *Med. Chem.* 12(6) (2016) 585-591

- [21] F. Dimiza, F. Rerdih, V. Tangouli, I. Turel, D. Kessissoglou, G. Psoms, Interaction of copper(II) with non-steroidal anti-inflammatory drugs naproxen and diclofenac: Synthesis, structure, DNA-and albumin-binding, *J. Inorg. Biochem.*, 105 (2011) 476-489
- [22] J. Feng, X. Du, H. Liu, X. Sui, C. Zhang, Y. Tang, J. Zhang, Manganese-mefenamic acid complexes exhibit high lipoxygenase inhibitory activity, *Dalton Trans.*, 43, (2014) 10930
- [23] C.N. Banti, S.K. Hadjikakou, Non-steroidal anti-inflammatory drugs (NSAIDs) in metal complexes and their effect at the cellular level, *Eur.J.Inorg.Chem.* (2016) 3048-3071
- [24] S. Rafique, M. Idrees, A. Nasim, H. Akbar, A. Athar, Transition metal complexes as potential therapeutic agents, *Biotechnol. Mol. Biol. Rev.*, 5(2), 38-45 (2010)
- [25] C.N. Banti, S.K. Hadjikakou, Anti-proliferative and anti-tumor activity of silver(I) compounds, *Metallomics*, 5(2013) 569-596
- [26] E.A. Sandilands, D.N. BATERMAN, Non-steroidal anti-inflammatory drugs, *Medicine*, 44(3) (2016), 185-186
- [27] A. Viegas, J. Manso, M.C. Corvo, M.B. Marques E.J. Cabrita, Binding of Ibuprofen, Ketolac and Diclofenac to COX-1 and COX-2 Studied by Saturation Transfer Difference NMR, *J. Med. Chem.*, 54 (2011) 8555-8562S
- [28] P.M. Brooks, Nonsteroidal Antiinflammatory Drugs- Differences and similarities, *N. Engl. J. Med.*, 324, 1716-1725 (2007)
- [29] C. Patrono, B. Rocca, Non-steroidal, anti-inflammatory drugs: Past, present, future, *Pharm. Res.*, 59 (2009) 258-289
- [30] L. Abuhijleh, J. Khalaf, Copper(II) complexes of the anti-inflammatory drugs naproxen and 3-pyridylmethanol as auxiliary ligand. Characterization, superoxide dismutase and catecholase- mimetic activities, *Europ. J. Med. Chem.*, 45 (2010) 3811-3817
- [31] J. Feng, X. Du, H.Liu, Y. Tang, J. Zhang, Manganese-mefenamic acid complexes exhibit high lipoxygenase inhibitory activity, *Dalton Trans.*, 43(28) (2014), 10930-10939
- [32] A. Topaçli, S. Ide, Molecular structures of metal complexes with mefenamic acid, *J. Pharm. Biomed. Anal.* 21 (1999) 975-982
- [33] A. L. Ong, A. H. Kamaruddin, S. Bhatia, Current technologies for the production of (S)-ketoprofen: Process perspective, *Process Biochem.*, 40 (2005) 3526-3535
- [34] K. Nakamoto, *Infrared and Raman Spectra of Inorganic and Coordination Compounds*, Wiley, New York 2009
- [35] International Centre for Diffraction Data (ICDD), reference code : 00-001-1164
- [36] I.M.Pereira E Silva, D. De Moraes Profirio, R.E.F. De Paiva, M. Lancellotti, A.L. Barboza Formiga, P.P.Corbi, , A silver complex with ibuprofen: Synthesis, solid state characterization, DFT calculations and antibacterial assays *J.Mol.Struct.*1049 (2013) 1-6