

The usefulness of subfossil Cladocera remains in Younger Dryas climatic reconstructions in central Poland

DOMINIK PAWŁOWSKI

*Institute of Geology, Adam Mickiewicz University, ul. B. Krygowskiego 12, 61-680 Poznań, Poland.
dominikp@amu.edu.pl*

ABSTRACT:

Pawłowski, D. 2017. The usefulness of subfossil Cladocera remains in Younger Dryas climatic reconstructions in central Poland. *Acta Geologica Polonica*, **67** (4), 567–584. Warszawa.

Cladoceran-based paleotemperature estimates for the Younger Dryas for ten sections of paleo-oxbow lakes, valley mires, and lacustrine sediments from central Poland are presented, and their potential usefulness to climatostratigraphy is described. Most of the changes observed in the cladoceran assemblages are responses to climate changes. The cladoceran-based temperature reconstructions reflect cold conditions during the Younger Dryas (YD) and allow a division of this period into two phases: an older colder phase, between 12,800 and 12,000 cal yrs BP, and a younger, warmer phase, between ca. 12,000 and 11,500 cal yrs BP. The geomorphological features of the study sites and local environmental forces are also taken into consideration. The cladoceran-inferred summer temperature estimates from all of the study sites correspond closely with the available climate reconstruction for the YD in central Europe.

Key words: Younger Dryas; Cladocera; Temperature; Oxbow lake; Climate.

INTRODUCTION

Since the later decades of the twentieth century, subfossil Cladocera analysis has been one of the most important methods in paleoecological studies (Frey 1986; Pawłowski 2011). The cladocerans (Branchiopoda, Cladocera) are the most abundant crustaceans preserved in lake sediments (Birks and Birks 1980). The group has been widely used as a proxy for reconstructing paleoclimates, as well as ecological and hydrological changes (Korhola and Rautio 2001; Mirosław-Grabowska *et al.* 2015), because their ecological preferences are relatively well understood. The response of cladocerans to changes in temperature and humidity is significantly more rapid than that of other proxies, e.g., pollen (Birks and Ammann 2000). Cladocerans can be used to quantitatively infer paleotemperature fluctuations and provide a reliable alternative method if a regional

transfer function is not available (Nevalainen *et al.* 2012; Pawłowski *et al.* 2015a).

The Younger Dryas (YD), being a transition from the Last Glaciation to the Holocene, is characterized by a series of rapid climatic changes. This has had a strong influence on local ecosystems and the aquatic invertebrate fauna (Szeroczyńska 1985; Goslar *et al.* 1998; Isarin and Bohncke 1999; Isarin and Rensen 1999; Milecka *et al.* 2011; Feurdean *et al.* 2014; Mirosław-Grabowska and Zawisza 2013; Gandouin *et al.* 2016; Stivrins *et al.* 2016). The river valleys of central Europe underwent significant geological and hydrological transformation (Starkel 1990, 2002; Macklin *et al.* 2006; Starkel *et al.* 2007; Notebaert and Verstraeten 2010; Kaiser *et al.* 2012; Turner *et al.* 2013). In response to temperature fluctuations, this caused biotic changes and alterations in river environments (Pawłowski *et al.* 2015b, c). However, knowledge of European mires in river valleys is relatively lim-



Text-fig. 1. Locality map of the sections studied

ited, despite the fact that most mires (e.g., in central Poland) are located in river valleys (Żurek 1987). Oxbow lake and valley mire sediments provide comparable alternatives to lake and mire sedimentary deposits (Millet *et al.* 2012) for the study of past environmental change (Gandouin *et al.* 2007; Engels *et al.* 2008; Pawłowski *et al.* 2012).

Studies of cladoceran-inferred paleotemperature in the YD and Early Holocene valley mires of central Poland have already begun (Pawłowski *et al.* 2015a, 2016a). Although a few studies have used cladoceran-inferred paleotemperatures in the Late Glacial and Holocene, there have been no comparisons of Cladocera reconstructions in different geological and geomorphological situations in a region.

This study presents data from a number of sites situated in a range of geographical settings in a specific region with the aim of reconstructing climate variability during the YD. The aim of the study is thus to answer the question of how to compare the climate signal coming from lakes and mires located in river valleys with those on plateaus of glacial and aeolian origin in the same region, and thereby how to compare this climatic signal with that on a wider regional scale. This study thus also explores the reliability of cladoceran-inferred summer temperatures for reconstructing past climatic conditions.

STUDY AREA


The investigated sites are located in the Łódź Region, central Poland (Text-fig. 1), which possesses a good record of the Saalian Glaciations, par-

ticularly the Wartanian Cold Stage of the Odranian Glaciation. All sites are located at approximately the same altitude (100–200 m a.s.l.; Table 1), but represent various geological and geomorphological situations (Forysiak 2012). There are lakes located (1) in river valleys – both active and in fossil valleys – and (2) on plateaus of both glacial and aeolian origin. Most of the lakes represent sites located in Polish river valleys: (1) Świerczyna, Pawłowa, Grabica, and Ldzań are on the valley floor of the Grabia River, (2) Korzeń is in the lower reaches of the Widawka River, (3) Kolonia Behcice is in the middle reach of the Ner River, and (4) Koźmin is in the middle reach of the Warta River (Text-fig. 1). The Grabia, Ner, and Widawka Rivers are semi-natural, partly unregulated tributaries of the Warta River (within the Odra River catchment). Most of the floodplains in these river valleys are occupied by a mire with peat pools and paleo-oxbow lakes, which are overgrown by riparian forests; they are located in cut-off paleochannels and also have different surface areas. Geologically, these river valleys are mostly filled with Late Glacial deposits and Holocene alluvium. The Saalian till is present on surrounding plains; locally it is overlain by glaciofluvial sands, and gravels surround the valley reaches (Forysiak 2012; Pawłowski *et al.* 2016a).

The two river valleys sites described here, Koźmin and Ługi, are unique. Koźmin represents a site with a fossil forest that persisted in the YD, in the potentially favourable growing conditions in the floodplain of the Warta River (Dzieduszyńska *et al.* 2014). The Ługi mire represents sites located in the fossil Jadwiczna-Pichna River valley (part of the Warta River valley), which functioned as a proglacial water outflow during the Saalian Glaciation recession and as the main tract of the Warta River Valley in the Late Plenivistulian (Klatkova and Załoba 1991; Forysiak 2012).

The two other sites, at Żabieniec and Rąbień, are located in the morainic uplands near Łódź (Text-fig. 1; Table 1). Żabieniec is on a plateau to the east of Łódź, in an oval depression surrounded by morainic hills (Forysiak 2012). This kettle hole was probably formed during the Late Saalian (Wartanian) and has a very irregular bottom configuration. The oldest sediments recorded were deposited at the end of the Upper Pleniweichselian. Rąbień, in turn, is located in the morainic Łask Upland to the west of Łódź (Text-fig. 1). This basin of aeolian origin is situated in an oval depression surrounded by dunes, which was formed during the Older Dryas (Forysiak 2012).

Detailed studies of the geology and relief of



Text-fig. 2. Percentages of Cladocera from the Rąbień, Pawłowa, Kolonia Bechcice, Koźmin, Korzeń, and Żabieniec sites

Site name	Latitude	Longitude	Altitude (m a.s.l.)	Location	Length of core (cm) / ca. length of studied period (cm)	Resolution of analysis (cm)
Grabica	51°29'11" N	19°32'12" E	215	active river valleys	200 / 80	4
Ldzań	51°35'31" N	19°13'58" E	173	active river valleys	100 / 44	4
Pawłowa	51°30'19" N	19°19'59" E	184	active river valleys	450 / 160	4
Świerczyna	51°28'02" N	18°59'89" E	146	active river valleys	336 / 176	4
Kolonia Behcice	51°45'12" N	19°14'26" E	154	active river valleys	139 / 30	4
Korzeń	51°28'44" N	18°53'25" E	139	active river valleys	200 / 90	10
Koźmin	52°04'51" N	18°40'30" E	97	active river valleys	50 / 30	4
Ługi	51°43'52" N	18°42'46" E	123	fossil river valley	300 / 150	10
Rąbień	51°48'20" N	19°18'05" E	189	plateau of aeolian origin	620 / 65	4
Żabieniec	51°51'01" N	19°46'38" E	180	plateau of glacial origin	1600 / 80	4

Table 1. Sites location and basic sampling information

the area, with analysis of the sediments from these sites using the multiproxy approach and radiocarbon datings, have been presented by Forysiak (2012), Pawłowski (2012a, b), Dzeduszyńska *et al.* (2014), Pawłowski *et al.* (2014, 2015a, b, 2016a, b).

The sites are no more than about 80 km apart from each other. They are thus characterized by the same climate condition. The climate is transitional (Woś 1999) with more continental than oceanic influences. Mean annual rainfall ranges from 500 to 600 mm; the growth period lasts for 210 days. The mean monthly air temperature varies from -3°C in January to 17.9°C in July, while the mean annual temperature is 7.6–8.0°C (Kłysik 2001). Snow cover is present for an average of 50 to 70 days annually.

MATERIALS AND METHODS

The study sites have previously undergone Cladocera analysis (Pawłowski 2012a, 2012b; Dzeduszyńska *et al.* 2014; Pawłowski *et al.* 2015a, b, 2016a, b). However, the cladoceran-inferred paleotemperature estimates from most study sites – Żabieniec, Rąbień, Korzeń, Ługi, Kolonia Behcice, and Koźmin – have not yet been presented. A new paleotemperature estimation from the YD was therefore constructed on the basis of the previous methods and is presented here. The results (Text-figs 2–4) concern only the YD periods, as the subfossil concentration was either too low or the remains were absent from the upper core section following peat accumulation (the low abundance of Cladocera made a cladoceran-inferred reconstruction impossible). The cladoceran-inferred temperature reconstructions are thus presented for this period.

Cladocera analysis

Sediment cores for the cladoceran studies were collected from all sites (Text-fig. 1) where the thickest organic deposits were found. In the laboratory, the cores were sampled at 4-cm; only two cores were taken at 10-cm resolution (Table 1). The sediment samples for Cladocera analysis were processed according to the standard procedure (Frey 1986); slides were prepared from 0.1 ml of each sample and examined with a microscope (100× magnification). The taxonomy of the cladoceran remains in this paper follows that presented by Szeroczyńska and Sarmaja-Korjonen (2007), and the ecological preferences of the cladoceran taxa were determined based on Bjerring *et al.* (2009). The results were plotted on a percentage diagram using POLPAL software (Walanus and Nalepka 1999).

Stratigraphy


The chronology of the sediments from the study sites was determined on the basis of AMS ¹⁴C dates (Table 2) and pollen biostratigraphy. Radiocarbon dates are limited to the YD period and are given in calibrated years BP; conventional radiocarbon dates were calibrated using the newest version of OxCal 4.2.3 (Bronk Ramsey 2009) and the IntCal13 calibration curve (Reimer *et al.* 2013). Although the YD period is presented here, the age-depth models for all studied sites were constructed based on all dates from the whole cores (for details, see the references in Table 2).

Cladoceran-inferred temperature reconstruction

A Finnish Cladocera-based mean July air temperature (T_{July}) transfer function was applied to

Site	Depth b.g.l. (cm)	¹⁴ C age (BP)	Calibrated age (95.40% probability) cal. BP	Source
GRAB	115	9150±50	10,486–10,225	Pawłowski <i>et al.</i> 2016a
	136	10,160±70	12,096–11,406	
	160	10,340±120	12,568–11,717	
	200	10,420±90	12,573–11,998	
LDZ	56	9930±50	11,605–11,234	Pawłowski <i>et al.</i> 2016a
	79	10,700±60	12,725–12,563	
	95	11,290±120	13,409–12,884	
PAW	208	9900±80	11,700–11,189	Pawłowski <i>et al.</i> 2016b
	264	10,140±80	12,079–11,395	
	336	11,690±120	13,759–13,292	
ŚWIERCZYNA	121	9500±50	11,080–10,931 (32.3%)	Pawłowski <i>et al.</i> 2015a, 2015b
			10,879–10,646 (59.2%)	
			10,630–10,589 (3.9%)	
	148	10,010±70	11,799–11,785 (0.6%)	
			11,774–11,250 (94.8%)	
	176	10,150±90	12,124–11,389 (94.7%)	
			11,377–11,353 (0.7%)	
	196	10,130±90	12,080–11,341	
	232	10,370±80	12,535–11,964	
272	10,360±60	12,514–12,496 (0.9%)		
		12,423–11,984 (94.5%)		
328	10,420±60	12,535–12,076		
KB	190–194	9520±90	11,149–10,581	Pawłowski 2012b; Płóciennik <i>et al.</i> 2016
	220	11,020±230	13,384–12,544	
KOŹMIN	200–204	10,000±80	11,821–11,240	Dzieduszyńska <i>et al.</i> 2014
	228–232	10,570±50	12,629–12,410	
	232–236	10,710±60	12,744–12,547	
	236–240	10,850±60	12,897–12,599	
KO	110	9710±110	11,323–10,715	Forysiak 2012; Pawłowski 2012b
ŁUGI	89–90	7430±90	8394–8043	Forysiak 2012
	140	10,110±130	12,225–11,244	
	279–281	13,820±130	17,130–16,302	
RAB	405	10,060±40	11,805–11,355	Forysiak 2012; Michczyńska <i>et al.</i> 2014
	475	11,180±45	13,150–12,962	
	505	11,625±45	13,569–13,353	
ŻAB	826	9130±50	10,479–10,468 (1.2%)	Forysiak 2012; Pawłowski 2012a
			10,423–10,204 (94.2%)	
	1036	11,860±60	13,858–13,479	

Table 2. Radiocarbon dating of samples from the studied sites (GRAB: Grabica; LDZ: Ldzań; PAW: Pawłowa KB: Kolonia Bechcice; KO: Korzeń; RAB: Rąbień; ŻAB: Żabieniec)




Text-fig. 3. Percentages of Cladocera from the Świerczyna and Ldzań sites

the Cladocera assemblages from all the study sites to create a quantitative estimate of past July air temperatures (Nevalainen *et al.* 2012). A weighted averaging-partial least squares regression (WA-PLS) technique model was used. The cladoceran-based T_{July} inference model had an r_{jack}^2 of 0.67, RMSEP of 0.86°C, and mean and maximum biases of -0.017°C and 1.732°C, respectively (Luoto *et al.* 2011).

The reliability of the temperature reconstructions was tested by MAT (modern analogue analysis) and

goodness-of-fit (GoF) analysis (Brooks *et al.* 2012) and evaluated by examining the squared residual distance of the modern and fossil passive samples in a canonical correspondence analysis (CCA), with the environmental variable of interest as the sole constraining variable (lack-of-fit N 10% of the extreme values in the modern calibration set). It was tested to see whether the samples have good modern analogs (threshold for good assemblage N 10% chord distance). CANOCO and C2 software, respectively, were used for these procedures (ter Braak and Šmilauer 2002; Juggins 2007).



Text-fig. 4. Percentages of Cladocera from the Ługi and Grabica sites

RESULTS

Cladoceran-inferred temperature reconstruction

The cladoceran-inferred mean July temperature (T_{July}) at the sites located in the river valleys (Kozmin, Ługi, Korzeń, Pawłowa, Ldzań and Grabica) reached a high of approximately 14.8°C (Text-fig. 5; Table 3) at the end of the Allerød. Similar values were reached at the upland hill sites (Rąbień and Żabieniec). In the YD, the mean summer temperature in the river valleys decreased to between 13.9°C to 14.7°C (Table 3). In turn, the upland hill sites had an average T_{July} of 14.3°C . The mean summer temperature at all studied sites decreased from ca. 12,800 cal yrs BP to ca. 12,500–12,000 cal yrs BP, when the lowest values were found (Text-fig. 5). From the mid-YD (ca. 12,000 cal yrs BP), all sites show an increasing trend for T_{July} and higher values in the late YD (ca. 11,500 cal yrs BP). In the Early Holocene, the cladoceran-based T_{July} showed higher temperatures than in the YD at all study sites. This indicates an increase of up to 15.7°C in the river valleys (Warta and Grabia valleys), and 15.3°C in the uplands (Żabieniec) (Table 3).

In the cladoceran-based mean July temperature reconstruction, only ca. 25% samples from the sites show a good fit to temperature (Text-fig. 6). For this reason, even though there are good modern analogs

in the modern training set, most of the cladoceran-inferred temperatures must be treated with caution.

DISCUSSION

Cladoceran-inferred temperature reconstitution and regional comparison

The YD cooling is visible in the cladoceran-based temperature reconstructions at all the central Poland study sites (Table 3). The temperature reconstructions from lakes located on active river valley beds show a drop in the July temperature (starting from 12,800 cal yrs BP) and reflect the beginning of the YD cooling (Text-fig. 5). Between ca. 12,500 and 12,300 cal yrs BP, the record from the river valleys is characterized by decreases in summer temperatures from 14°C to 12°C . After that, summer temperatures show a pattern fluctuating between 12.2°C and 16.5°C (with an average of 14.3°C). Finally, the summer temperature increases to 15°C at the end of the YD. The estimates from lakes located on (both active and fossil) river valley beds correlate well with the estimates from plateaus, whether of glacial (Żabieniec) or aeolian origin (Rąbień) (Table 3; Text-fig. 5). This confirms that the temperature estimates from study sites are consistent. These reconstructions are also similar to

	Upland Hills		River Valleys							
	Rąbień this study	Żabieniec this study	Koźmin this study	Ługi this study	Korzeń this study	Kolonia Behcice this study	Świerczyna (Pawłowski <i>et al.</i> , 2015a)	Grabica (Pawłowski <i>et al.</i> , 2016a)	Ldzań (Pawłowski <i>et al.</i> , 2016a)	Pawłowa (Pawłowski <i>et al.</i> , 2016b)
EH	15.1°C	15.3°C	15.7°C	15.1°C	14.9°C	15.3°C	14.7°C	14.8°C	14.7°C	15.5°C
YD	14.3°C	14.3°C	14.6°C	14.4°C	14.0°C	14.5°C	14.2°C	14.7°C	13.9°C	14.0°C
LA	14.5°C	14.6°C	14.8°C	14.3°C	14.5°C				14.5°C	14.3°C

Table 3. Mean T_{July} estimation for the YD (gray field) from central Poland based on cladoceran-inferred model. EH: Early Holocene; YD: Younger Dryas; LA: Late Allerød

most YD paleotemperatures reconstructed for the YD for central Poland, based on chironomid models that show the YD mean average July air temperature to have been ca. 14°C (Żabieniec, Płóciennik *et al.* 2011; Koźmin, Dzieduszyńska *et al.* 2014). However, at sites like Rąbień and Świerczyna, which are dominated by the chironomid *Corynocera ambigua* (80–90%), the reconstructed summer temperatures are much lower (at ca. 8°C) than at nearby localities. This is because they are closely associated with the local environmental conditions (i.e., high levels of mineral matter, flooding, and weakly developed plant communities) that affected the chironomid assemblages (Pawłowski *et al.* 2015a).


The cladoceran-inferred summer temperature reconstruction from paleo-oxbow lakes in central Poland seems to be reliable and comparable to the record from plateau sites that are not exposed to the direct activity of rivers. There is insufficient radiocarbon data from all the study sites to allow plausible discussion of the correlation of annual or near-annual resolution records; however, it can be suggested that most study sites reflect fluctuating conditions during the YD. This period can thus be divided into a first phase, associated with lower temperature (to 12,000 cal yrs BP), and a second warmer YD phase (ca. 12,000–11,500 cal yrs BP). These observations confirm previous investigations from the region, which show similar patterns of temperature and precipitation (Płóciennik *et al.* 2011; Pawłowski *et al.* 2015a). Additionally, this trend is consistent with the bipartite climate division of the period (a cold first part followed by a warmer second part) detected by Peyron *et al.* (2005) in records from central and northwestern Europe. This situation also has been linked with changes in atmospheric circulation regimes (Bakke *et al.* 2009; Lane *et al.* 2013).

Between 12,500 and 12,300 cal yrs BP, the pollen-based reconstructions from central Polish river valleys are characterized by decreases in winter temperatures from -11 to -22°C (Pawłowski *et al.* 2015a).

These estimates are very close to the results of the climate stimulations of Isarin and Bohncke (1999), who reported a decrease in the YD winter temperatures to -25°C in northwestern and central Europe. This confirms the general trend toward cold and dry conditions over central and eastern Europe during the YD, driven by enhanced continentality. The decline in temperatures in the YD was found to be more pronounced in winter than in summer (Feurdean *et al.* 2014); the winter temperature estimates seem to be similar throughout Europe (Pawłowski *et al.* 2015a).

The cladoceran-based summer temperature reconstructions from central Poland (12.2–16.5°C) are similar to the cladoceran-based estimates for eastern Poland (11.5–14.5°C; Łukie Lake; Zawiska *et al.* 2014) and to the chironomid-inferred summer temperature estimates for the mid-latitudes (50–55°N) of east and central-southern Europe (Heiri *et al.* 2014). Additionally, the YD temperature values from central Poland studied here are very similar to the estimates from lowlands and mid-elevation (up to 700 m a.s.l.), central European and southern European lakes (Table 4), whose temperatures ranged between 12°C and 16°C (Stergoju, Romania; Feurdean *et al.* 2008), 13.5°C and 16°C (Kraków-Wieluń Upland, Poland; Lorenc 2008), 12.1°C and 14.4°C (15.7–17.3°C, Żabieniec, Poland; Płóciennik *et al.* 2011), 14.8°C and 15.8°C (Koźmin, Poland; Dzieduszyńska *et al.* 2014), 13.5°C and 14.5°C (Hijkermeer, Netherlands; Heiri *et al.* 2007), 13°C and 16°C (Friedländer Große Wiese, Germany; van Asch *et al.* 2012), 14°C and 15°C (Lautrey, France; Heiri and Millet 2005), 15°C and 15.5°C (Ech, France; Millet *et al.* 2012), and ca. 16°C (Lago di Piccolo, Italy; Larocque and Finsinger 2008). In turn, the summer estimates in northern and eastern Europe were significantly lower (Table 4): 8–10°C (England; Lang *et al.* 2010); 7.5–10°C (Lough Nadourcan, Ireland; Watson *et al.* 2010); 10–12°C (Sergeevskoe, Belarus; Veski *et al.* 2015); 11–14°C (Kurjanovas, Latvia; Veski *et al.* 2015); and 9.5–13°C (Nakri, Estonia; Veski *et al.* 2015).

RIVER VALLEYS



UPLAND HILLS


Lithology

- 1 – fluvial sands
- 2 – peat with sands
- 3 – peat
- 4 – gyttja
- 5 – gyttja with sands
- 6 – detritus-clay gyttja
- 7 – clay gyttja
- 8 – sedge peat with sand
- 9 – sedge-moss peat with sands
- 10 – detrital-calcareous gyttja
- 11 – detrital gyttja
- 12 – reed peat
- 13 – organic mud
- 14 – sandy silt
- 15 – reed-sedge peat


Chronostratigraphy

- LA – Late Allerød
- YD – Younger Dryas
- EH – Early Holocene

Żabieniec



Rąbień



Text-fig. 5. Summary diagram demonstrating the cladoceran-inferred summer temperature reconstructions from central Poland. Records are plotted versus the age-depth model. Black lines: cladoceran-inferred models; red lines: smoothed trend of reconstruction; error bar estimates are shown as black horizontal spans


	Central and southern Europe								
	Koźmin ** Poland	Żabieniec ** Poland Norwegian TS Swiss TS	Krakowsko- Wieluńska Upland *** Poland	Łukie Lake * Poland	Hijkermeer ** Netherlands	Friedländer Grobe Wiese ** Germany	Ech ** France	Lautrey ** France	Lago di Piccolo ** Italy
YD	14.8–15.8°C	12.1–14.4°C 15.7–17.3°C	13.5–16°C	11.5–14.5°C	13.5–14.5°C	13–16°C	15–15.5°C	14–15°C	16°C
	Central and southern Europe				Northern and eastern Europe				
	Gościąg *** Poland	Witów *** Poland	Stereoiu *** Romania	5 sites in England **	Lough Nadourcan ** Ireland	Sergeevskoe *** Belarus	Kurjanovas ** / *** Latvia	Nakri ** / *** Estonia	
YD	10–14°C	12°C	12–16°C	8–10°C	7.5–10°C	10–12°C	11–14°C 9–10.5°C	9.5–13°C 9.5–11.5°C	

Table 4. Temperature estimation for the YD (gray field) from Europe (without the Alpine and Norway regions) based on the cladoceran-inferred model (*), the chironomid-inferred model (**), and other indicators: pollen-inferred models, selected plant indicators, bird bones (***). References to temperature estimates sources are cited in the text

Similarly, Isarin and Bohncke (1999) suggested summer temperatures of about 10–14°C in north and central Europe during the YD. In turn, Ralska-Jasiewiczowa *et al.* (1998) suggest 10°C for Poland in the first part of the YD and about 12–14°C in the second part (Wasylikowa 1964) (Table 4). However, this estimate was based on the thermal requirements of selected climate indicator plant species, and should thus be treated with caution when compared with other results or reconstructions (Pawłowski *et al.* 2015a). Overall, the central and southern Europe climate trend during the YD differs from those in the northern and eastern part of Europe. This thermal offset is probably due to latitudinal shifts and the restructuring in the oceanic surface current circulation over the North Atlantic area (Gandouin *et al.* 2016). The estimates from central Poland, which are slightly higher than the reconstructions for northern and eastern Europe, could also be explained by the influence of more continental than oceanic air masses affecting the weather conditions in the eastern and northern parts of Europe. Further, the enhanced continentality could have resulted in a temporary increase in summer temperatures and lower winter temperatures, resulting in a more rapid reaction of aquatic invertebrates to this event, but a complete lack of reaction from vegetation. According to the results from central Poland, the cladoceran-based model infers a rise in the YD summer temperature slightly earlier than does the pollen-based model (Pawłowski *et al.* 2015a). It is probable that this is related to the high sensitivity of cladocerans

to changing conditions and to their short life cycle, which allows them to respond more quickly to climate change. Finally, it is possible that discrepancies have arisen from the different methods of estimation used in cladoceran, chironomid, and pollen/plant reconstructions; a model which is constructed to infer summer mean temperatures reflects a true mean and not individual measurements; otherwise, chance factors may strongly influence the resulting model. Additionally, the new findings of temperate-adapted species, such as *Quercus* and *Fagus* from central Europe in the YD (Robin *et al.* 2016), may suggest higher temperature estimates than the reconstructions known to us at present. The Cladocera-based temperature scenario thus seems to be realistic and appears to be the most plausible.

During ca. 12,000–11,500 cal yrs BP, a gradual warming occurred (Text-fig. 5). This warming likely resulted in strongly contrasting seasonal hydrological conditions: summer temperatures increased (13.5–16°C), corresponding to a gradual rise in winter temperatures and in annual precipitation (which was, however, variable in spring and summer) in the Grabia River valley (Pawłowski *et al.* 2015a). A similar shift in the mid-YD as a climatic amelioration associated with changes driven by the large-scale reorganizations of atmospheric and oceanic circulation patterns has been noted in Kråkenes (Norway; Bakke *et al.* 2009) and Meerfelder Maar (Germany; Lane *et al.* 2013). Although these atmospheric reorganizations, which resulted in rapid alternations between glacial growth and melting during the later YD, were



Text-fig. 6. Modern analogue analysis (MAT; red line) and goodness-of-fit (GoF; blue line) of the fossil assemblages to temperature. The vertical red dotted line indicates 10% of the squared chord distances of the fossil sample to the samples in the modern calibration dataset. The vertical blue dotted line indicates 10% of the squared residual distances of the modern samples to the first axis in a CCA. The samples to the right of the lines have no good modern analogue and have respectively a poor or very poor fit-to-temperature. The units for GoF are 'squared residual distance'; for MAT, 'squared chord distance'

expressed most clearly on the margins of the polar and Atlantic air masses, it is possible that these climate shifts could also have influenced periodicity in central Poland during the YD. In addition, this situation confirms that the mid-YD transition was locally abrupt but time-transgressive across Europe. A similar warming trend at the end of the YD has been

widely observed in other lakes throughout Europe (Bohncke *et al.* 1993; Goslar *et al.* 1993; Ralska-Jasiewiczowa *et al.* 1998; Lotter *et al.* 2000; Velichko *et al.* 2002; Birks and Ammann 2000; Birks *et al.* 2010; Neugebauer *et al.* 2012; Brooks and Langdon 2014; Ganduoin *et al.* 2016). Most authors have suggested a probable strong warming (ca. +2–3°C) in the second

and terminal parts of the YD. The sites investigated here are located within a transition zone between the influences of the North Atlantic and the continental climatic regimes, resulting in variable weather conditions. Some of the results thus reflect the climate dynamics of western, northwestern, and southern European sites (Magny *et al.* 2006; Neugebauer *et al.* 2012; Hošek *et al.* 2014). These results suggest that Atlantic air masses may have penetrated to the east during the second part of the YD and led to oceanic influence on terrestrial ecosystems, causing changes in seasonality (i.e., relatively warm and wet winters, and an increase in spring precipitation).

The beginning of the Holocene at the study sites was characterized by an increase in summer temperatures and precipitation, as well as by prolonged growing seasons. This could indicate a decrease in the influence of the continental climate and may imply a temporary climatic connection of the study area with the circum-North Atlantic air masses (Pawłowski *et al.* 2015a).

Reliability of the cladoceran-based temperature reconstruction

The GoF and MAT analyses suggest that temperature may be one of the major drivers of change in the cladoceran assemblages. However, most subfossil samples had no 'close' analogue in the modern calibration data set (Text-fig. 6). These lack-of-fit measures indicate that the fossil cladoceran assemblages in those samples may be responding to changes in an environmental variable other than temperature. Temperature reconstructions from samples where goodness-of-fit values are particularly extreme may thus be less reliable; they should be considered tentative and interpreted with caution. It is however possible that, in samples lacking a close modern analogue in the training set, the dominant taxa in the fossil sample are less well-represented in the modern training set. Rather than suggesting that temperature is not a strong driver of these samples, it may be that the temperature optima of these taxa has not been well-estimated; the temperature estimate of these samples would therefore be less reliable. For this reason, it is necessary to exclude drivers associated with local changes or climate conditions. The local factors (e.g., water level changes and influence of the rivers, habitat modification, macrophyte abundance, eutrophication, acidity, predation of fish, and presence of CaCO₃) may also be crucial in determining cladoceran assemblages, and therefore the Cladocera-based temperature estimation could be dubious.

The greatest differences in the estimated YD temperature occurs between the sites located in the river valleys – Ldzań (13.9°C), Grabica, and Koźmin (at approximately 14.7°C). There are a few possible explanations. The first may be related to the geomorphological position of the sites, changes in the riverbeds, or large individual floods. An increase in fluvial activity during the YD has been widely reported from Polish and European valleys (e.g. Huisink 2000; Andres *et al.* 2001; Starkel 2002, 2011; Borisova *et al.* 2006; Gao *et al.* 2007; Starkel *et al.* 2007, 2015; Pawłowski *et al.* 2015b; Petera-Zganiacz *et al.* 2015). The floodwaters may significantly decrease the water temperature in the oxbow lake, as at the Ldzań site. In contrast to other sites investigated here, which are located much further from active river channels (from 600 m to 2000 m), Ldzań is situated very close (100 m) to the present-day Grabia River channel (Pawłowski *et al.* 2014). The impact of the river could thus have been more significant on the water temperature of this oxbow lake than at the other sites. The second possible explanation is associated with the density and composition of vegetation, which affects the duration of snow cover and seasonal snowmelt. In the YD, the Koźmin site was a very shallow periodic flood basin with the riparian forest, which persisted through the YD under potentially good growing conditions in the floodplain of the Warta River, despite the generally open landscape of central Poland (Dzieduszyńska *et al.* 2014). It is therefore possible that places such as Koźmin, which offer vegetation associations some protection and refuge from climate extremes, could affect the T_{July} estimations, which could be slightly higher than in other paleo-oxbow lakes in floodplains. The peak in cladoceran density could also be associated with periods when the water was relatively warm; an increase in water temperature generally results in rich macrovegetation covering the littoral zone, allowing an increase in the number of macrophyte-associated species. This situation does not support the idea that the slightly higher frequency of Cladocera at sites on plateaus or isolated from fluvial influences (Żabieniec and Rąbień) constitutes the only reason for higher temperature estimates; similar T_{July} estimations are, after all, present at paleo-oxbow lakes. The third possible explanation for the higher T_{July} estimate at Koźmin and Grabica is connected with fact that these sites were very shallow pools during the YD. Water level changes seem to be a very important factor affecting the cladoceran community (Pawłowski *et al.* 2016a). In extremely shallow water, the inferred values and residuals in the

Cladocera-based calibration set most likely cause an overestimation of temperature. In such water pools, Cladocera are represented mainly by taxa such as *A. affinis*, *Ch. sphaericus*, *Al. excisa*, and *A. guttata* var. *tuberculata*. These benthic taxa are abundant in temperate lakes and mires, and their T_{July} optima are similar (ca. 14.5–15.2°C). Their presence in such shallow sites could lead to overestimations in temperature, as their high frequency is presumably related to nonclimatic limnological factors (such as macrophyte density and low pH); however, these changes may also result from climate-related variables (such as T_{July}). Similar situations have been observed in the Cladocera-inferred water depth models (Luoto *et al.* 2011; Nevalainen 2011; Pawłowski *et al.* 2016a) as “edge effects”. Such effects suggest that the Cladocera-based inference model can most reliably quantify past water level changes at intermediate depths relative to the training set depth gradient (because they lead to overestimation at shallow sites and an underestimation at deeper sites). It is possible that this could be applied to temperature estimations. However, such “edge effects” in the nonlinear distortions at the ends of the gradients constitute a problem inherent to all unimodal-based calibration methods (Lotter *et al.* 1997).

It is very interesting that the presence of zooplanktivorous fish seems not to be the main influence on the cladoceran-based temperature model. Fish generally affect zooplankton communities and have major cascading effects on food webs (Jeppesen *et al.* 2001; Davidson *et al.* 2007). Although fish predator species are known only at Pawłowa (Pawłowski *et al.* 2016b), it cannot be excluded that there was at least temporary fish predation at other sites. Size-selective predation on larger cladoceran species could increase the relative abundance of some planktonic species. For example, the rise of *Bosmina longirostris* in concert with the density of zooplanktivorous fish could also result in an absence of *Daphnia pulex* (Davidson *et al.* 2010). However, changes associated with size-selective predation of larger species (e.g., *Daphnia* spp.) and increases in small-bodied Bosminidae, as described at Pawłowa (Pawłowski *et al.* 2016b), are likely to have affected the T_{July} reconstruction only to a small extent, because these planktonic taxa have similar T_{July} optima of around 14.9–15.9°C. These remarks also apply to a number of larger benthic chydorid taxa, in particular *A. harpae*, *E. lamellatus*, *Pleuroxus* spp., and *Ch. sphaericus*; a slight increase in zooplanktivorous fish density may have resulted in the absence of these taxa. These

cladocerans are abundant in temperate lakes and mires and are strongly associated with the presence of aquatic macrophytes at lower fish densities (Davidson *et al.* 2010). However, their T_{July} optima are similar to those of small-bodied Bosminidae, ca. 14.5–15.7°C. Similarly, there is no link between low *Daphnia* incidences and the very low calcium content in the water of the studied lakes (Jeziorski *et al.* 2008) in the YD, which could have influenced the Cladocera-based temperature estimation.

The high CaCO_3 content at the Ługi and Pawłowa sites (the presence of detrital-calcareous gyttja) is also unlikely to have affected the T_{July} reconstruction in the YD (Text-fig. 5). The CaCO_3 enrichment of sediments at Pawłowa can be attributed to bio-chemical calcite precipitation due to the activity of autotrophic organisms in the lake (e.g., *Chara*), inputs of detrital carbonates, and leaching of carbonate-rich raw soils in the lake’s catchment (Pawłowski *et al.* 2016b). In contrast to Pawłowa, there is no insight into the direct mechanism involved in calcium carbonate precipitation in Ługi. It is thus possible, though less plausible, that these changes refer to a change in the temperatures in spring and summer when most autochthonous carbonates are precipitated within the lake; however, in the sites studied, this problem applies only to the Allerød period (Text-fig. 5).

CONCLUSIONS

This study concludes that the distribution of cladoceran species from different lakes located on active river valley beds, in fossil valleys, and on plateaus of glacial and aeolian origin is mainly associated with changes in temperature. Although other environmental changes in physical, chemical, and biological lake dynamics could have affected the cladoceran assemblages, temperature remains the most important control variable in the system of interest. These aquatic organisms thus have potential as quantitative climate indicators in the YD. The cladoceran-inferred summer temperature estimates show two phases during the YD: a first colder phase between 12,800 and 12,000 cal yrs BP, and a second warmer phase between ca. 12,000 and 11,500 cal yrs BP. In the YD, the cladoceran-based summer temperature reconstructions in central Poland show a decreasing trend and had an average estimate of ca. 14°C. The lowermost estimates, at ca. 12°C, are between 12,500 and 12,300 cal yrs BP. In turn, the highest estimates, at ca. 15°C, are at the end of the YD.

Acknowledgments

I am grateful to the reviewers for their critical and helpful comments. I thank the National Science Centre (NCN) for funding the study entitled “The role of valley mires in reconstructing paleohydrological events in the light of selected paleoecological studies” (no 2011/01/B/ST10/04905). I also thank, and am indebted to, Tomi Luoto of the Department of Biological and Environmental Science, University of Jyväskylä, Finland, for his help in processing the Cladocera-based mean July air temperature transfer function. Special thanks also go to Steve Brooks from the Department of Life Sciences, Natural History Museum, London, UK, for his help in processing the reliability of the temperature reconstructions (GoF and MAT) and for several important suggestions.

REFERENCES

- Andres, W., Bos, J.A.A., Houben, P., Kalis, A.J., Nolte, S., Rittweger, H. and Wunderlich, J. 2001. Environmental change and fluvial activity during the Younger Dryas in central Germany. *Quaternary International*, **79**, 89–100.
- Bakke, J., Lie, Ø., Heegaard, E., Dokken, T., Haug, G.H., Birks, H.H., Dulski, P. and Nilsen, T. 2009. Rapid oceanic and atmospheric changes during the Younger Dryas cold period. *Nature Geoscience*, **2**, 202–205.
- Birks, H.H. and Birks, H.J.B. 1980. Quaternary palaeoecology, pp. 95–120. The Blackburn Press; New York.
- Birks, H.H. and Ammann, B. 2000. Two terrestrial records of rapid climatic change during the glacial–Holocene transition (14,000–9,000 calendar years B.P.) from Europe. *Proceedings of the National Academy of Sciences of USA*, **97**, 1390–1394.
- Birks, H.J.B., Heiri, O., Seppä, H. and Bjune, A.E. 2010. Strengths and weaknesses of quantitative climate reconstructions based on late-Quaternary biological proxies. *The Open Ecology Journal*, **3**, 68–110.
- Bjerring, R., Becares, E., Declerck, S., Gross, E.M., Hansson, L.-A., Kairesalo, T., Nykänen, M., Halkiewicz, A., Kornijów, R., Conde-Porcuna, J.M., Seferlis, M., Noges, T., Moss, B., Amsinck, S.L., Odgaard, B.V. and Jeppesen, E. 2009. Subfossil Cladocera in relation to contemporary environmental variables in 54 Pan-European lakes. *Freshwater Biology*, **54**, 2401–2417.
- Bohncke, S., Vandenbergh, J. and Huijzer, A.S. 1993. Periglacial environments during the Weichselian Late Glacial in the Maas valley, the Netherlands. *Geologie en Mijnbouw*, **72**, 193–210.
- Borisova, O., Sidorchuk, A. and Panin, A. 2006. Palaeohydrology of the Seim River basin, Mid-Russian Upland, based on palaeochannel morphology and palynological data. *Catena*, **66**, 53–73.
- Bronk Ramsey, C. 2009. Bayesian analysis of radiocarbon dates. *Radiocarbon*, **51**, 337–360.
- Brooks, S.J. and Langdon, P.G. 2014. Summer temperature gradients in northwest Europe during the Lateglacial to early Holocene transition (15–8 ka BP) inferred from chironomid assemblages. *Quaternary International*, **341**, 80–90.
- Brooks, S.J., Matthews, I.P., Birks, H.H. and Birks, H.J.B. 2012. High resolution late-glacial and early Holocene summer air temperatures from Scotland inferred from chironomid midge assemblages. *Quaternary Science Reviews*, **41**, 67–82.
- Davidson, T.A., Sayer, C.D., Perrow, M.R., Bramm, M. and Jeppesen, E. 2007. Are controls on species composition similar for contemporary and sub-fossil cladoceran assemblages? A study of 39 shallow lakes of contrasting trophic status. *Journal of Paleolimnology*, **38**, 117–134.
- Davidson, T., Sayer, C.D., Perrow, P., Bramm, M. and Jeppesen, E. 2010. The simultaneous inference of zooplanktivorous fish and macrophyte density from sub-fossil cladoceran assemblages: a multivariate regression tree approach. *Freshwater Biology*, **55**, 546–564.
- Dzieduszyńska, D.A., Kittel, P., Petera-Zganiacz, J., Brooks, S.J., Korzeń, K., Krapiec, M., Pawłowski, D., Płaza, D.K., Płóciennik, M., Stachowicz-Rybka, R. and Twardy, J. 2014. Environmental influence on forest development and decline in the Warta River valley (Central Poland) during the Late Weichselian. *Quaternary International*, **324**, 99–114.
- Engels, S., Bohncke, S.J.P., Heiri, O. and Nyman, M. 2008. Intra-regional variability in chironomid-inferred temperature estimates and the influence of river inundations on lacustrine chironomid assemblages. *Journal of Paleolimnology*, **40**, 129–142.
- Forysiak, J. 2012. Zapis zmian środowiska przyrodniczego późnego Wistulianu i holocenu w osadach torfowisk regionu łódzkiego. *Acta Geographica Lodziensia*, **99**, 1–164. [In Polish with English summary]
- Frey, D.G. 1986. Cladocera analysis. In: B.E. Berglund (Ed.), *Handbook of Holocene paleoecology and paleohydrology*, pp. 667–692. John Wiley and Sons; Chichester.
- Feurdean, A., Klotz, S., Brewer, S., Mosbrugger, V., Tămaş, T. and Wohlfarth, B. 2008. Lateglacial climate development in NW Romania – comparative results from three quantitative pollen-based methods. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **265**, 121–133.
- Feurdean, A., Persoiu, A., Tanțău, J., Stevens, T., Magyari, E.K., Onac, B.B., Markovic, S., Andric, M., Connor, S., Galka, M., Hoek, W.Z., Lamentowicz, M., Sümegi, P., Persoiu, J., Kolaczek, P., Kuneš, P., Marinova, E., Slowinski, M., Michczyńska, D., Stancikaitė, M., Svensson, A., Veski, S., Fărcaș, S., Tămaş, T., Zernitskaya, V., Timar, A., Tonkov, S., Toth, M., Willis, K.J., Płóciennik, M. and Gaudeny, T. 2014. Climate variability and associated vegetation response throughout Central and Eastern Europe (CEE) between 60 and 8 ka. *Quaternary Science Reviews*, **106**, 206–224.

- Gandouin, E., Ponel, P., Franquet, E., van Vliet-Lanoë, B., Andrieu-Ponel, V., Keen, D.H., Brulhet, J. and Brocandel M. 2007. Chironomid responses (Insect: Diptera) to Younger Dryas and Holocene environmental changes in a river floodplain from northern France (St-Momelin, St-Omer basin). *Holocene*, **17**, 331–347.
- Gandouin, E., Rioual, P., Pailles, C., Brooks, S.J., Ponel, P., Guiter, F., Djamali, M., Andrieu-Ponel, V., Birks, H.J.B., Leydet, M., Belkacem, D., Haas, J.N., Van der Putten, N. and de Beaulieu, J.L. 2016. Environmental and climate reconstruction of the late-glacial–Holocene transition from a lake sediment sequence in Aubrac, French Massif Central: Chironomid and diatom evidence. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **461**, 292–309.
- Gao, C., Boreham, S., Preece, R.C., Gibbard, P.L. and Briant, R.M. 2007. Fluvial response to rapid climate change during the Devensian (Weichselian) Lateglacial in the River Great Ouse, S England, UK. *Sedimentary Geology*, **202**, 193–210.
- Goslar, T., Kuc, T., Ralska-Jasiewiczowa, M., Rozanski, K., Arnold, M., Bard, E., van Geel, B., Pazdur, M.F., Szeroczyńska, K., Wicik, B., Więckowski, K. and Walanus, A. 1993. High resolution lacustrine record of the Late Glacial/Holocene transition in Central Europe. *Quaternary Science Reviews*, **12**, 295–305.
- Goslar, T., Ralska-Jasiewiczowa, M., Starkel, L., Demske, D., Kuc, T., Łącka, B., Szeroczyńska, K., Wicik, B. and Więckowski, K. 1998. Discussion of the Late-Glacial record in the Lake Gościąg sediments. In: M. Ralska-Jasiewiczowa, T. Goslar, T. Madeyska and L. Starkel (Eds), *Lake Gościąg, central Poland, A monographic study*, pp. 171–180. IB PAN; Kraków.
- Heiri, O. and Millet, L. 2005. Reconstruction of Late Glacial summer temperatures from chironomid assemblages in Lac Lautrey (Jura, France). *Journal of Quaternary Science*, **20**, 1–12.
- Heiri, O., Cremer, H., Engels, S., Hoek, W.Z., Peeters, W. and Lotter, A.F. 2007. Lateglacial summer temperatures in the Northwest European lowlands: a chironomid record from Hijkermeer, the Netherlands. *Quaternary Science Reviews*, **26**, 2420–2437.
- Heiri, O., Brooks, S. J., Renssen, H., Bedford, A., Hazekamp, M., Ilyashuk, B., Jeffers, E.S., Lang, B., Kirilova, E., Kuiper, S., Millet, L., Samartin, S., Toth, M., Verbruggen, F., Watson, J.E., van Asch, N., Lammertsma, E., Amon, L., Birks, H.H., Birks, H.J.B., Mortensen, M.F., Hoek, W.Z., Magyari, E., Sobrino, C.M., Seppä, H., Tinner, W., Tonkov, S., Veski, S. and Lotter, A.F. 2014. Validation of climate model-inferred regional temperature change for late-glacial Europe. *Nature Communication*, **5**, 1–7.
- Hošek, J., Pokorný, P., Kubovčík, V., Horáček, I., Žáčková, P., Kadlec, J., Rojik, F., Lisá, L. and Bučkuliaková, S. 2014. Late glacial climatic and environmental changes in eastern-central Europe: correlation of multiple biotic and abiotic proxies from the Lake Švarcenberk, Czech Republic. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **396**, 155–172.
- Huisink, M. 2000. Changing river styles in response to Weichselian climate changes in the Vecht valley, eastern Netherlands. *Sedimentary Geology*, **133**, 115–134.
- Isarin, R.F.B. and Bohncke, S.J.P. 1999. Mean July temperatures during the Younger Dryas in northwestern and central Europe as inferred from climate indicator plant species. *Quaternary Research*, **51**, 158–173.
- Isarin, R.F.B. and Rensen, H. 1999. Reconstructing and modelling Late Weichselian climates: the Younger Dryas in Europe as a case study. *Earth-Science Reviews*, **48**, 1–38.
- Jeppesen, E., Christoffersen, K., Landkildehus, F., Lauridsen, T., Amsinck, S., Riget, F. and Søndergaard, M. 2001. Fish and crustaceans in northeast Greenland lakes with special emphasis on interactions between Arctic charr (*Salvelinus alpinus*), *Lepidurus arcticus* and benthic chydorids. *Hydrobiologia*, **442**, 329–337.
- Jeziorski, A., Yan, N.D., Paterson, A.M., DeSellas, A.M., Turner, M., Jeffries, D.S., Keller, B., Weeber, R.C., McNicol, D.K., Palmer, M.E., McIver, K., Arsenau, K., Ginn, B.K., Cumming, B.F. and Smol, J.P. 2008. The widespread threat of calcium decline in fresh waters. *Science*, **322**, 1374–1377.
- Juggins, S. 2007. C2 Version 1.5 User Guide. Software for Ecological and Palaeoecological Data Analysis and Visualisation. Newcastle University, Newcastle upon Tyne, pp. 1–73.
- Kaiser, K., Lorenz, S., Germer, S., Juschus, O., Küster, M., Libra, J., Bens, O. and Hütti, R.F. 2012. Late Quaternary evolution of rivers, lakes and peatlands in northeast Germany reflecting past climates and human impact – an overview. *E&G Quaternary Science Journal*, **61**, 103–132.
- Klatkova, H. and Załoba, M. 1991. Kształtowanie budowy geologicznej i rzeźby południowego obrzeżenia Basenu Uniejowskiego. In: W. Stankowski (Ed.), *Przemiany środowiska geograficznego obszaru Konin-Turek*, pp. 33–44. Wyd. Nauk UAM; Poznań.
- Kłysik, K. 2001. Warunki klimatyczne. In: S. Liszewski (Ed.), *Zarys monografii województwa łódzkiego*, pp. 68–81. Łódzkie Towarzystwo Naukowe; Łódź.
- Korhola, A. and Rautio, M., 2001. Cladocera and other branchiopod crustaceans. In: J.P. Smol, H.J.B. Birks, and W.M. Last (Eds), *Tracking Environmental Change Using Lake Sediments, Zoological Indicators*, 4, pp. 5–41. Kluwer Academic Publishers; Dordrecht.
- Lane, C.S., Brauer, A., Blockley, S.P.E. and Dulski, P. 2013. Volcanic ash reveals time-transgressive abrupt climate change during the Younger Dryas. *Geology*, **41**, 1251–1254.
- Lang, B., Brooks, S.J., Bedford, A., Jones, R.T., Birks, H.J.B. and Marshall, J.D. 2010. Regional consistency in Late-glacial chironomid-inferred temperatures from five sites in north-west England. *Quaternary Science Reviews*, **29**, 1528–1538.

- Larocque, I. and Finsinger, W. 2008. Late-glacial chironomid-based temperature reconstructions for Lago Piccolo di Avigliana in the southwestern Alps (Italy). *Palaeogeography, Palaeoclimatology, Palaeoecology*, **257**, 207–223.
- Lorenc, M. 2008. Omożliwościach rekonstrukcji paleotemperatur wistulianu na podstawie kopalnych zespołów ptaków. *Geologos*, **14**, 91–106.
- Lotter, A.F., Birks, H.J.B., Hofmann, W. and Marchetto, A. 1997. Modern diatom, cladocera, chironomid, and chrysophyte cyst assemblages as quantitative indicators for the reconstruction of past environmental conditions in the Alps. I. Climate. *Journal of Paleolimnology*, **18**, 395–420.
- Lotter, A.F., Birks, H.J.B., Eicher, U., Hofmann, W., Schwander, J. and Wick, L. 2000. Younger Dryas and Allerød summer temperatures at Gerzensee (Switzerland) inferred from fossil pollen and cladoceran assemblages. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **159**, 349–361.
- Luoto, T.P., Nevalainen, L., Kultti, S. and Sarmaja-Korjonen, K. 2011. An evaluation of the influence of water depth and river inflow on quantitative Cladocera-based temperature and lake level inferences in a shallow boreal lake. *Hydrobiologia*, **676**, 143–154.
- Macklin, M.G., Benito, G., Gregory, K. J., Johnstone, E., Lewin, J., Michczyńska, D.J., Soja, R., Starkel, L. and Thornycroft, V.R. 2006. Past hydrological events reflected in the Holocene fluvial record of Europe. *Catena*, **66**, 145–154.
- Magny, M., Aalbersberg, G., Begeot, C., Benoit-Ruffaldi, P., Bossuet, G., Disnar, J.R., Heiri, O., Laggoun-Defarge, F., Mazier, F., Millet, L., Peyron, O., Vanniere, B. and Walter-Simonnet, A.V. 2006. Environmental and climatic changes in the Jura mountains (eastern France) during the Late glacial–Holocene transition: a multi-proxy record from Lake Lautrey. *Quaternary Science Reviews*, **25**, 414–445.
- Michczyńska, D.J., Borówka, R.K., Okupny, D., Obremska, M., Forsytek, J., Pawłowski, D., Płóciennik, M., Słowiński, M., Żurek, S., Brook, S.J., Michczyński, A. and Witkowski, A., 2014. The environment changes and chronology of the Late Vistulian (Weichselian) and Early Holocene sediments in the Rąbień mire, Central Poland. In: A. Moreno, B.L. Valero-Garcés and S.O. Rasmussen (Eds), INTIMATE: Open Workshop and COST Action ES0907 Final Event, pp. 54–55. Zaragoza
- Milecka, K., Kowalewski, G. and Szeroczyńska, K. 2011. Climate-related changes during the Late Glacial and early Holocene in northern Poland, as derived from the sediments of Lake Sierzywk. *Hydrobiologia*, **676**, 187–202.
- Millet, L., Rius, D., Galop, D., Heiri, O. and Brooks, S.J. 2012. Chironomid-based reconstruction of Lateglacial summer temperatures from the Ech palaeolake record (French western Pyrenees). *Palaeogeography, Palaeoclimatology, Palaeoecology*, **315–316**, 86–99.
- Mirowska-Grabowska, J. and Zawisza, E. 2013. Late Glacial–early Holocene environmental changes in Charzykowskie Lake (northern Poland) based on oxygen and carbon isotopes and Cladocera data. *Quaternary International*, **328–329**, 156–166.
- Mirowska-Grabowska, J., Niska, M. and Kupryjanowicz, M. 2015. Reaction of lake environment on the climatic cooling-Transition from the Eemian Interglacial to Early Vistulian on the basis of Solniki palaeolake sediments (NE Poland). *Quaternary International*, **386**, 158–170.
- Nevalainen, L. 2011. Intra-lake heterogeneity of sedimentary cladoceran (Crustacea) assemblages forced by local hydrology. *Hydrobiologia*, **676**, 9–22.
- Nevalainen, L., Luoto, T.P., Kultti, S. and Sarmaja-Korjonen, K. 2012. Do subfossil Cladocera and chydorid ephippia disentangle Holocene climate trends? *Holocene*, **22**, 291–299.
- Neugebauer, I., Brauer, A., Dräger, N., Dulski, P., Wulf, S., Plessen, B., Mingram, J., Herzschuh, U. and Brande, A. 2012. A Younger Dryas varve chronology from the Rehwiess palaeolake record in NE-Germany. *Quaternary Science Reviews*, **36**, 91–102.
- Notebaert, B. and Verstraeten, G. 2010. Sensitivity of West and Central European river systems to environmental changes during the Holocene: a review. *Earth-Science Reviews*, **103**, 163–182.
- Pawłowski, D. 2011. Evolution of an Eemian lake based on Cladocera analysis (Konin area, Central Poland). *Acta Geologica Polonica*, **61**, 441–450.
- Pawłowski, D. 2012a. Early development of Late Vistulian (Weichselian) lacustrine sediments in the Żabieniec swamp (central Poland). *Geochronometria*, **39**, 197–211.
- Pawłowski, D. 2012b. Younger Dryas Cladocera assemblages from two valley mires in central Poland and their potential significance for climate reconstructions. *Geologos*, **18**, 237–249.
- Pawłowski, D., Kloss, M., Obremska, M., Szymanowski, M. and Żurek, S. 2012. Evolution of small valley mire in central Poland as a result of hydroclimatic oscillations. *Geochronometria*, **39**, 133–148.
- Pawłowski D., Okupny D., Włodarski W. and Zieliński T. 2014. Spatial variability of selected physicochemical parameters within peat deposits in small valley mire: a geostatistical approach. *Geologos*, **20**, 269–288.
- Pawłowski, D., Płóciennik, M., Brooks, S.J., Luoto, T.P., Milecka, K., Nevalainen, L., Peyron, O., Self, A. and Zieliński, T. 2015a. A multiproxy study of Younger Dryas and Early Holocene climatic conditions from the Grabia River palaeo-oxbow lake (central Poland). *Palaeogeography, Palaeoclimatology, Palaeoecology*, **438**, 34–50.
- Pawłowski, D., Kowalewski, G., Milecka, K., Płóciennik, M., Woszczyk, M., Zieliński, T., Okupny, D., Włodarski, W. and Forsytek, J. 2015b. A reconstruction of the palaeohydrological conditions of a floodplain: a multi-proxy study from the Grabia River valley mire, central Poland. *Boreas*, **44**, 543–562.

- Pawłowski, D., Milecka, K., Kittel, P., Woszczyk, M. and Szychalski, W., 2015c. Palaeoecological record of natural changes and human impact in a small river valley in Central Poland. *Quaternary International*, **370**, 12–28.
- Pawłowski, D., Borówka, R.K., Kowalewski, G., Luoto, T.P., Milecka, K., Nevalainen, L., Okupny, D., Płóciennik, M., Woszczyk, M., Tomkowiak, J. and Zieliński, T. 2016a. The response of flood-plain ecosystems to the Late Glacial and Early Holocene hydrological changes: A case study from a small Central European river valley. *Catena*, **147**, 411–428.
- Pawłowski, D., Borówka, R.K., Kowalewski, G., Luoto, T.P., Milecka, K., Nevalainen, L., Okupny, D., Zieliński, T. and Tomkowiak, J. 2016b. Late Weichselian and Holocene record of the paleoenvironmental changes in a small river valley in Central Poland. *Quaternary Science Reviews*, **135**, 24–40.
- Peyron, O., Bégeot, C., Brewer, S., Heiri, O., Magny, M., Millet, L., Ruffaldi, P., Van Campo, E. and Yu, G. 2005. Late-Glacial climatic changes in Eastern France (Lake Lautrey) from pollen, lake-levels, and chironomids. *Quaternary Research*, **64**, 197–211.
- Petera-Zganiacz, J., Dzieduszyńska, D. Twardy, J., Pawłowski, D., Płóciennik, M., Lutyńska, M. and Kittel, P. 2015. Younger Dryas flood events: a case study from the middle Warta River Valley (central Poland). *Quaternary International*, **386**, 55–69.
- Płóciennik, M., Self, A., Birks, H.J.B. and Brooks, S.J. 2011. Chironomidae (Insecta: Diptera) succession in Żabieniec bog and its palaeo-lake (central Poland) through the Late Weichselian and Holocene. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **307**, 150–167.
- Płóciennik, M., Kittel, P., Borówka, R.K., Cywa, K., Forysiak, J., Okupny, D., Obremska, M., Pawłowski, D., Stachowicz-Rybka, R., Szperna, R. and Witkowski, A., 2016. Warunki ekologiczne subkopalnego koryta Kolonia Bechcice na tle hydrologii środkowej doliny Neru. *Acta Geographica Lodziensia*, **105**, 107–124.
- Ralska-Jasiewiczowa, M., Goslar, T., Madeyska, T. and Starkel, L. 1998. Lake Gościąg, Central Poland. A Monographic Study. Part 1. pp. 1–340. W. Szafer Institute of Botany, Polish Academy of Science; Kraków.
- Reimer, P.J., Bard, E., Bayliss, A., Beck, J.W., Blackwell, P.G., Bronk Ramsey, C., Grootes, P.M., Guilderson, T.P., Hafflidsen, H., Hajdas, I., Hatté, C., Heaton, T.J., Hoffmann, D.L., Hogg, A.G., Hughen, K.A., Kaiser, F., Kromer, B., Manning, S.W., Niu, M., Reimer, R.W., Richards, D.A., Scott, E.M., Southon, J.R., Staff, R.A., Turney, C.S.M. and van der Plicht, J. 2013. IntCal13 and Marine13 Radiocarbon Age Calibration Curves 0–50,000 Years cal BP. *Radiocarbon*, **55**, 1869–1887.
- Robin, V., Nadeau, M.-J., Grootes, P.M., Bork, H.-R. and Nelle, O. 2016. Too early and too northerly: evidence of temperate trees in northern Central Europe during the Younger Dryas. *New Phytologist*, **212**, 259–68.
- Starkel, L. (Ed.) 1990. Evolution of the Vistula River valley during the last 15000 years. *Geographical Studies*, **3** (Special Issue 5), 1–220.
- Starkel, L. 2002. Younger Dryas–Preboreal transition documented in the fluvial environment of Polish rivers. *Global and Planetary Change*, **35**, 157–167.
- Starkel, L. 2011. Present-day events and the evaluation of Holocene palaeoclimatic proxy data. *Quaternary International*, **229**, 2–7.
- Starkel, L., Gębica, P. and Superson, J. 2007. Last Glacial–Interglacial cycle in the evolution of river valleys in southern and central Poland. *Quaternary Science Reviews*, **26**, 2924–2936.
- Starkel, L., Michczyńska, D., Gębica, P., Kiss, T., Panin, A. and Perşoiu, I. 2015. Climatic fluctuations reflected in the evolution of fluvial systems of Central-Eastern Europe (60–8 ka cal BP). *Quaternary International*, **388**, 97–118.
- Stivrins, N., Soinenen, J., Amon, L., Fontana, S.L., Gryguc, G., Heikkilä, M., Heiri, O., Kisielién, D., Reitalu, T., Stančikaitė, M., Veski, S. and Seppä, H. 2016. Biotic turnover rates during the Pleistocene–Holocene transition. *Quaternary Science Reviews*, **151**, 100–110.
- Szeroczyńska, K., 1985. Cladocera jako wskaźnik ekologiczny w późnoczwartorzędowych osadach jeziornych Polski Północnej. *Acta Palaeontologica Polonica*, **30**, 3–69. [In Polish with English summary]
- Szeroczyńska, K. and Sarmaja-Korjonen, K. 2007. Atlas of subfossil Cladocera from central and northern Europe. Friends of Lower Vistula Society; Świecie.
- Ter Braak, C.J.F. and Šmilauer, P. 2002. CANOCO Reference Manual and Cano Draw for Windows User's Guide: Software for Canonical Community Ordination (Version 4.5). Microcomputer Power, Ithaca, NY, USA.
- Turner, F., Tolksdorf, J.F., Viehberg, F., Schwalb, A., Kaiser, K., Bittmann, F., van Bramann, U., Pott, R., Staesche, U., Breest, K. and Veil, S. 2013. Lateglacial/early Holocene fluvial reactions of the Jeetzel river (Elbe valley, northern Germany) to abrupt climatic and environmental changes. *Quaternary Science Reviews*, **60**, 91–109.
- van Asch, N., Kloos, M.E., Heiri, O., De Klerk, P. and Hoek, W.Z. 2012. The Younger Dryas cooling in northeast Germany: summer temperature and environmental changes in the Friedländer Große Wiese region. *Journal of Quaternary Science*, **27**, 531–543.
- Velichko, A.A., Catto, N., Drenova, A.N., Klimanov, V.A., Kremenetski, K.V. and Nechaev, V.P. 2002. Climate changes in East Europe and Siberia at the late glacial–Holocene transition. *Quaternary International*, **91**, 75–99.
- Veski, S., Seppä, H., Stančikaitė, M., Zernitskaya, V., Reitalu, T., Gryguc, G., Heinsalu, A., Stivrins, N., Amon, L., Vassiljev, J. and Heiri, O. 2015. Quantitative summer and winter

- temperature reconstructions from pollen and chironomid data between 15 and 8 ka BP in the Baltic-Belarus area. *Quaternary International*, **388**, 4–11.
- Walanus, A. and Nalepka, D. 1999. Polpal program for counting pollen grains, diagrams plotting and numerical analysis. *Acta Palaeobotanica Suppl.*, **2**, 659–661.
- Wasylikowa, K. 1964. Roślinność i klimat późnego glacjału w środkowej Polsce na podstawie badań w Witowie koło Łęczycy. *Biuletyn Peryglacjalny*, **13**, 261–417.
- Watson, J.E., Brooks, S.J., Whitehouse, N.J., Reimer, P.J., Birks, H.J.B. and Turney, C. 2010. Chironomid-inferred late-glacial summer air temperatures from Lough Nadourcan, Co. Donegal, Ireland. *Journal of Quaternary Science*, **25**, 1200–1210.
- Woś, A. 1999. *Klimat Polski*. Wydawnictwo Naukowe PWN; Warszawa.
- Zawiska, I., Słowiński, M., Correa-Metrio, A., Obremska, M., Luoto, T., Nevalainen, L., Woszczyk, M. and Milecka, K. 2014. The response of a shallow lake and its catchment to Late Glacial climate changes – A case study from eastern Poland. *Catena*, **126**, 1–10.
- Żurek, S. 1987. Złoza torfowe Polski na tle stref torfowych Europy. *Dokumentacja Geograficzna IGiPZ PAN*, **4**, 1–84.

Manuscript submitted: 15th December 2016

Revised version accepted: 15th November 2017