

Matúš PEŠKO¹, Katarína KRÁĽOVÁ¹ and Elena MASAROVICHOVÁ¹

PHYTOTOXIC EFFECT OF SOME METAL IONS ON SELECTED RAPESEED CULTIVARS REGISTERED IN SLOVAKIA

WPLYW FITOTOKSYCZNOŚCI JONÓW NIEKTÓRYCH METALI NA WYBRANE ODMIANY UPRAWNE RZEPAKU ZAREJESTROWANE NA SŁOWACJI

Abstract: The aim of this study was to investigate the phytotoxic effects of seven metal ions (Cd(II), Cr(VI), Cu(II), Hg(II), Ni(II), Pb(II) and Zn(II)) on length of roots of five rapeseed (*Brassica napus* L. *subsp. napus*) cultivars registered in Slovakia (Atlantic, Baldur, Californium, Oponent and Verona). The phytotoxic effect of metals was evaluated using IC₅₀ values. The studied metal ions inhibited germination and root growth of rapeseed seedlings. In general, the toxicity of metal ions decreased in the following order Cu > Cr > Hg > Cd > Pb > Ni > Zn. Atlantic, Baldur and Californium were more sensitive to Cd than to Ni, for Oponent and Verona higher toxicity exhibited Ni. From the studied rapeseed cultivars Atlantic and Californium were found to be most sensitive to tested metals. On the other hand, high tolerance to metal treatment was determined for Baldur. Czech cultivar Oponent showed high tolerance to Cd, Cr, Cu and Pb, but it was sensitive to Hg and Ni. The above-mentioned results confirmed differences in the metal tolerance of tested rapeseed cultivars.

Keywords: toxic metals, *Brassica napus* L., phytotoxicity, root growth

Rapeseed (*Brassica napus* L. *subsp. napus*) is the most important oil crop in Slovakia with a wide spectrum of utilization: as an agricultural crop (food industry), worthwhile nutritive animal fodder (rapeseed press cakes) and technical crop. From rapeseed oil biocomponent FAME for biodiesel is prepared [1, 2]. Some *Brassica* species (*Brassica napus* and *Raphanus sativus*) showed a moderate tolerance against several metals and thus, rapeseed can be used for remediation of soils contaminated by toxic metals [3, 4]. If soils contaminated with heavy metals are phytoremediated by oil crops (such as *Brassica* spp.), biodiesel production from the resulting plant oil could be a viable option to generate bioenergy [5].

Elongation growth is a complex process including turgor changes, synthesis of cell membrane components and content of growth regulators. Excess of heavy metals inhibits elongation plant growth [6, 7]. Some authors consider that inhibition of the cell cycle is the basis for growth inhibition [8]. Moreover, many toxic metal ions are efficient catalysts in the formation of several reactive oxygen species and other free radicals which adversely affect physiological processes in plants, including plant growth [6, 9].

Uptake of toxic amounts of metal by higher plants or algae can result in inhibition of several enzymes and in increase in activity of others. Metal accumulation in the cellular compartment is a prerequisite for enzyme inhibition *in vivo*. Binding of toxic metals to sulfhydryl groups, involved in the catalytic action of structural integrity of enzymes results in inhibition of enzyme activity. On the other hand, the induction of some enzymes (eg

¹ Faculty of Natural Sciences, Comenius University Bratislava, Mlynská dolina CH-2, SK-842 15 Bratislava, Slovakia, phone +421 2 60296340, fax +421 2 65429064, email: pesko@fns.uniba.sk

SOD, CAT, etc.) is considered to play a significant role in the stress metabolism, induced by metal phytotoxicity [10].

This study is aimed to investigate the effect of seven metals ions (Cd(II), Cr(VI), Cu(II), Hg(II), Ni(II), Pb(II) and Zn(II)) on root growth of five cultivars of *Brassica napus* plants registered in Slovakia (Atlantic, Baldur, Californium, Oponent and Verona) and to determine differences in intraspecific sensitivity of rapeseed to metal induced stress.

Material and methods

For experiments following compounds were used: $\text{Cd}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$, $\text{K}_2\text{Cr}_2\text{O}_7$, $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$, HgCl_2 , $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$, $\text{Pb}(\text{NO}_3)_2$ and $\text{ZnCl}_2 \cdot \text{H}_2\text{O}$. These analytical reagent-grade chemicals purchased from Lachema (Brno, Czech Republic) were employed for the preparation of all solutions.

For experiments seeds of following five rapeseed cultivars registered in Slovakia were used: Atlantic, Baldur, Californium, Oponent and Verona. The seeds were purchased from Slovak Centrum of Agricultural Production, Research Institute of Plant Production in Piestany, Slovakia. Atlantic is a medium-early and high type genotype suitable for potato production region; Baldur is a medium-early and medium-high type of genotype suitable for all production regions; Californium is a medium-early to early and medium-high type of genotype suitable for maize production region; Oponent is a late and high type of genotype suitable for potato production region and Verona is a late and medium-high genotype suitable for maize and potato production regions [2, 11].

The seeds of *Brassica napus* were placed in Petri dishes with a 14 cm diameter and filter paper on the bottom. In each Petri dish 58 seeds were evenly displayed on the surface of filter paper and the amount of solution used was 10 cm^3 per dish. Each concentration was duplicated. After 72 hours exposure at mean air temperature (25°C) in the dark the length of roots and shoots was measured. The applied concentration range of studied compounds was 10^{-6} to $10^{-3} \text{ mol dm}^{-3}$.

Results and discussion

Previously it was found that rapeseed is a suitable plant species for investigations of metal phytotoxicity [12]. In the present study *root tolerance index* (RTI) was calculated as the mean seminal root length in the metal treatment divided by the mean seminal root length in the control. From the dependence of RTI on the concentration of applied toxic metals IC_{50} values, ie molar concentrations of the studied metals causing a 50 % inhibition of the studied parameter, were determined (Table 1).

In general, the toxicity of metal ions decreased in the following order $\text{Cu} > \text{Cr} > \text{Hg} > \text{Cd} > \text{Pb} > \text{Ni} > \text{Zn}$. Atlantic (A), Baldur (B) and Californium (C) were more sensitive to Cd than to Ni, for Oponent (O) and Verona (V) higher toxicity exhibited Ni. The sensitivity of studied cultivars treated with toxic metals decreased as follows: for Cd: $A > C > V > B > O$; for Cr: $A = C = V > B = O$; for Pb: $A > V > C > B > O$, for Zn: $A > C > O > V > B$, for Cu: $A > C = V > B > O$, for Hg: $O > C > A = V > B$ and for Ni: $O = A > V > B > C$. Thus, it can be concluded that from the studied rapeseed cultivars Atlantic and Californium were found to be most sensitive to tested metals. On the other hand, overall high tolerance to metal treatment was determined for Baldur. Czech cultivar Oponent showed high tolerance to Cd, Cr, Pb and Cu but it was sensitive to Hg and Ni.

Table 1
 IC₅₀ values related to root growth inhibition of five rapeseed cultivars by tested metals (C.L. - confidence limits)

Metal	IC ₅₀ ± C.L. _{0.05} [mmol dm ⁻³]				
	Atlantic	Baldur	Californium	Oponent	Verona
Cd	0.554 0.295÷1.090	0.659 0.320÷1.47	0.571 0.200÷1.912	0.684 0.427÷1.180	0.615 0.353÷1.150
Cr	0.114 0.056÷0.264	0.122 0.048÷0.400	0.114 0.060÷0.252	0.124 0.060÷0.308	0.128 0.054÷0.324
Cu	0.065 0.037÷0.125	0.088 0.061÷0.145	0.071 0.039÷0.146	0.089 0.047÷0.208	0.072 0.036÷0.192
Hg	0.392 0.222÷0.742	0.539 0.230÷1.542	0.375 0.175÷0.826	0.369 0.237÷0.588	0.396 0.192÷0.878
Ni	0.609 0.443÷1.078	0.644 0.379÷1.142	0.748 0.488÷1.204	0.607 0.428÷0.868	0.612 0.359÷1.084
Pb	0.581 0.299÷1.261	0.736 0.376÷1.575	0.701 0.416÷1.240	0.880 0.644÷1.241	0.696 0.351÷1.480
Zn	0.803 0.404÷1.863	1.168 0.808÷1.757	0.834 0.410÷1.905	0.850 0.576÷1.347	1.130 0.596÷2.287

Toxic effect of metal ions depends on their physico-chemical properties. Nieboer and Richardson [13] classified metals based on their ionic and covalent bonding tendencies and donor-atom preference of metals. These authors used the dependence of the covalent index *versus* the ionic index as a base for classification of metal and metalloid ions to three classes. Metals of class A are oxygen donor-atom seekers, whereas those of class B are nitrogen and sulphur seekers and metals ranged to borderline metals are characterized by ambivalent affinity for all three donor-atom. According to Nieboer et al [14] from the investigated seven metals Hg²⁺ was classified as a metal of class B, the other ones as borderline (intermediate) metals. On the other hand, Duffus [15] assigned Cd²⁺ and Pb²⁺ to the metals of class B.

Ivanov et al [16] investigated the effect of different metal ions on root growth inhibition of maize plants and found that the inhibitory effectiveness decreased in the following order: Cu²⁺ > Hg²⁺ > Cd²⁺ > Zn²⁺ and the IC₅₀ values related to the inhibition of root growth showed correlation with the affinity of metal ions to -SH groups. The rank of inhibitory effectiveness for phytotoxic effects of above metals on maize roots was the same as determined in our experiment for rapeseed plants. It is known that toxic metals act as mitotic inhibitors what is reflected in the reduction of root growth. Jiang et al [17] observed toxic effect of CuSO₄ (10⁻⁴÷10⁻² mol dm⁻³) on the morphology of chromosomes. Doncheva [18] found that copper interrupts the nuclei formation at the crucial- G₁/S transition point of the cell cycle, when it prevents their entry into mitosis. Reduced root growth could be connected with copper effect on proliferation of root meristem cells. Yadav and Srivastava [19] confirmed inhibitory effect of Cd²⁺ ions on mitotic index as well as on active mitotic index for *Hordeum vulgare* and *Setaria italica*. These authors observed different types of mitotic aberration due to Cd²⁺ action. Ni²⁺ ions belong to strong mitotic inhibitors causing modification on membrane permeability, inhibition of ascorbate peroxidase. However, these negative effects could be reduced by synthetic cytokinins [20].

Summarizing the obtained results it can be concluded that root tolerance index (RTI) was found to serve as good biomarker for evaluating the relative toxicity of toxic metals to rapeseed cultivars.

Acknowledgements

This study was financially supported by Development fund of Comenius University in Bratislava under the contract No. UK/209/2011 and Sanofi Aventis Pharma Slovakia.

References

- [1] Masarovičová E., Kráľová K., Brestič M. and Olšovská K.: *Production potential of rapeseed in Slovakia environmental conditions from the aspect of „FAME“ production*. Int. Symp. Motor fuels 2008. Tatranské Matliare, 23.06-26.06.2008, 520-536 (in Slovak).
- [2] Masarovičová E., Kráľová K., Malovcová Ľ., Surovčík J., Brestič M. and Olšovská K.: *Genotypic differences of rapeseed from the aspect of oil content, oil quality and seed production*. Šudyová V. and Gregová E. (Eds.). Proc. 15th Sci. Conf. with int. particip. „New information related to genetic and breeding of agricultural crops“. 12.-13.11.2008, SCPV - VÚRV Piešťany, 19-22 (in Slovak).
- [3] Kráľová K. and Masarovičová E.: *Minimalization of hazards for the environment related to rapeseed cultivation for biofuel purpose*. Manová A., Čacho F. (eds.), Book of abstracts, 28. Sci. Conf. „Industrial toxicology 08. Tatranská Štrba, 18.06-20.6.2008, 311-315 (in Slovak).
- [4] Marchiol L., Assolari S., Sacco P. and Zerbi G.: *Phytoextraction of heavy metals by canola (Brassica napus) and radish (Raphanus sativus) grown on multi-contaminated soil*. Environ. Pollut., 2004, **132**, 21-27.
- [5] Van Ginneken L., Meers E., Guisson R., Ruttens A., Elst K., Tack F.M.G., Van Gronsveld J., Diels L. and Dejonghe W.: *Phytoremediation for heavy metal-contaminated soils combined with bioenergy production*. J. Environ. Eng. Landsc., 2007, **15**, 227-236.
- [6] Maksymiec W.: *Effect of copper on cellular processes in higher plants*. Photosynthetica, 1997, **34**, 321-342.
- [7] Panda S.K. and Choudhury S.: *Chromium stress in plants*. Braz. J. Plant Physiol., 2005, **17**, 85-98.
- [8] Punz W.F. and Sieghart H.: *The response of roots of herbaceous plant species to heavy metals*. Environ. Exp. Bot., 1993, **33**, 85-98.
- [9] Benavides M.P., Gallego S.M. and Tomaro M.L.: *Cadmium toxicity in plants*. Braz. J. Plant Physiol., 2005, **17**, 21-34.
- [10] Van Assche F. and Clijsters H.: *Effects of metals on enzyme activity in plants*. Plant Cell Environ., 1990, **13**, 195-206.
- [11] Characterization of registered rapeseed cultivars. Central Controlling and Testing Institute in Agriculture (ÚKSÚP), Bratislava 2009 (in Slovak).
- [12] Škopíková A., Kráľová K. and Masarovičová E.: *Phytotoxic effects of selenium oxoacids and some of their salts on growth of Brassica L. seedlings*. Ecol. Chem. Eng., 2008, **15**, 221-226.
- [13] Nieboer E. and Richardson D.: *The replacement of the nondescript term 'heavy metals' by a biologically and chemically significant classification of metal ions*. Environ. Pollut., B 1980, **1**, 3-26.
- [14] Nieboer E., Fletcher G.G. and Thomassen Y.: *Relevance of reactivity determinants to exposure assessment and biological monitoring of the elements*. J. Environ. Monit., 1999, **1**, 1-14.
- [15] Duffus J.H.: *“Heavy metals” - a meaningless term?* Pure Appl. Chem., 2002, **74**, 793-807.
- [16] Ivanov V.B., Bystrova E.I. and Seregin I.V.: *Comparative impacts of heavy metals on root growth as related to their specificity and selectivity*. Russ. J. Plant Physiol., 2003, **50**, 398-406.
- [17] Jiang W., Liu D. and Liu X.: *Effects of copper on root growth, cell division, and nucleolus of Zea mays*. Biol. Plant., 2001, **44**, 105-109.
- [18] Doncheva S.: *Copper-induced alterations in structure and proliferation of maize root meristem cells*. J. Plant Physiol., 1998, **153**, 482-487.
- [19] Yadav P. and Srivastava A.K.: *Cadmium induced mitotic anomalies in Hordeum vulgare and Setaria italica*. J. Environ. Biol., 1998, **19**, 25-32.
- [20] Lukatkin A.S., Gracheva N.V., Grishenkova N.N., Dukhovskis P.V. and Brazaitite A.A.: *Cytokinin-like growth regulators mitigate toxic action of zinc and nickel ions on maize seedlings*. Russ. J. Plant Physiol., 2007, **54**, 381-387.