

Upper Homeric (Silurian) high-resolution correlation using cyclostratigraphy: an example from western Lithuania

SIGITAS RADZEVIČIUS*, BRIGITA TUMAKOVAITĖ and ANDREJ SPIRIDONOV

Department of Geology and Mineralogy, Vilnius University, M.K. Čiurlionio 21/27, Vilnius, LT03101, Lithuania

**E-mail: sigitas.radzevicius@gf.vu.lt*

ABSTRACT:

Radzevičius, S., Tumakovaitė, B. and Spiridonov, A. 2017. Upper Homeric (Silurian) high-resolution correlation using cyclostratigraphy: an example from western Lithuania. *Acta Geologica Polonica*, **67** (2), 307–322. Warszawa.

The Gėluva regional stage stratigraphically corresponds to the late Wenlock. This time interval witnessed significant graptolite extinctions and turnovers of conodont faunas, as well as a large positive Mulde carbon isotopic excursion. Thus, the development of a detailed stratigraphy is a necessary step in understanding the complex patterns of regional and global variations in the sediments accumulating during the time interval studied. Therefore, in this contribution we present a cyclostratigraphic analysis of gamma ray (GR) logs from four wells, which are located in the deep water facies belt of the Lithuanian part of the Silurian Baltic Basin of the Gėluva regional stage. The analysis was performed using REDFIT spectral estimation, continuous wavelet transform and signal filtering techniques. As a result two 4th order and five 5th order cycles were distinguished and named in all sections. The correlation of cycles between sections was calibrated with the graptolite biozones. The comparative analysis revealed that intra-basinal cyclostratigraphic correlation could achieve resolution of the order of several tens of thousands of years.

Key words: Lithuania; Silurian; Upper Homeric; Cyclostratigraphy; Graptolite biozones.

INTRODUCTION

The Silurian is the shortest but one of the most dynamic periods in the Paleozoic Era of the Earth's history. There were several biotic extinctions and radiations, with climate changes, glaciations, and sea level fluctuation episodes in this period (Jeppsson 1987, 1993, 1998; Johnson *et al.* 1991; Kiipli *et al.* 2010; Aldridge *et al.* 1993; Loydell 1998, 2007; Munnecke *et al.* 2003; Spiridonov *et al.* 2015). One of such events is documented in the middle–upper Homeric (Calner *et al.* 2006; Melchin *et al.* 2012). The middle–upper Homeric spans the stratigraphical interval from the *parvus* to *ludensis* biozones (Text-fig. 1).

Although biostratigraphic, chemostratigraphic

and radiometric age dating investigations of the Silurian Period are intensive, the Silurian, as with all Paleozoic Era systems is unlucky with respect to the amount of cyclostratigraphic or astrochronologic studies performed so far (Hinnov and Ogg 2007). One of the first reports proving the Milankovitch orbital cycles for the Late Ordovician–Early Silurian is presented by G. E. Williams (1991). He, using Fourier analysis, investigated the cyclic halite and dolomite sequences of the Mallowa Salt from the Canning Basin, Western Australia. In addition, there is a magnetosusceptibility study of cyclic rock units, which links these cycles to the Earth's obliquity or eccentricity cycles at the Silurian–Devonian boundary in the Czech Republic (Crick *et al.* 2001). The ma-

System	Series	Stages	Graptolite biozones		Regional stages	Formations		
			Generalized	Lithuania		West Lithuania	Central Lithuania	East Lithuania
SILURIAN	Ludlow	Gorstian	<i>nilssoni</i>	<i>nilssoni</i>	Dubysa	Rusnė	Dubysa	Neris
			Wenlock	Homerian	<i>lundgreni</i>	<i>lundgreni</i>	Jaagarahu	Ragainė
	<i>ludensis</i>	<i>ludensis</i>			Gėluva	Siesartis	Gėluva	Širvinta
	<i>deubeli</i>	<i>deubeli</i>						Nevėžis
	<i>praedeubeli</i>	<i>praedeubeli</i>						
	<i>nassa parvus</i>	<i>nassa parvus</i>						
	Ančia Mb.	Ragainė	Riga	Birštonas				

Text-fig. 1. Correlation of the Lithuanian graptolite biozones with generalized graptolite biozones (Koren' *et al.* 1996) and with regional stages and formations

rine rhythmic sequences of the Riksu Formation in Estonia (Wenlock) were interpreted as being caused by Milankovitch cycles (Nestor *et al.* 2001). These studies did not use any statistical testing methods and were rather subjective in their approach. A more quantitative approach using Fischer plots analysis revealed the third order cycles in the Silurian of Siberia (Artyushkov and Chekhovich 2004). An investigation of Late Ordovician–Early Silurian tempestites in the Anticosti foreland basin (Canada) revealed evidence of astronomically caused cyclicality (Long 2007). There is a cyclostratigraphic study, which used gamma ray logs and time series techniques for the stratigraphical refinement of the Upper Wenlock in Lithuania (Radzevičius *et al.* 2014c). Also, cyclostratigraphical approaches were used recently in characterizing macroecological dynamics of conodonts in the middle–upper Pridoli of Lithuania (Spiridonov *et al.* 2016).


The appearance of new precise radiometric age dating (Cramer *et al.* 2012, 2015), and the latest Silurian Geochronological Scale (Melchin *et al.* 2012) allows us to distinguish and understand cyclic sedimentary patterns in the upper Homerian precisely.

The purpose of this study is to test the natural gamma ray record for the presence of cycles in the upper Homerian strata of Lithuania, and using cyclostratigraphic and graptolite biostratigraphic approaches to produce high-resolution correlation of the upper Homerian strata of the same territory. In order to achieve these goals, four core sections from outer shelfal settings were studied using spectral analytical (REDFIT, continuous wavelet transform) and signal filtering techniques (Gaussian filtering).

GEOLOGICAL BACKGROUND

During the Homerian, the Baltica palaeocontinent was located in the southern hemisphere near the equator (Cocks and Torsvik 2002). The Lithuanian territory was in the eastern part of the Baltic Silurian sedimentary basin and was located on the western part of the Baltica palaeocontinent (Text-fig. 2A).

A graptolite extinction is distinguished at the end of the *lundgreni* Biozone called the *lundgreni* Event (Koren' 1987) or the Big Crisis (German *Große Krise*) (Jaeger 1991). About 95% of graptolite species became extinct during the *lundgreni* Event (Lenz and Kozłowska-Dawidziuk 2001). Only two species of the genera *Pristiograptus* and *Gothograptus* survived the *lundgreni* extinction event, and consequently it is the biggest graptolite crisis in the Silurian Period (Cooper *et al.* 2014). Based on the conodont data, this ecological perturbation is named the Mulde Bioevent (Jeppsson *et al.* 1995). The Mulde Bioevent was distinguished following detailed investigations of Gotland conodonts, which showed the conodont turnover occurring in three event levels or datum points (Calner and Jeppsson 2003; Jeppsson and Calner 2002). Acritarchs and prasinophytes experienced reorganisation in their communities (Masiak 1998, 1999; Porębska *et al.* 2004; Calner *et al.* 2006). However, recent studies have revealed that the *lundgreni* Event (the beginning of the Mulde interval) probably had a rather minor effect on the phytoplankton (Venckutė-Aleksienė *et al.* 2016). Major reductions in radiolarian biodiversity were documented during the middle Homerian of Arctic Canada (Lenz *et al.* 2006). There are well documented changes of



Text-fig. 2. A – Facies map of the western margin of the East European Platform during *Gothograptus nassa* time (after Einasto *et al.* 1986); B – Structural map of the top the Wenlock (Bitinas 2013) and location of boreholes

other biotic groups after the *lundgreni* Event as well (Kaljo *et al.* 1996).

During the discussed event there was major perturbation of the global carbon cycle in the middle–upper Homerian, which was called the upper Homerian positive carbon isotopic ($\delta^{13}\text{C}$) excursion (Kaljo *et al.* 1997). The middle–upper Homerian positive $\delta^{13}\text{C}$ excursion for the first time was identified in the Anglo-

Welsh area (U.K.) (Corfield *et al.* 1992) and now is well known around the world. This excursion is characterized by two distinct positive peaks of $\delta^{13}\text{C}$ values (Cramer *et al.* 2011).

The Gėluva regional stage corresponds to the middle–upper Homerian in Lithuania and includes *parvus–ludensis* interval (Radzevičius 2013). The unit-stratotype of the Gėluva regional stage is in

the Jakšiai-104 borehole in the 813–776.9 m interval (Paškevičius *et al.* 1994), and the boundary-stratotype is the Vilkaviškis-61 borehole (Kojelė *et al.* 2014). There were marked facies changes across Lithuania during the Homerian, resulting in the use of three stratal schemes for the western, central and eastern parts of the country (Text-fig. 1). In western Lithuania, the upper part of the Ančia Member (Upper Riga Formation) and the Siesartis Formation belong to the Gėluva regional stage. The Ančia Member is represented by microlaminated limestone (Lapinskas and Paškevičius 1976), and the Siesartis Formation is composed of dark grey marls (Paškevičius and Lapinskas 1978). In the central part of Lithuania, the Gėluva Formation is composed of greenish grey marls and light grey nodular limestones (Lapinskas *et al.* 1985). In eastern Lithuania, the equivalent strata comprise the Nevėžis and Širvinta formations (Text-fig. 1). The Nevėžis Formation is represented by grey marls, nodular limestones, and clayey dolostones with lenses of gypsum (Lapinskas and Paškevičius 1976). The Širvinta Formation is composed of reddish, laminated dolomitic marls with lenses of gypsum (Paškevičius 1979). Previously, the upper part of the Birštonas Formation was assigned to the Gėluva regional stage and the upper part of Nevėžis Formation, with the Širvinta Formation was assigned to the Ludlow (Paškevičius 1997). According to new biostratigraphical and geochemical ($\delta^{13}\text{C}$) data, the Birštonas Formation is now assigned to the Jaagarahu regional stage and the Nėris and Širvinta formations are assigned to the Gėluva regional stage (Radzevičius *et al.* 2014b; Radzevičius *et al.* 2016).

During the studied time interval, local paleoenvironments in the studied area changed from clayey outer shelf facies in west Lithuania to the carbonatic lagunal and sabkha facies in east Lithuania (Text-fig. 2A). Moreover, the Silurian Baltic Basin experienced a long-term (second order) trend of progressive shallowing during the Silurian. In Lithuania the lateral boundaries of facies, therefore, migrated to the west (Lazauskienė *et al.* 2003).

MATERIAL

The investigated wells are located in western Lithuania (from north to the south): Šiupyliai-69 (φ 56°2'59.97"N; λ 23°5'23.99"E), Kurtuvėnai-161 (φ 55°44'36.11"N; λ 23°5'23.99"E), Viduklė-61 (φ 55°23'43.05"N; λ 22°54'37.01"E) and Vilkaviškis-131 (φ 54°45'51.94"N; λ 22°50'55.99"E). All wells were drilled in the outer shelf facies zone (Text-fig. 2B).


In all investigated boreholes the upper part of the Riga Formation with the Ančia Member was recognized. The Siesartis Formation is present in the Šiupyliai-69, Kurtuvėnai-161 and Viduklė-61 wells and the Gėluva Formation in the Vilkaviškis-131 well (Text-fig. 3). The Dubysa Formation overlaps the Siesartis and Gėluva formations. In the investigated interval, all core material studied is composed of marls with different concentrations of the clay component and different dolomitization levels. The distribution of graptolites has been studied in detail and published elsewhere (Text-fig. 3).

Several bentonites layers occur in the investigated interval of the middle–upper Homerian (Text-fig. 3). According to geochemical data, the Grötlingbo Bentonite has been identified at 1308 m depth, just above the Ančia Member, in the *parvus* Biozone in the Viduklė-61 well (Kipli *et al.* 2008). The Grötlingbo Bentonite is an important radiometrically dated bentonite layer (Cramer *et al.* 2012) of Homerian age first described from the Gotland (Snäll 1977). There are bentonite layers at the same level in the Šiupyliai-69 and Vilkaviškis-131 boreholes. The geochemical analysis of these layers has not been done, but these bentonites in the *parvus* Biozone are interpreted as representing the Grötlingbo Bentonite.

METHODS

The gamma ray log data was used to reflect the variations in the contribution of clayey and detrital component in the time series analyses. This is a first order simplification, because the cumulative gamma signal reflects the summed contribution of gamma dosage from the radioactive isotopes of U, Th, and K, which have varying geochemical and consequently sedimentological properties (Kozłowski and Sobieñ 2012). Moreover, it is widely recognized that the relative amount of terrigenous and clayey material increases in the distal parts of the Silurian Baltic Basin in comparison to the proximal parts (Paškevičius 1997). Therefore, this simplification is justified in these sedimentary settings.


In order to reveal cyclical patterns of sedimentation in the strata of the Gėluva regional stage three complimentary spectral analytical approaches were used: (1) REDFIT amplitude spectrum estimation, (2) continuous wavelet transform, and (3) band-pass signal filtering. The REDFIT (red noise fitting) spectral estimation algorithm, which utilizes the Lomb-Scargle Fourier transform, was used because it is robust to the variations in the sampling rate (Schulz



Text-fig. 3. The correlation of the upper Homerian biostratigraphy, lithology and gamma ray log of Viduklė-61 (after Radzevičius *et al.* 2014c), Vilkaviškis-161 (after Radzevičius 2006), Kurtuvėnai-161 and Šiupyliai-69 (after Radzevičius and Paškevičius 2005)

and Mudelsee 2002). The continuous wavelet transform was used because it reveals the so called “evolutionary” spectrum of a time series, where frequency and amplitude modulation can be distinguished. Performing wavelet analyses, Morlet mother wavelet was used because of its superior characteristics in determining periodicities in the frequency domain. The REDFIT and continuous wavelet spectra were estimated using the PAST (PALaeontological STATistics) statistical package (Hammer *et al.* 2001). Before the spectral analyses, in order to stationarize them, all stratigraphic series were detrended by subtracting the second order polynomial trend.

For the purposes of cyclostratigraphic subdivision and correlation in all three sections two longest (4th and 5th order) cyclicities, which were distinguished earlier in one of the investigated cores using the real part of the wavelet scalogram and Lomb periodogram, were used (Radzevičius *et al.* 2014c). Cycles as well as non-periodic features with longer duration (or stratigraphic span) are less prone to the distortions caused by the erosion, non-deposition and variations in sedimentation rates (Jerolmack and Paola 2010). In order to reconstruct the shapes of cycles from the natural gamma ray data a signal band-pass filtering procedure was performed. The natural gamma trend



Text-fig. 4. REDFIT spectra of the gamma ray logs in the investigated wells. CI – confidence interval


was filtered in two statistically significant frequency pass bands in each core. For this purpose a Gaussian filter, which is described as a “zero phase” filter, was used (Kodama and Hinnov 2014). This filter does not distort the stratigraphic positions of heights and troughs of oscillations. The filtering of signals was performed using package “astrochron” in the R statistical computational environment (Meyers 2015; R Development Core Team 2015).

Filtered statistically significant cyclicities were used in the high resolution correlation of deep water (graptolite) facies of the Lithuanian part of the Baltic Basin. The order of sedimentary cycles was determined using available chronostratigraphic information and using conventional ordering nomenclature. Cycles were called 4th order when their estimated duration was of the order of several hundreds of thousands of years, and cycles were named 5th order when their estimated duration varied from several tens to a couple of hundreds of thousands of years (Miall 2010). All distinguished 4th and 5th order cycles were named using the following nomenclature: $GI_{i,j}$. Here GI is for Gėluva regional stage, “*i*” denotes the order of the cycle (in this case 4th or 5th) and “*j*” denotes the number of a cycle. This scheme allows us to compare cycles with other stratigraphic features in a similar manner

to the event datum (Jeppsson 1998). In this way there is integration of cyclostratigraphic information into the high-resolution event stratigraphy (Cramer *et al.* 2015). Cycles are non-unique in their nature (Miall 2010) in the sense that they are distinguished using only information about their amplitude and phase relations. Thus there is a possibility of missing some cycles. In the future there is a possibility that some gaps in the record will be distinguished. Our scheme could easily accommodate such changes by interpolating new cycles between distinguished ones (i.e. if there is an additional cycle of the same order between cycles $GI_{5,1}$ and $GI_{5,2}$ it will be named $GI_{5,1.1}$).

RESULTS

In this study the two cyclicities with the lowest frequencies and the highest power were focused on. According to the REDFIT analysis (Text-fig. 4), there are several additional high frequency cyclicities whose spectral power passed the 95 and 99% confidence intervals. However, high frequency features, as was mentioned in the methods section, are more sensitive to sedimentological distortion. The period lengths of 4th order cycles in all four analysed sections



Text-fig. 5. Upper Homeric biostratigraphy, lithology, gamma-ray log data, wavelet scalogram (colour intensity shows spectral strength) and filtered cycles in the Viduklė-61 well. Legend see Text-fig. 3

reached 20 m. The 5th order cycles were more variable in their period length: Viduklė-61, Šiupiliai-69, and Vilkaviškis-131 had 6.7 m long cycles; and Kurtuvėnai-161 had 7.5 m long 5th order cycles.

Cyclicities were distinguished in the Jaagarahu, Gėluva and Dubysa regional stages. In this study only the cyclicities of the Gėluva regional stage are

focused on. The descriptions of the cycles of the Gėluva regional stage are given below.

Viduklė-61 well

Two 4th order cycles of the Gėluva regional stage in Viduklė-61 well were distinguished (Text-fig. 5).

The base of first 4th order cycle (Gl_{4,1}) coincides with the minimum values of GR in the Ančia Member in the lowermost part of the *parvus* Biozone. The Gl_{4,1} ended in the upper part of *praedeubeli* Biozone. The base of Gl_{4,2} cycle coincides with top of Gl_{4,1} cycle. The top of the Gl_{4,2} cycle coincides with the base of the *nilssoni* Biozone and the Dubysa regional stage.

Five 5th order cycles are distinguished in the Gėluva regional stage (Text-fig. 5). The base of Gl_{5,1} is a little bit lower than the base of Gl_{4,1} cycle in the base of the Ančia Member. The cycle Gl_{5,1} ends in the topmost part of the *parvus* Biozone. Gl_{5,2} cycle corresponds to the *nassa* Biozone. The base of the Gl_{5,3} coincides with the *praedeubeli* Biozone. The cycle Gl_{5,4} corresponds to the *deubeli* Biozone. The base of Gl_{5,4} is near the base of the *ludensis* Biozone and it ends a little bit lower than the *nilssoni* Biozone and here coincides with the top of Gl_{4,2} cycle boundary.

The distinguished cycles of the Viduklė-61 well are used as the typical or standards for the Gėluva regional stage. Viduklė-61 well has been chosen because it includes the boundary stratotype of the Gėluva regional stage (Kojelė et al. 2014), the type section of the Siesartis Formation (Paškevičius and Lapinskas 1978), and this core is well explored chemo- (Martma et al. 2005) and biostratigraphically (Radzevičius et al. 2014a).

Kurtuvėnai-161 well

Two 4th order cycles were distinguished in the Gėluva regional stage in the Kurtuvėnai-161 well (Text-fig. 6). The base of Gl_{4,1} is in the Ančia Member, the *parvus* Biozone. The top of Gl_{4,1} cycle is in the upper part of the *praedeubeli* Biozone. The top of Gl_{4,2} cycle is near the *nilssoni* Biozone base. It is difficult to correlate the top of Gl_{4,2} with the graptolite biozones because graptolites are absent in the 1297.8–1284.7 m interval, but the first appearance of *Neodiversograptus nilssoni* (Barrande) marking the beginning of the Ludlow is at a depth of 1284.7 m. Therefore the lower boundary of the *nilssoni* Biozone cannot be higher in the section.

Five 5th order cycles are distinguished in the Gėluva regional stage interval. The base of Gl_{5,1} is lower than the base of Gl_{4,1} cycle, near the base of the *parvus* Biozone (Text-fig. 6). The Gl_{5,2} starts in the *nassa* Biozone and ends in the lower part of *praedeubeli* Biozone. The base of Gl_{5,3} coincides with the top of Gl_{5,2} cycle. The top of Gl_{5,3} is in the upper part of the *praedeubeli* Biozone, higher than the base of Gl_{4,2} cycle. The top position of

Gl_{5,4} is difficult to correlate with the graptolite biozones too. There are no graptolites in the interval, although the Gl_{5,4}, and the Gl_{5,5} cycles can be related to the *deubeli* and *ludensis* graptolite biozones. The top of Gl_{5,5} cycle is lower than the top of Gl_{4,2} cycle, near the boundary of the Gėluva and Dubysa regional stages.

The numbers and boundary positions of the 4th and 5th order cycles of the Gėluva regional stage in the Kurtuvėnai-161 well are similar to those in the Viduklė-61 well.


Šiupyliai-69 well

Two 4th order cycles of the Gėluva regional stage is distinguished in the Šiupyliai-69 well (Text-fig. 7). The base of Gl_{4,1} cycle is in the Ančia Member, near the base of the *parvus* Biozone. The top of Gl_{4,1} is in the lower part of the *deubeli* Biozone. The Gl_{4,2} cycle is distinguished from the lower part of the *deubeli* Biozone to the middle *nilssoni* Biozone. The top of Gl_{4,2} is about 10 m higher than the Gėluva and Dubysa regional stage boundary. Therefore significant part of the Gl_{4,2} cycle is in the Dubysa regional stage. According to the described patterns, the upper boundary of Gl_{4,2} in Šiupyliai-69 well is higher than in Viduklė-61 and Kurtuvėnai-161 wells.

Five 5th order cycles are distinguished in the Gėluva regional stage (Text-fig. 7). The base of Gl_{5,1} is in the Ančia Member, in the middle part of the *parvus* Biozone. It is a little bit higher than the base of Gl_{4,1} cycle. The base of Gl_{5,2} is in the middle part of the *nassa* Biozone. The cycle Gl_{5,3} is in the middle *praedeubeli* and middle *deubeli* biozones interval. The base of Gl_{5,4} cycle is in the middle part of the *deubeli* Biozone and the top in the middle of the *ludensis* Biozone. The top of Gl_{5,5} is higher than the boundary of the Gėluva and the Dubysa regional stages, in the lower part of the *nilssoni* Biozone.

Vilkaviškis-131 well

There are two 4th order cycles in the Gėluva regional stage of the Vilkaviškis-131 well section (Text-fig. 8). The base of Gl_{4,1} is in the Ančia Member at the lower boundary of the *parvus* Biozone. Gl_{4,1} ends in the lower part of *deubeli* Biozone. The base of Gl_{4,2} is in the *deubeli* Biozone. The position of the Gl_{4,2} cycle upper boundary is difficult to correlate with the graptolite biozones. In this part of the section (starting from 1067.3 m depth) graptolites are absent. Therefore the boundary between the Gėluva




Text-fig. 6. Upper Homeric biostratigraphy, lithology, gamma-ray log data, wavelet scalogram and filtered cycles in the Kurtuvėnai-161 well. Legend see Text-fig. 3

and the Dubysa regional stages are distinguished based on the lithological changes (Gėluva–Dubysa formations).

There are five 5th order cycles in the studied interval (Text-fig. 8). The base of Gl_{5,1} is in the upper part of the *parvus* Biozone, a little bit higher than the base of the Gl_{4,1} cycle. The base of Gl_{5,2} is located in

the upper part of the *nassa* Biozone. The Gl_{5,3} corresponds to the *deubeli* Biozone. There is no possibility of correlating Gl_{5,4} and Gl_{5,5} with the graptolite biozones because graptolites are absent there. However, the upper boundary of Gl_{5,5} is lower than the upper boundary of the Gl_{4,2} cycle, near the Gėluva and Dubysa regional stages boundary.




Text-fig. 7. Upper Homeric biostratigraphy, lithology, gamma-ray log data, wavelet scalogram and filtered cycles in the Šiupyliai-69 well. Legend see Text-fig. 3

DISCUSSION AND CONCLUSIONS

All studied middle–upper Wenlock sections yielded the same set of 4th and 5th order cycles: two long and five shorter cycles each (Text-fig. 9). Lower (4th) order cycles (GI_{4,1} and GI_{4,2}) are broadly compa-


rable to the Ho2 and Ho3 sequences recognized in different sections of Europe and North America (Barrick 1997; Calner 1999; Cramer *et al.* 2011; Cramer *et al.* 2012; Marshall *et al.* 2012; Ray and Butcher 2010). However, according to the precise cyclostratigraphical placement, based on the spectral analysis and sig-



Text-fig. 8. Upper Homeric biostratigraphy, lithology, gamma-ray log data, wavelet scalogram and filtered cycles in the Vilkaiviškis-131 well. Legend see Text-fig. 3

nal filtering, the base of the $Gl_{4,2}$ cycle is stratigraphically lower than that proposed for the comparable sequence Ho3. In our case the maximum regression is observed in the *praedeubeli* or *deubeli* biozones and not in the *ludensis* Biozone as was shown by Cramer *et al.* (2011). The differences could be of a double

nature. Firstly, a difference could arise because of different scaling schemes for graptolite zones. Secondly, stratigraphical sequences are distinguished based on mainly qualitative observations and typological schemes, (e.g., maximum regressive surface), which could conflate dynamic features that belong to dif-



Text-fig. 9. Correlation of 4th and 5th order cycles with graptolite biozones in the studied boreholes

ferent hierarchical orders when they superpose each other (Catuneanu 2006). Spectral analyses, on the other hand, are based on the Fourier theorem, which states that the studied continuous signal could be decomposed into the sum of independent components (simple functions), representing amplitude contributions of different frequencies of the signal.

It should be noted that the occurrence of two 4th order cycles in the Gėluva regional stage corresponds well to the bentonite based correlation of the lower two sedimentary cycles (Iide Beds and Viita+Kuusnõmme Beds) in the Rootsikũla Stage of the shallow water Estonian sections in the upper part of the Homerian (Kiipli *et al.* 2011).

The 4th order cycles revealed could be correlated between the different wells studied with high accuracy and precision. However, some features, especially in the upper part of the section, have a lower stratigraphic time significance potential. The base of the Gl_{4,1} cycle corresponds approximately to the lower boundary of the *parvus* Biozone. The base of the Gl_{4,2} cycle in the type section (Viduklė-61) and also in the Kurtuvėnai-161 section is distinguished in the middle of the *praedeubeli* Biozone. In the more near-shore successions of the Šiupyliai-69 and the Vilkaviškis-131 wells the base of the Gl_{4,2} cycle is observed stratigraphically higher – in the *deubeli* Biozone. Similarly in those two sections the upper boundary of the Gl_{4,2} cycle is offset further into the Ludlow. This phase delay of cycles in shallower sections in comparison to deeper sections is also observed in the patterns of the 5th order cycles. For example, in the Vilkaviškis-131 succession 5th order cycles are significantly (by 180°) displaced in phase in comparison to the Viduklė-61 section.

There could be several possible reasons for differences in phase relations between cycles, which are driven by the same forcing mechanism, i.e., Milankovitch mechanisms (Herbert 1994; Hinnov and Hilgen 2012). One of the possible sources of non-in-phase responses of sedimentary systems is progressive variation in the potential of carbonate production on the offshore-nearshore gradient. This could be related to the variations in pH and also in the availability of light for photosymbiotic carbonate building organisms. This could result in non-linear fluxes of carbonates as a response to possibly linear external forcing. It was shown that the sensitivity of carbonate content (which is approximately inversely reflected in the gamma ray curves) as a palaeoclimatic proxy strongly depends on the average carbonate concentrations in the studied section (Herbert 1994). Changes in accommodation spaces due to eustatic variations also could cause out of phase sedimentation patterns in different parts of a basin (Kidwell 1988), as well as spatial colonization dynamics of carbonate material producing organisms, which react to the sea level perturbations in a lagged manner (Tipper 1997).

The patterns of cyclic sedimentation revealed, which are constrained by high-resolution graptolite stratigraphy, undoubtedly show a great potential of spectral method based cyclostratigraphy in the correlation of offshore strata of the Silurian Period. Based on the previous estimations of the durations of the cycles of the Gėluva regional stage (Radzevičius *et al.* 2014c), core sections could be correlated with

the stratigraphic precision in order of ~ 70 Ka (approximately half of a 5th order cycle). The combination of cyclostratigraphy with other tools, such as biostratigraphy and δ¹³C chemostratigraphy could reduce this correlation error even further.

Acknowledgements

This study is a part of the project “Event Stratigraphy in the Silurian Sedimentary Basin of Lithuania” and a contribution to the IGCP-591 (The Early to Middle Palaeozoic Revolution). This study was sponsored by the Lithuanian Academy of Sciences grant for young scientist (A.S.). We would like to thank T. Kiipli (Tallinn University of Technology) and W. Kozłowski (University of Warsaw) for significantly improving the manuscript.

REFERENCES

- Aldridge, R., Jeppsson, L. and Dornig, K. 1993. Early Silurian oceanic episodes and events. *Journal of the Geological Society*, **150**, 501–513.
- Artushkov, E.V. and Chekhovich, P.A. 2004. Mechanisms of sea depth changes in Silurian epeiric basins of East Siberia. *Geologiya i Geofizika (Russian Geology and Geophysics)*, **45** (11), 1273–1291 (1219–1236).
- Barrick, J.E. 1997. Wenlock (Silurian) depositional sequences, eustatic events, and biotic change on the southern shelf of North America. *Geological Society of America Special Papers*, **321**, 47–66.
- Bitinas, J. 2013. Structural map of the top of Upper Wenlockian of Lower Silurian of the West Lithuania area. Lithuanian Geological Survey.
- Calner, M. 1999. Stratigraphy, facies development, and depositional dynamics of the Late Wenlock Fröjel Formation, Gotland, Sweden. *GFF*, **121**, 13–24.
- Calner, M. and Jeppsson, L. 2003. Carbonate platform evolution and conodont stratigraphy during the middle Silurian Mulde Event, Gotland, Sweden. *Geological Magazine*, **140**, 173–203.
- Calner, M., Kozłowska, A., Masiak, M. and Schmitz, B. 2006. A shoreline to deep basin correlation chart for the middle Silurian coupled extinction-stable isotopic event. *GFF*, **128**, 79–84.
- Catuneanu, O. 2006. Principles of sequence stratigraphy, 386 pp. Elsevier; Amsterdam.
- Cooper, R.A., Sadler, P.M., Munnecke, A. and Crampton, J.S. 2014. Graptoloid evolutionary rates track Ordovician Silurian global climate change. *Geological Magazine*, **151**, 349–364.
- Corfield, R.M., Siveter, D.J., Cartlidge, J.E. and McKerrow,

- W.S. 1992. Carbon isotope excursion near the Wenlock–Ludlow (Silurian) boundary in the Anglo-Welsh area. *Geology*, **20**, 371–374.
- Cramer, B.D., Condon, D.J., Söderlund, U., Marshall, C., Worton, G.J., Thomas, A.T., Calner, M., Ray, D.C., Perrier, V. and Boomer, I. 2012. U-Pb (zircon) age constraints on the timing and duration of Wenlock (Silurian) paleocommunity collapse and recovery during the “Big Crisis”, *Geological Society of America Bulletin*, **124**, 1841–1857.
- Cramer, B.D., Vandenbroucke, T.R.A. and Ludvigson, G.A. 2015. High-Resolution Event Stratigraphy (HiRES) and the quantification of stratigraphic uncertainty: Silurian examples of the quest for precision in stratigraphy. *Earth-Science Reviews*, **141**, 136–153.
- Cramer, B.D., Brett, C.E., Melchin, M.J., Maennik, P., Kleffner, M.A., McLaughlin, P.I., Loydell, D.K., Munnecke, A., Jeppsson, L. and Corradini, C. 2011. Revised correlation of Silurian Provincial Series of North America with global and regional chronostratigraphic units and $\delta^{13}\text{C}_{\text{carb}}$ chemostratigraphy. *Lethaia*, **44**, 185–202.
- Crick, R.E., Ellwood, B.B., Hladil, J., El Hassani, A., Hroudra, F. and Chlupáč, I. 2001. Magnetostratigraphy susceptibility of the Pridolian–Lochkovian (Silurian–Devonian) GSSP (Klonk, Czech Republic) and a coeval sequence in Anti-Atlas Morocco. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **167**, 73–100.
- Einasto, P.E., Abushik, A.F., Kaljo, D.L., Koren, T.N., Modzalevskaya, T.L. and Nestor, H.E. 1986. Silurian sedimentation and the fauna of the East Baltic and Podolian marginal basin: a comparison. In: D.L. Kaljo and E. Klaamann (Eds), *Theory and Practice of Ecostratigraphy*, pp. 65–72. Institute of Geology, Academy of Sciences of the Estonian SSR; Tallin. [In Russian]
- Hammer, Ø., Harper, D.A.T. and Ryan, P.D. 2001. PAST-Palaeontological Statistics, ver. 3.0. *Palaeontologia Electronica*, **4**, 9.
- Herbert, T.D. 1994. Reading orbital signals distorted by sedimentation: models and examples. In: P.L. DeBoer and D.G. Smith (Eds), *Orbital Forcing and Cyclic Sequences*, pp. 483–507. Wiley; Oxford.
- Hinnov, L.A. and Hilgen, F.J. 2012. Cyclostratigraphy and astrochronology. In: F.M. Gradstein, J.G. Ogg and M. Schmitz (Eds.), *The Geologic Time Scale 2012*, 2-volume set, pp. 63–83. Elsevier; Amsterdam.
- Hinnov, L.A. and Ogg, J.G. 2007. Cyclostratigraphy and the Astronomical Time Scale. *Stratigraphy*, **4**, 239–251.
- Jaeger, H. 1991. Neue Standard-Graptolithenzonenfolge nach der ‘Großen Krise’ an der Wenlock/Ludlow-Grenze (Silur). *Neues Jahrbuch für Geologie und Paläontologie Abhandlungen*, **182**, 303–354.
- Jeppsson, L. 1998. Silurian oceanic events: summary of general characteristics. In: J.M. Landing (Ed.), *Silurian cycles: Linkages of dynamic stratigraphy with atmospheric, oceanic and tectonic changes*, pp. 239–257. New York State Museum; New York.
- Jeppsson, L. 1987. Lithological and conodont distributional evidence for episodes of anomalous oceanic conditions during the Silurian. In: R.J. Aldridge (Ed.), *Palaeobiology of Conodonts*, pp. 129–145. Ellis Horwood Ltd; Chichester.
- Jeppsson, L. 1993. Silurian events: the theory and the conodonts. *Proceedings of the Estonian Academy of Sciences*, **42**, 23–27.
- Jeppsson, L. 1998. Silurian oceanic events: summary of general characteristics. In: E.J.M. Landing (Ed.), *Silurian cycles: Linkages of dynamic stratigraphy with atmospheric, oceanic and tectonic changes*, pp. 239–257. New York State Museum; New York.
- Jeppsson, L., Aldridge, R. and Dorning, K. 1995. Wenlock (Silurian) oceanic episodes and events. *Journal of the Geological Society*, **152**, 487–498.
- Jeppsson, L. and Calner, M. 2002. The Silurian Mulde Event and a scenario for secundo-secundo events. *Earth and Environmental Science Transactions of the Royal Society of Edinburgh*, **93**, 135–154.
- Jerolmack, D.J. and Paola, C. 2010. Shredding of environmental signals by sediment transport. *Geophysical Research Letters*, **37**, 1–19.
- Johnson, M.E., Kaljo D. and Rong, J.-Y. 1991. Silurian eustasy. *Special Papers in Palaeontology*, **44**, 145–63.
- Kaljo, D., Boucot, A.J., Corfield, R.M., Le Herisse, A., Koren, T.N., Kriz, J., Männik, P., Märss, T., Nestor, V., Shaver, R.H., Siveter, D.J. and Viira, V. 1996. Silurian bio-events. In: O.H. Walliser (Ed.), *Global events and event stratigraphy in the Phanerozoic*, pp. 173–224. Springer; Berlin.
- Kaljo, D., Kiipli, T. and Martma, T. 1997. Carbon isotope event markers through the Wenlock-Pfidioli sequence in Ohesaare (Estonia) and Priekule (Latvia). *Palaeogeography, Palaeoclimatology, Palaeoecology*, **132**, 211–224.
- Kidwell, S.M. 1988. Reciprocal sedimentation and noncorrelative hiatuses in marine-paralic siliciclastics: Miocene outcrop evidence. *Geology*, **16**, 609–612.
- Kiipli, T.R., Einasto, T., Kallaste, V., Nestor, Perens, H. and Siir, S. 2011. Geochemistry and correlation of volcanic ash beds from the Rootsiküla Stage (Wenlock–Ludlow) in the eastern Baltic. *Estonian Journal of Earth Sciences*, **60** (4), 207–219.
- Kiipli, T., Kiipli, E. and Kaljo, D. 2010. Silurian sea level variations estimated using $\text{SiO}_2/\text{Al}_2\text{O}_3$ and $\text{K}_2\text{O}/\text{Al}_2\text{O}_3$ ratios in the Priekule drill core section, Latvia. *Bollettino della Società Paleontologica Italiana*, **49**, 55–63.
- Kiipli, T., Radzevičius, S., Kallaste, T., Motuza, V., Jeppsson, L. and Wickström, M.L. 2008. Wenlock bentonites in Lithuania and correlation with bentonites from sections in Estonia, Sweden and Norway. *GFF*, **130**, 203–210.
- Kodama, K.P. and Hinnov, L.A. 2014. *Rock Magnetic Cyclostratigraphy*, pp. 1–165. Wiley-Blackwell; Chichester.
- Kojelė, A., Radzevičius, S., Spiridonov, A. and Brazauskas, A.

2014. Gėluvos regioninio aukšto ribos (viršutinis homeris, silūras) stratotipas. *Geologijos akiračiai*, **1**, 10–21.
- Koren', T.N., Lenz, A.C., Loydell, D.K., Melchin, M.J., Stroch, P. and Teller, L. 1996. Generalized graptolite zonal sequence defining Silurian time intervals for global paleogeographic studies. *Lethaia*, **29**, 59–60.
- Koren', N.T. 1987. Graptolite dynamics in Silurian and Devonian time. *Bulletin of the Geological Society of Denmark*, **35**, 149–159.
- Kozłowski, W. and Sobień, K., 2012. Mid-Ludfordian coeval carbon isotope, natural gamma ray and magnetic susceptibility excursions in the Mielnik IG-1 borehole (Eastern Poland) – dustiness as a possible link between global climate and the Silurian carbon isotope record. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **339**, 74–97.
- Lapinskas, P. and Paškevičius, J. 1976. Silurian stratigraphic scheme of South Baltic. In: *Materialy po stratigrafii Pribaltiki*, pp. 44–45. LitNIGRI; Vilnius. [In Russian]
- Lapinskas, P., Paškevičius, J. and Jacyna, J. 1985. New data on the stratigraphy and correlation of the South-West Baltic Wenlockian and Ludlovian (Silurian). *Geologija* (Vilnius), **6**, 29–40.
- Lazauskienė, J., Šliaupa, S., Brazauskas, A. and Musteikis, P. 2003. Sequence stratigraphy of the Baltic Silurian succession; tectonic control on the foreland infill. *Geological Society London Special Publications*, **208**, 95–115.
- Lenz, A.C. and Kozłowska-Dawidziuk, A. 2001. Upper Wenlock (Silurian) graptolites of Arctic Canada: pre-extinction, *lundgreni* Biozone fauna. *Palaeontographica Canadiana*, **20**, 1–61.
- Lenz, A.C., Noble, P.J., Masiak, M., Poulson, S.R. and Kozłowska, A. 2006. The *lundgreni* Extinction Event: integration of paleontological and geochemical data from Arctic Canada. *GFF*, **128**, 153–158.
- Long, D.G.F. 2007. Tempestite frequency curves: a key to Late Ordovician and Early Silurian subsidence, sea-level change, and orbital forcing in the Anticosti foreland basin, Quebec, Canada. *Canadian Journal of Earth Sciences*, **44**, 413–431.
- Loydell, D.K. 1998. Early Silurian sea-level changes. *Geological Magazine*, **135**, 447–471.
- Loydell, D.K. 2007. Early Silurian positive $\delta^{13}\text{C}$ excursions and their relationship to glaciations, sea-level changes and extinction events. *Geological Journal*, **42**, 531–546.
- Marshall, C., Thomas, A., Boomer, I. and Ray, D. 2012. High resolution $\delta^{13}\text{C}$ stratigraphy of the Homeric (Wenlock) of the English Midlands and Wenlock Edge. *Bulletin of Geosciences*, **87**, 669–679.
- Masiak, M. 1998. Preliminary data on Silurian acritarch assemblages from the Holly Cross Mountains (Central Poland). *Temas Geológico-Mineros ITGE*, **23**, 105–106.
- Masiak, M. 1999. Zapis zdarzenia *Cyrtograptus lundgreni* w górnolenockim zespole akritarchowym Gór Świętokrzyskich. *Przegląd Geologiczny*, **47**, 359–360.
- Melchin, M.J., Sadler, P.M. and Cramer, B.D. 2012. The Silurian Period, In: F.M. Gradstein, J.G. Ogg, M. Schmitz, G. Ogg (Eds), *The Geologic Time Scale 2012*, 2-volume set, pp. 525–558. Elsevier; Amsterdam.
- Meyers, S.R. 2015. *Astrochron: A Computational Tool for Astrochronology* (Version 0.5).
- Miall, A.D. 2010. *The Geology of Stratigraphic Sequences* (2nd edition), pp. 1–522. Springer; Berlin.
- Munnecke, A., Samtleben, C. and Bickert, T. 2003. The Ireviken Event in the lower Silurian of Gotland, Sweden – relation to similar Palaeozoic and Proterozoic events. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **195**, 99–124.
- Nestor, H., Einasto, R., Nestor, V., Marss, T. and Viira, V. 2001. Description of the type section, cyclicity and correlation of the Riksu Formation (Wenlock, Estonia). *Proceedings of the Estonian Academy of Sciences, Geology*, **50**, 149–173.
- Paškevičius, J. and Lapinskas, P. 1978. New Silurian stratotypes of South Baltic States. *Nauchnye trudy vysshikh uchebnykh zavedenij Litovskoj SSR, Geologiya i Geografiya*, **14**, 123–128. [In Russian].
- Paškevičius, J. 1979. Biostratigraphy and graptolites of the Lithuanian Silurian, pp. 1–267. Mokslas; Vilnius.
- Paškevičius, J. 1997. The Geology of the Baltic Republics, pp. 1–387. Vilnius University and Geological Survey of Lithuania; Vilnius.
- Paškevičius, J., Lapinskas, P., Brazauskas, A., Musteikis, P. and Jacyna, J. 1994. Stratigraphic revision of the regional stages of the Upper Silurian part in the Baltic Basin. *Geologija* (Vilnius), **17**, 64–87.
- Porębska, E., Kozłowska-Dawidziuk, A. and Masiak, M. 2004. The *lundgreni* event in the Silurian of the East European Platform, Poland. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **213**, 271–294.
- R Development Core Team 2015. *R: A Language and Environment for Statistical Computing*. Version 3.1.3. R Foundation for Statistical Computing; Vienna.
- Radzevičius, S. 2006. Late Wenlock biostratigraphy and the *Pristiograptus virbalensis* group (Graptolithina) in Lithuania and the Holy Cross Mountains. *Geological Quarterly*, **50**, 333–344.
- Radzevičius, S. 2013. Silurian graptolite biozones of Lithuania: present and perspectives. *Geologija* (Vilnius), **55**, 41–49.
- Radzevičius, S. and Paškevičius, J. 2005. *Pristiograptus* (Graptoloidea) from the Upper Wenlock of the Baltic Countries. *Stratigraphy and Geological Correlation*, **13**, 159–169.
- Radzevičius, S., Spiridonov, A. and Brazauskas, A. 2014a. Integrated middle–upper Homeric (Silurian) stratigraphy of the Viduklė-61 well, Lithuania. *GFF*, **136**, 218–222.
- Radzevičius, S., Spiridonov, A., Brazauskas, A., Norkus, A., Meidla, T. and Ainsaar, L. 2014b. Upper Wenlock $\delta^{13}\text{C}$ chemostratigraphy, conodont biostratigraphy and palaeoecology

- gical dynamics in the Ledai-179 drill core (Eastern Lithuania). *Estonian Journal of Earth Sciences*, **63**, 293–299.
- Radzevičius, S., Spiridonov, A. and Brazauskas, A. 2014 c. Application of Wavelets to the Cyclostratigraphy of the Upper Homeric (Silurian) Gėluva Regional Stage in the Viduklė-61 Deep Well (Western Lithuania). In: J. Pais (Ed.), STRATI 2013, pp. 437–440. Springer, Berlin.
- Radzevičius, S., Spiridonov, A., Brazauskas, A., Dankina, D., Rimkus, A., Meidla, T. and Ainsaar, L. 2016. Integrated stratigraphy, conodont turnover and palaeoenvironments of the upper Wenlock and Ludlow in the shallow marine succession of the Vilkaviškis-134 core (Lithuania). *Newsletters on Stratigraphy*, **49**, 321–336.
- Ray, D. and Butcher, A. 2010. Sequence stratigraphy of the type Wenlock area (Silurian), England. *Bollettino della Società Paleontologica Italiana*, **49**, 47–54.
- Schulz, M. and Mudelsee, M. 2002. REDFIT: estimating red-noise spectra directly from unevenly spaced paleoclimatic time series. *Computers & Geosciences*, **28**, 421–426.
- Snäll, S. 1977. Silurian and Ordovician bentonites of Gotland (Sweden). *Stockholm Contributions in Geology*, **31**, 1–80.
- Spiridonov, A., Brazauskas, A. and Radzevičius, S. 2015. The Role of Temporal Abundance Structure and Habitat Preferences in the Survival of Conodonts during the Mid–Early Silurian Ireviken Mass Extinction Event. *PLoS ONE*, **10** (4): e0124146.
- Spiridonov, A., Brazauskas, A. and Radzevičius, S. 2016. Dynamics of abundance of the mid- to late Pridoli conodonts from the Eastern part of the Silurian Baltic Basin: multifractals, state shifts, and oscillations. *American Journal of Science*, **316**, 363–400.
- Tipper, J. C. 1997. Modeling carbonate platform sedimentation – lag comes naturally. *Geology*, **25**, 495–498.
- Venckutė-Aleksienė, A., Radzevičius, S. and Spiridonov, A. 2016. Dynamics of phytoplankton in relation to the upper Homeric (Lower Silurian) *lundgreni* event – an example from the Eastern Baltic Basin (Western Lithuania). *Marine Micropaleontology*, **126**, 31–41.
- Williams, G.E. 1991. Milankovitch-band cyclicity in bedded halite deposits contemporaneous with Late Ordovician–Early Silurian glaciation, Canning Basin, Western Australia. *Earth and Planetary Science Letters*, **103**, 143–155.

Manuscript submitted: 9th February 2016

Revised version accepted: 2nd February 2017