

DEPOSITIONAL ENVIRONMENTS, FACIES AND DIAGENESIS OF THE UPPER JURASSIC–LOWER CRETACEOUS CARBONATE DEPOSITS OF THE BUILA-VÂNTURARIȚA MASSIF, SOUTHERN CARPATHIANS (ROMANIA)

George PLEȘ, Ioan I. BUCUR & Emanoil SĂSĂRAN

Department of Geology, Babeș-Bolyai University, M. Kogălniceanu 1, 400084 Cluj-Napoca, Romania; e-mails: george.ples@ubbcluj.ro; ioan.bucur@ubbcluj.ro; emanoil.sasaran@ubbcluj.ro

Pleș, G., Bucur, I. I. & Săsăran, E., 2016. Depositional environments, facies and diagenesis of the Upper Jurassic–Lower Cretaceous carbonate deposits of the Buila-Vânturarița Massif, Southern Carpathians (Romania). *Annales Societatis Geologorum Poloniae*, 86: 165–183.

Abstract: The Buila-Vânturarița Massif consists of massive Upper Jurassic reef limestones (Kimmeridgian–Tithonian) and Lower Cretaceous (Berriasian–Valanginian, and Barremian–?Lower Aptian) deposits. Besides corals and stromatoporoids, a wide range of micro-encrusters and microbialites has contributed to their development. In this study, the authors describe briefly and interpret the main facies associations and present the microfossil assemblages that are important for age determination. The distribution of facies associations, corroborated with the micropalaeontological content and early diagenetic features, indicate different depositional environments. The carbonate successions show the evolution of the Late Jurassic–Early Cretaceous depositional environments from slope and reef-front to internal-platform sedimentary settings, including peritidal environments in the lowermost Cretaceous. Early diagenesis, represented by syndimentary cementation in the form of micritization (including cement crusts in the reef microframework), followed by dissolution, cementation and dolomitization in a meteoric regime, and void-filling late cementation during the burial stage.

Key words: Carbonate platforms, reefs, microfacies, micro-encrusters, carbonate diagenesis, Upper Jurassic, lowermost Cretaceous, Southern Carpathians, Romania.

Manuscript received 9 February 2015, accepted 14 April 2016

INTRODUCTION

The Buila-Vânturarița Massif, located in the central-northern part of Vâlcea County, Romania, is part of the Căpățâni Mountains, central Southern Carpathians (Fig. 1). Its lithology, geological history and geomorphology are unique by comparison with the features of the main mountain chain in the Căpățâni Mountains. The Buila-Vânturarița Massif consists of a 12-km-long and 0.5- to 2-km-wide, linear, calcareous crest, mainly built of Jurassic limestones. It is a fragment of the carbonate sedimentary system, known as the Getic Carbonate Platform (Săndulescu, 1984), which covered the Getic-Supragetic domain during the Late Jurassic–Early Cretaceous. The outline and NNE–SSW orientation of the morphology of the main Buila-Vânturarița crest are similar to those of the Piatra Craiului Massif, another part of the Getic Carbonate Platform, located more to the east. The studied massif is dominated by Kimmeridgian–Tithonian reef limestones. Synchronous limestones of other areas in the Tethys realm are known as “Štramberk”-type limestones in the Carpathians (see Uță and Bucur, 2003;

Șerban *et al.*, 2004; Bucur and Săsăran, 2005; Săsăran, 2006; Ivanova *et al.*, 2008; Pleș *et al.*, 2013; Săsăran *et al.*, 2014; Kołodziej, 2015a, b and literature therein), and as “Plassen”-type limestones in the Alps (e.g., Schlagintweit and Gawlick, 2008; Schlagintweit *et al.*, 2005). The term “Štramberk”-type limestones was even recently applied to limestones in Turkey (Masse *et al.*, 2015). In many occurrences, as is also the case with the Buila-Vânturarița Massif, the reef limestones are conformably overlain by lowermost Cretaceous (Berriasian–Valanginian) stratified limestones, followed by transgressive Urgonian limestones, Barremian–Lower Aptian in age (Dragastan, 2010; Pleș *et al.*, 2013; Mircescu *et al.*, 2014; Pleș and Schlagintweit, 2014). The aim of this paper is the identification and interpretation of the main microfacies types, micropalaeontological assemblages and diagenetic processes that affected these limestones, in order to reconstruct their depositional environments.




Fig. 1. Geological map of the Buila-Vânturarița Massif (modified after Lupu et al., 1978).

GEOLOGICAL SETTING

The Buila-Vânturarița Massif is part of the former Getic Nappe, a structural unit within the Median Dacides, in the Southern Carpathians (Săndulescu, 1984). In this massif, post-Triassic sedimentary deposits are mainly represented by Middle Jurassic–Oxfordian detrital and siliciclastic rocks, overlain by Kimmeridgian–lowermost Cretaceous reef limestones (Fig. 1). Boldur *et al.* (1968) identified Bajocian deposits in the south-eastern area. The Bajocian age is supported by the assemblages of ostreids, Phylloceratidae ammonites, and bivalve species of *Entolium* Meek, 1865 and *Camptochlamys* Arkell, 1930 (Dragastan, 2010). Bathonian deposits are present in smaller areas in the Buila-Vânturarița Massif and are often associated with Lower Callovian ones. Boldur *et al.* (1968) identified specimens of *Oecotraustes* sp. in a horizon of calcareous sandstones in the Bistrița Valley, indicating the presence of Bathonian–Callovian rocks within the succession. Boldur *et al.* (1968, 1970) identified Callovian deposits in the central-western part of the massif, in the Bistrița and Costești valleys, as well as south of Stogu Peak. These authors have described the following lithological succession for the Bathonian–Callovian: micaceous calcareous sandstones with *Oecotraustes* (Upper Bathonian–Lower Callovian); reddish and yellowish micaceous calcareous sandstones with *Grossouvria curvicosta* (Opper, 1857) and *G. subtilis* (Neumayr, 1871), and sandy limestones with *Bositra buchi* (Roemer, 1836) and *Phylloceras* sp. (Middle–Upper Callovian). Sediments assigned to the Oxfordian occur across relatively larger areas compared to the Middle Jurassic equivalents. They were grouped into the Buila and Bistrița members (Dragastan, 2010), two lithological units that are difficult to discriminate in outcrop. From the first studies concerning the Jurassic deposits in this massif, it was assumed that Oxfordian deposits were present at the base of the reef limestones, as limestones with siliceous nodules. Boldur *et al.* (1968) assigned the succession of red, marly limestones and pink limestones with layers of reddish jasper to the Oxfordian. Nevertheless, the associations described in these deposits by Dragastan (2010) show a wider stratigraphic distribution, including the Kimmeridgian–Tithonian interval. As a consequence, the presence of the Oxfordian in the white limestones succession is still unconfirmed. The Kimmeridgian–Tithonian deposits, consisting of massive reef limestones with thicknesses sometimes exceeding 300 m, dominate the calcareous succession of the Buila-Vânturarița Massif, making up the main crest. They contain a rich micropalaeontological fossil suite. Pleș *et al.* (2013) identified the following main species of micro-organisms in microbial crusts and encrusting microbial organisms: *Crescentiella morronensis* Crescenti, 1969, *Radimura cautica* Senowbari-Daryan et Schäfer, 1979, *Koskinobullina socialis* Cherchi et Schroeder, 1979, *Iberopora bodeuri* Granier et Berthou, 2002, *Bacinella*-type structures and/or *Lithocodium aggregatum* Elliott, 1956. The microbial structures and the encrusting organisms were essential for the development of these bioconstructions. Pleș *et al.* (2013) defined the latter as “coral-microbial-microencruster boundstones”. They are similar to the Upper Jurassic–lowermost Cretaceous carbonate deposits of other regions in the




Fig. 2. Location of the sections studied.

intra-Tethys domain. In this region, lowermost Cretaceous (Berriasian–Valanginian) sediments cover small areas; they are overlain transgressively by Urgonian limestones (Dragastan, 1980; Uță and Bucur, 2003). Dragastan (1980, 2010) identified the following micro-organisms as characterizing the Berriasian–Valanginian interval: foraminifera [*Mohlerina basiliensis* (Mohler, 1938), *Andersenolina elongata* (Leupold, 1935), *A. alpina* (Leupold, 1935), *A. delphinensis* (Arnaud-Vanneau, Boisseau et Darsac, 1988), *Mayncina bulgarica* ng1033 Laug, Peybernes et Rey, 1980, and *Kaminskia acuta* Neagu, 1999], calcareous algae [*Salpingoporella annulata* Carozzi, 1953, *Rajkaella iailensis* (Maslov, 1965), *Macroporella praturloni* Dragastan, 1978, *Actinoporella podolica* (Alth, 1878), *Felixporidium atanasii* Dragastan, 1999, *Fagetiella angulata* Dragastan, 1988] and micro-encrusting organisms. Dragastan (2010) considered that Hauterivian deposits also should be present within the Lower Cretaceous deposits of the Buila-Vânturarița Massif. In the opinion of the present authors, the foraminiferal assemblage including *Haplophragmoides joukowskyi* Charollais, Bronnimann et Zaninetti, 1966, *Neotrocholina* sp., *Patellina turriculata* Dieni et Massari, 1966, *Andersenolina histeri* Neagu, 1994 and *Kaminskia exigua* Neagu, 1999 cannot be considered to be typical for the Hauterivian thus, its presence in the succession leaves questions about the correctness of the stratigraphy. Uță and Bucur (2003) confirmed the presence of Barremian–?Lower Aptian in the area, based on the occurrence of the species *Vercorsella hensoni* (Dalbiez, 1958), *Vercorsella camposaurii* (Sartoni et Crescenti, 1962), *Charentia* sp., *Everticycla-mmima* sp. and *Falsolikanela danilovae* Radoičić, 1969. In the north-

eastern part of the massif, Upper Cretaceous deposits lie transgressively and unconformably on top of the Upper Jurassic–Lower Cretaceous limestones (Todiriță–Mihăilescu, 1973). Within these deposits, Boldur *et al.* (1970) recognized three individual sedimentary complexes: Vraconian–Cenomanian–Turonian, Coniacian–Santonian, and Campanian–Maastrichtian. Mesozoic deposits are covered by transgressive Eocene sedimentary rocks (Popescu, 1954).

MATERIAL AND METHODS

The authors collected 1,250 samples from profiles in eight different areas in the Buila-Vânturarița Massif: P1 – Bistrița Gorges; P2 – Arnota Mountain; P3 – Costești Gorges; P4 – Cacova sector; P5 – Albu Mountain; P6 – Ștevioara sector; P7 – Curmătura Oale, and P8 – Cheii Gorges (Fig. 2). The sampling procedure was established on the basis of local features in each of the profiles/transects. In most cases, the profiles run along valley bottoms, where the carbonate deposits are well-exposed. From these samples, the authors prepared 1,270 thin sections for petrographic analysis and the study of microfacies and diagenetic processes.

AGE CONSTRAINTS

Reef-type deposits were found in all eight profiles, making up the most significant calcareous succession in the Buila-Vânturarița Massif. The most important microfossils for biostratigraphy (Table 1) led the authors to assign a Kimmeridgian–Tithonian age to the sequence, on the basis of the association of foraminifera and calcareous algae [*Salpingoporella pygmaea* (Gümbel, 1891), *Clypeina sulcata* (Alth, 1881), *Nipponophycus ramosus* Yabe et Toyama, 1928, *Coscinoconus alpinus* Leupold, 1935, *Mohlerina basiliensis*] that are typical for Upper Jurassic carbonate-platform facies in many European regions (Dragastan, 1975; Bucur, 1978; Kołodziej and Decrouez, 1997; Carras and Georgala, 1998; Krajewski, 2000; Uță and Bucur, 2003; Olivier *et al.*, 2004; Bucur and Săsăran, 2005; Săsăran, 2006; Catincuț *et al.*, 2010; Ivanova and Kołodziej, 2010; Rusciadelli *et al.*, 2011; Turi *et al.*, 2011; Pleș *et al.*, 2013; Pleș and Schlagintweit, 2014).

The carbonate deposits overlying the reef limestones include a micropalaeontological association composed of *Coscinoconus delphinensis* (Arnaud-Vanneau, Boisseau et Darsac, 1988), *Montsalevia salevensis* (Charrolais, Brönnimann et Zaninetti, 1987), *Protopeneloplis ultragranulata* (Gorbachik, 1971), *Charentia evoluta* (Gorbachik, 1968) and *Bacinella*-type structures. To these, Dragastan (1980) added *Salpingoporella annulata*, *Rajkaella iailensis*, *Macroporella praturloni*, and *Actinoporella podolica*. The entire micropalaeontological association is typical for lowermost Cretaceous (Berriasian–Valanginian) rocks, as it also has been recognised in other areas within the facies types assigned to this stratigraphic interval, for example, in the Romanian Carpathians, the Hațeg-Pui and Brașov-Dâmbovicioara-Piatra Craiului areas, and in other areas of the Tethyan domain by Bucur (1993), Chiocchini *et al.* (1994),

Bucur *et al.* (1995), Bulot *et al.* (1997), Kołodziej and Decrouez (1997); Ivanova (1999), Moshammer and Schlagintweit (1999), Schlagintweit and Ebli (1999), Schroeder *et al.* (2000), Pop and Bucur (2001), Radoičić (2005), Dragastan (2010), Bucur *et al.* (2014a, b, c) and Mircescu *et al.* (2014).

In some areas within the Buila-Vânturarița Massif (Arnota Mountain, Costesti Gorges, Cacova Mountain, Albu Mountain, Ștevioara sector and Curmătura Oale) the authors identified limestones that transgressively overlie the Upper Jurassic–lowermost Cretaceous. In such deposits the authors have identified specimens of *Paracoskinolina? jourdanensis* Foury et Moullade, 1966 and *Vercorsella camposaurii*. Dragastan (1980), and Uță and Bucur (2003) also found *Vercorsella henoni*, *Paracoskinolina* sp., *Everticyclammina* sp., *Salpingoporella muehlbergii* (Lorenz, 1902), *Falsolikanella danilovae*, as well as rudist fragments. On the basis of this association, the deposits on top of the Upper Jurassic–lowermost Cretaceous succession were assigned by Dragastan (1980), and Uță and Bucur (2003) to the Barremian–?Lower Aptian interval. Similar associations were described elsewhere (Bucur *et al.*, 1993; Bodrogi *et al.*, 1994; Turi *et al.*, 2011; Lazăr *et al.*, 2012; Michetiuc *et al.*, 2012; or Bruchental *et al.*, 2014).

FACIES AND MICROFACIES ANALYSIS

The authors identified five major microfacies associations (MFA), three in the Upper Jurassic (MFA1, MFA2, MFA3) and two in the Lower Cretaceous (MFA4 and MFA5) deposits, respectively (Figs 3, 4).

MFA1 – Fine bioclastic grainstone/packstone

This facies was identified on Albu Mountain (P5), in the Ștevioara sector (P6) and in the Curmătura Oale profile (P7). In the Albu Mountain area (P5), MFA1 is represented mostly by fine bioclastic grainstones and packstones with small fragments of echinoids, calcified sponges, foraminifera and *Crescentiella morronensis*. In the Ștevioara sector (P6), such deposits dominate the lower half of the carbonate succession and they are represented by fine-grained, bioclastic limestones with peloids, foraminifera and *Crescentiella morronensis*. In the well-sorted deposits, the authors identified the following microfacies types: fine, bioclastic-peloidal grainstone; bioclastic packstone/grainstone with thin microbial crusts; and bioclastic packstone with small fragments of bioconstructing organisms (sclerosponges or encrusting micro-organisms). The abundance of microbial laminated crusts in these limestones is noteworthy. In the Curmătura Oale profile (P7), the fine-grained microfacies types (MFA1) also occurs in the lower part of the succession, as interlayers within coarser deposits. These limestones have similar characteristics to the ones previously described: well sorted, microbial content, fragments of corals and encrusting micro-organisms. The main MFA1 microfacies types in this profile are represented by fine, bioclastic grainstone with foraminifera, peloids or *Crescentiella*-type structures and bioclastic packstone with fragments of sclerosponges and echinoids.




Fig. 3. General succession of the limestone deposits of the Buila-Vânturarița Massif (sections P1 to P4).





Fig. 4. General succession of the limestone deposits of the Buila-Vânturarița Massif (sections P5 to P8).

Table 1

Main Upper Jurassic–Lower Cretaceous microfossils of the Buila-Vânturarița Massif and their stratigraphic range

		Biostratigraphical range			
		KIMMERIDGIAN	TITHONIAN	BERRIASIAN	VALANGINIAN
Foraminifera	<i>Charentia evoluta</i> (GORBATCHIK)		█	█	
	<i>Protopenoplis ultragranulata</i> (GORBATCHIK)		█		
	<i>Coscinoconus alpinus</i> LEUPOLD	█			
	<i>Bulloporella aff. laevis</i> SOLLAS	█			
	<i>Lituola? baculiformis</i> SCHLAGINTWEIT & GAWLICK	█			
	<i>Mohlerina basiliensis</i> (MOHLER)	█			
	<i>Troglotella incrustans</i> WERNLI & FOOKES	█			
	<i>Montsalevia salevensis</i> CHAROLLAIS, BRÖNNIMANN & ZANNINETI			█	
	<i>Thaumatoporella parvovesiculifera</i> RANIERI	█			
	<i>Salpingoporella pymaea</i> (GUEMBEL)	█			
	<i>Glypeina sulcata</i> (ALTH)	█			
Calcareous algae	<i>Nipponophycus ramosus</i> YABE & TOYAMA	█		█	█
	" <i>Solenopora</i> " sp.	█		█	█
	<i>Calcastella jachenhausenensis</i> REITNER		█		
	<i>Neuropora lusitanica</i> TERMIER & TERMIER	█			
	<i>Perturbatacrusta leini</i> SCHLAGINTWEIT & GAWLICK	█			
Sclerospoges	<i>Thalamopora lusitanica</i> TERMIER & TERMIER	█			
	<i>Crescentiella morronensis</i> (CRESCENTI)	█			
Microencrusters	<i>Radiomura cautica</i> SENOWBARI-DARYAN & SCHAEFER	█			
	<i>Koskinobullina socialis</i> CHERCHI & SCHROEDER	█			
	<i>Iberopora bodeuri</i> GRANIER & BERTHO	█			



Interpretation

Most probably, the fine-grained facies types (MFA1), identified in the three sections, are the results of moderate and concentrated density flows (turbiditic deposits). The clasts originated in the reef area and were probably deposited in the lower part of the reef slope. In some cases, they are associated with deposits showing MFA2 and MFA3 facies types. On the basis of their association with reef-slope rudstones (MFA2), these deposits can be assigned to the “slope carbonate apron” model defined by Mullins and Cook (1986). Comparable microfacies types were documented by Bucur *et al.* (2010a) in the Upper Jurassic carbonate deposits of the Mateiaş area, indicating similar depositional environments.

MFA2 – Coarse bioclastic-intraclastic grainstone/rudstone

This microfacies type is commonly recognized throughout the entire carbonate succession in the Buila-Vânturariţa Massif, in all eight sections studied. In the Bistriţa Gorges (P1) it consists mainly of reef debris levels ubiquitous throughout the entire succession, showing tabular and layer-like geometries (Pleş *et al.*, 2013). The microfacies are intra-bioclastic rudstone and coarse, intra-bioclastic grainstone. These limestones are mainly composed of coral fragments, sclerosponges and large intraclasts of coral-microbial boundstone, stromatolitic-thrombolitic microbial crusts and bioclastic packstone with *Crescentiella morronensis*. Clasts are angular to subangular and variable in size (up to 1.5 cm in diameter). The deposits are poorly sorted, with a chaotic orientation of clasts. These limestones contain a micropalaeontological association of foraminifera [*Charentia evoluta*, *Lituola? baculiformis* Schlagintweit et Gawlick, 2009, *Mohlerina basiliensis* (Fig. 5B), *Coscinoconus alpinus*, *C. delphinensis* (Fig. 5A), *Troglotella incrustans* Wernli et Fookes, 1992, *Bullopore laevis* Sollas, 1877, *Lenticulina* sp. (Fig. 5C), *Coscinophragma* sp. (Fig. 5D), *Ammobaculites* sp.], calcareous algae [*Clypeina sulcata* (Fig. 5J), *Salpingoporella pygmaea* (Fig. 5H), *Thaumatoporella parvovesiculifera* (Ranieri, 1927), *Nipponophycus ramosus* (Fig. 5I), “*Solenopora*” sp.), sclerosponges (*Perturbatacrusta leini* Schlagintweit et Gawlick, 2011, *Murania reitneri* Schlagintweit, 2004, *Neuropora lusitanica* Termier et Termier, 1985, *Thalamopora lusitanica* Termier et Termier, 1985 (Fig. 5K), *Calcistella jachenhausenensis* Reitner, 1992), gastropods, bivalves and worm tubes (*Mercierella dacica* Dragastan, 1967 and *Terebella* sp.). In the Arnota Mountain profile (P2), the authors identi-

fied the following microfacies types: intra-bioclastic floatstone, bio-intraclastic rudstone, and less frequently, bioclastic grainstone. In the Costeşti Gorges (P3), rudstone levels (MFA2) are very thick; they were mainly observed in the lower and middle part of the profile. Facies assigned to the MFA2 type dominate the upper half of the Cacova sector succession (P4) and Albu Mountain (P5), consisting of coarse grainstone and bioclastic rudstone with fragments of reef macro-organisms. In the Ştevioara (P6) and Curmătura Oale (P7) sectors, the coarse deposits (MFA2) are found in the middle part of the succession. The main biota present in these microfacies types consist of organisms related to the reef, represented by fragments of corals and sclerosponges, microproblematic organisms (*Crescentiella morronensis*, *Lithocodium aggregatum* or *Bacinella*-type micro-organisms), benthic foraminifera, as well as some molluscs (gastropods or fragments of bivalves). In the lower and middle part of the Cheia Valley succession (P8), coarse deposits of MFA2 dominate.

Interpretation

The MFA2-type deposits are significantly thick and they consist of poorly-sorted angular clasts represented by intraclasts and bioclastic fragments. Constant wave action and currents caused erosion during reef growth, so many fragments were redeposited and incorporated within these rudstones (Turnšek *et al.*, 1981). Resedimentation of the reef clasts can be linked to small debris flows in an upper slope environment (Morsilli and Bosellini, 1997). The encrusting nature of many of the micro-organisms, associated with syndepositional cement crusts, have stabilized and reinforced these slope carbonates. The reef debris deposits (MFA2) in association with coral-stromatoporoid-microencruster boundstones (MFA3) are interpreted as an equivalent of the fore-reef/upper-slope “Plassen”-type limestones, described by Schlagintweit and Gawlick (2008) from the Northern Calcareous Alps. They also resemble the platform-margin *Ellipsactinia* facies of the Central Apennines (Rusciadelli *et al.*, 2011), characterized by similar sedimentary settings. Other close facies associations have been described from several areas of the Romanian Carpathians (Bucur, 1978; Bucur and Săsăran, 2005; Bucur *et al.*, 2010a, b; Bucur and Săsăran, 2012).

MFA3 – Coral-stromatoporoid-microencruster boundstone

In the Bistriţa Gorges profile (P1), boundstone levels (MFA3) occur as interlayers in the lower and upper part of

Fig. 5. Main Upper Jurassic–Lower Cretaceous microfossils of the Buila-Vânturariţa Massif. **A.** *Coscinoconus delphinensis* (sample 794 – Valea Cheii Gorges – P8). **B.** *Mohlerina basiliensis* (sample 1053 – Ştevioara sector – P6). **C.** *Lenticulina* sp. (sample 38 – Bistriţa Gorges – P1). **D.** *Coscinophragma* sp. (sample 1040 – Ştevioara sector – P6). **E.** *Protopenneroplis ultragranulata* (sample 879 – Curmătura Oale – P7). **F.** *Vercorsella camposaurii* (sample 890 – Curmătura Oale – P7). **G.** *Paracoskinolina? jourdanensis* (sample 457 – Costeşti Gorges – P3). **H.** *Salpingoporella pygmaea* (sample 1054 – Valea Cheii Gorges – P8). **I.** *Nipponophycus ramosus* (sample 472 – Costeşti Gorges – P3). **J.** *Clypeina sulcata* (sample 1054 – Ştevioara sector – P6). **K.** *Thalamopora lusitanica* (sample 19b – Bistriţa Gorges – P1). **L.** *Ellipsactinia* sp. (sample 114 – Bistriţa Gorges – P1).

the succession, on top of the intraclastic-bioclastic rudstone/grainstone facies types (MFA2). The corals and sponges are intensely encrusted by problematic micro-organisms (*Crescentiella morronensis*, *Radiomura cautica*, *Koskinobullina socialis*, *Lithocodium aggregatum*, *Iberopora bodeuri*, or *Bacinella*-type structures). Additionally, the authors identified stromatolitic- and thrombolitic-type structures. Peloids, bioclasts and silt-sized carbonate intraclasts embedded in stromatolitic crusts were recognized. Other organisms are represented by stromatoporoids (*Ellipsactinia* sp., *Calci-stella jachenhausenensis*), ?udoteacean algae (*Nipponophycus ramosus*), dasycladales (*Salpingoporella pygmaea*), benthic foraminifera (*Lituola? baculiformis*, *Mohlerina basiliensis*, *Coscinoconus alpinus*, *Troglotella incrustans*, *Lenticulina* sp., *Coscinophragma* sp.), molluscs, echinoid fragments and spines, or worm tubes (including *Terebella* sp.). An important feature of the bioconstructed limestones from Bistrița Gorges is represented by the presence of *Epiphyton*-type micro-organisms. These were recently identified for the first time in the Upper Jurassic limestones (Kimmeridgian–Tithonian) in Romania (Săsară et al., 2014). In the Arnota Mountain profile (P2) MFA3 is less frequent. The main micro-encrusters involved in the development and consolidation of this deposit are *Lithocodium aggregatum* and *Crescentiella morronensis*. MFA3-type boundstones were mainly observed in the lower half of the Costești Gorges section (P3), interlayered with MFA2-type deposits. They consist of coral-microencruster boundstones with internal wackestone/packstone sediment and boundstones with stromatoporoids and encrusting organisms. In the Cacova sector (P4) the boundstone levels (MFA3) are thin and were observed only at the base and the top of the succession. The most representative facies types consist of coral-microbial boundstones with thrombolitic crusts and stromatoporoids and brecciated boundstones with problematic micro-organisms (*Crescentiella morronensis*). In the Ștevioara profile (P6), a horizon of coral-microbial boundstone with stromatoporoids (MFA3) conformably overlies the fine-grained deposits (MFA1). Here, the following microfacies types occur: coral-microbial boundstone and boundstone with stromatoporoids (*Ellipsactinia* sp., *Calci-stella jachenhausenensis*, *Neuropora lusitanica* or *Actinostromaria* sp.) and micro-encrusters. MFA3 is present along the entire succession in the Curmătura Oale profile (P7), as interlayers within bio-intraclastic rudstones (MFA2) and micritic fenestral deposits (belonging to MFA4). Boundstones dominate the lower part of the profile as banks of various thicknesses. The dominant microfacies type is coral-microbial framestone and boundstone with stromatoporoids and microbial crusts. Only sporadically, leiolitic or stromatolitic structures were noticed. In the Cheii Valley profile (P8), the deposits that the authors assigned to MFA3 occur solely at the end of the lower part of the profile. They are represented by coral-microbial and pure, microbial boundstone. Microscopically, as opposed to most of the MFA2- and MFA3-type limestone samples investigated in the region, thin sections from Cheii Valley (P8) show lesser amounts of microbial crusts. Crustiform fabrics associated with various encrusting micro-organisms are rare, while boundstones with stromatoporoids are absent. The microbial structures are repre-

sented by leiolitic microstructures, as well as fine-laminated mesostructures.

Interpretation

The bioconstruction levels (MFA3) from the Buila-Vânturarița Massif are generally characterized by crustose and encrusting fabrics, associated with widely developed cement crusts. The intrareef sediment of MFA3-type deposits is mud-dominated, represented by bioclastic wackestone with various reef organisms, such as fragments of corals and sclerosponges, foraminifera, calcareous algae (*Salpingoporella pygmaea* and/or *Nipponophycus ramosus*), as well as encrusting micro-organisms or clotted microbial structures. The micro- and macropalaeontological assemblages from the MFA3 deposits, represent a typical frontal reef or reef-crest association, as previously documented by Morsilli and Bosellini (1997), Schlagintweit and Gawlick (2008) or Pleș et al. (2013). The abundance of calcareous sponges [*Perturbatacrusta leini*, *Ellipsactinia* sp. (Fig. 5L), *Neuropora lusitanica*, *Thalamopora lusitanica* or *Cylicopsis verticalis* Turnšek, 1968], makes possible the correlation of the analysed boundstones (MFA3) with bioconstructions of the stromatoporoid zones from many Late Jurassic barrier-reef models of the intra-Tethys domain. Thus, the MFA3 microfacies association may correspond to the “Stromatopores Unit” of Rusciadelli et al., (2011) from the external zone (upper slope-reef crest transition) of the Marsica area reef (Central Apennines), characterized by abundant stromatoporoids (Ellipsactinia Limestones). MFA3-type deposits also resemble facies associations 4 and 5 (boundstones and coarse grainstones with *Ellipsactinia* or *Sphaeractinia*) of Morsilli and Bosellini (1997), and with the *Actinostromaria* Zone described by Turnšek et al. (1981) from Slovenia.

MFA4 – Peritidal bioclastic-oncoidal-peloidal wackestone

This microfacies association was assigned to the lowermost Cretaceous. They have been identified only in the profiles in the Arnota Mountain (P2), Cacova (P4), Ștevioara (P6), Curmătura Oale (P7) sectors, and Cheia Valley (P8), where they are developed conformably on top of the Jurassic reef limestones. For example, in the profile at Arnota Mountain (P2), the peloidal-bioclastic wackestones and the fenestral-microbial bindstones represent the main microfacies (MFA4) the top of the profile. In the Cacova (P4) and Ștevioara (P6) sectors, MFA4 includes the following microfacies: peloidal wackestones with fenestrae, bioclastic-peloidal wackestones with micritized rivulariacean-type cyanobacteria (*Cayeuxia* sp.), peloidal-intraclastic wackestones, fenestral wackestones, and bioclastic-peloidal packstones. Foraminifera [*Protopenneroplis ultragranulata* (Fig. 5E) or *Montsalevia salevensis*], cyanobacteria, crustacean coprolites (*Favreina* sp.) and *Bacinella*-type structures represent the main biotic components. In the upper part of the Curmătura Oale profile (P7) the authors identified microbial limestones with fenestrae, which we also assigned to the MFA4. These are associated with (or, towards the top gradually pass into) bioclastic fenestral wackestone, peloidal bioclastic wackestone/packstone and non-fossiliferous mud-












Fig. 6. Distribution of the main facies associations within the carbonate succession of the Buila-Vânturarița Massif. **A.** Fine bioclastic-peloidal grainstone with *Crescentiella morronensis* (MFA2) (sample 778 – Albu Mountain – P5). **B.** Coarse intraclastic-bioclastic rudstone with microbial structures and echinoid fragments (MFA3) (sample 489 – Costești Gorges – P3). **C.** Stromatopora-microencruster boundstone with *Neuropora lusitanica*, *Perturbatacrusta leini* and *Crescentiella*-type structures (MFA4); note the distinct micritic envelopes surrounding the cyanobacteria fragments (sample 548 – Costești Gorges – P3). **D.** Bioclastic floatstone with rivulariacean-type cyanobacteria oncoids (MFA4); note the distinct micritic envelopes surrounding the cyanobacteria fragments (sample 548 – Costești Gorges – P3). **E.** Bioclastic packstone with foraminifera (MFA5) (sample 889 – Curmătura Oale – P7).



stone. In the carbonate succession from Cheii Valley, the following microfacies occur: peloidal-bioclastic wackestone with cyanobacteria and micritic envelopes; bioclastic wackestone with fractures and cement-filled cavities; fractured mudstone, and brecciated peloidal-intraclastic wackestone. Here, the biotic component is represented by foraminifera, microbialites, as well as species of green algae (*Clypeina sulcata*).

Interpretation

The facies types assigned to MFA4 are present in the upper part of the profiles studied, being deposited in peritidal environments with reduced hydrodynamics (Shinn, 1983; Hardie and Shinn, 1986). These limestones are frequently associated with the facies of intertidal ponds and in places with supratidal deposits. Overall, they most probably characterize lagoonal sedimentary settings (Morsilli and Bosellini, 1997). The abundant rivularia-type cyanobacteria (often with clear envelopes or fine micritic cortexes), microbial oncoids and fenestral structures indicate a transition with more protected depositional environments of these micrite-dominated limestones (Flügel, 2004).

MFA5 – Bioclastic packstone/wackestone

This microfacies type is found in six out of the eight profiles studied. They transgressively overlie Upper Jurassic–lowermost Cretaceous deposits (MFA1, MFA2, MFA3). In the Arnota Mountain (P2) and the Costești Gorges (P3), the Lower Cretaceous deposits (MFA5) are similar to those of MFA4-type, at the tops of the successions. Microfacies types are represented by bioclastic-peloidal wackestone with foraminifera [*Paracoskinolina? jourdanensis* (Fig. 5G) and *Vercorsella camposaurii* (Fig. 5F)], fractured, intraclastic wackestone, bioclastic-oncoidal wackestone with rivularia-type cyanobacteria and peloidal-fenestral wackestone. In the Cacova sector (P4), the authors have identified bioclastic-peloidal wackestone/packstone with foraminifera (*Paracoskinolina? jourdanensis*) and rudist fragments, peloidal fenestral wackestone and fractured wackestone. The

deposits assigned to MFA5 crop out in small areas on top of the successions in Albu Mountain (P5), Ștevioara (P6) and Curmătura Oale (P7) sectors. Here, fossils are relatively scarce, the main bioclasts being represented by benthic foraminifera. Microfacies types are: bioclastic peloidal wackestone with foraminifera, bioclastic microbial wackestone and fractured bioclastic packstone.






Interpretation

The carbonate deposits assigned to MFA5 formed in a shallow, subtidal environment of the platform shelf. Taxonomic diversity of microfossils is significantly reduced as compared to the previously described deposits (MFA1 to MFA4). Micrite-dominated facies types, characterized by micritization processes, were noticed towards the top of this microfacies association type.

FACIES DISTRIBUTION, SEDIMENTARY SETTINGS AND EVOLUTION OF STUDIED CARBONATE PLATFORM

Starting with the lower part of the carbonate succession of the Buila-Vânturarița Massif, finer deposits of granular flows with turbiditic character (MFA1) occur. They were recognized in three of the sections analysed (Fig. 6). This feature points to an environment with relatively high energy, most probably located towards the base of the reef slope (Mullins and Cook, 1986; Bucur *et al.*, 2010a). On top of these deposits, thick levels of reef rudstones (MFA2), interlayered with coral-microencruster boundstones with stromatoporoids (MFA3), are present. Bioconstructions (MFA3) occur on top of MFA2-type facies that evidence the instability of the reefal slope. The associated distribution of reef facies types (MFA2 and MFA3), the thickness of the rudstone levels and the presence of typical microfossil associations are all arguments for a patch-reef development of most of the bioconstructions in Buila-Vânturarița Massif. The genesis of these bioconstructions was strictly controlled

Fig. 7. Diagenetic features. **A.** Large cement crust formed as a result of radial-fibrous cement recrystallization (RFC) in a reef-slope rudstone; note fine laminated peloidal sediment in the cavity formed by cement growth indicating syndepositional cementation (sample 21 - Bistriței Gorges - P1). **B.** Neomorphic features inside a coral fragment; after dissolution of the septa the interseptal spaces were subsequently filled with granular cement (arrow) (sample 1050 - Ștevioara sector - P6). **C.** Syndepositional radial-fibrous cement (RFC) succeeded by a recrystallization phase; notice the relic features of the primary radial-fibrous cementation (arrows) followed by a granular spar cement coarsening towards the centre of the crust; CM - *Crescentiella morronensis* encrusting syndepositional cement (sample 14b - Bistriței Gorges - P1). **D.** Multiple cement generations in a reef-slope facies; the first generation of cement is represented by a fine rim of isopachous-fibrous cement (IFC) surrounding the oncoids, followed by a brownish microcrystalline cement (MC); the third generation of cement is composed of bladed isopachous spar crystals (BS) succeeded by a final stage of clear blocky calcite cementation (BC) (sample 696 - Albu Mountain - P5). **E.** Meteoric dissolution resulted in cavities (DC) bordered by rims of non-isopachous dog-tooth cement (NIDC), locally associated with syntaxial cement overgrowing echinoid fragments (SOC); the internal filling of the cavities is made up of fine, marl-silty sediment, probably of vadose origin (sample 151 - Bistriței Gorges - P1). **F.** Mixed meteoric and shallow burial features in a coral-microbial boundstone; the meteoric diagenesis is represented by several dissolution cavities (DC); the corals were partially fractured during initial burial, followed by a last stage of cementation inside the fissures, represented by clear blocky crystals (sample 13a - Bistriței Gorges - P1). **G.** Peritidal oncoidal wackestone with gravitational cements (GC) that hang from the bottoms of the grains; the remaining pores were filled with granular spar (sample 332 - Arnota Mountain - P2). **H.** Early dolomitization; some of the rhombohedral dolomite crystals are partially filled with vadose calcite (dedolomitization), probably triggered by the change of meteoric regime into a normal-marine one (sample 943 - Curmătura Oale - P7).



by stabilization of the reef debris. Thus, the encrusting character or the sediment-binding function that most bioconstructing organisms play. The relatively reduced thickness of these bioconstructed levels (MFA3) may be also connected to the nature of the reef framework-forming organisms. In contrast to the Upper Jurassic microbialite-dominated reefs of the northern Tethyan realm (Leinfelder *et al.*, 1993), the cement-supported microencruster frameworks, which characterize the MFA2 and MFA3 facies associations, show many similarities with the Late Jurassic “Plassen”-type microencruster boundstones of the Northern Calcareous Alps (Schlagintweit and Gawlick, 2008; Pleş *et al.*, 2013). Therefore, laminated microbial structures are subordinate to encrusting micro-organisms associated with syndimentary cement crusts in the development and consolidation of these deposits. On the basis of the composition and textural-structural features, the described facies associations (MFA2 and MFA3) were assigned to fore-reef slope environments, close to the proximal areas of the reef crest (Turnšek *et al.*, 1981; Schlagintweit and Gawlick, 2008; Rusciadelli *et al.*, 2011; Pleş *et al.*, 2013). The presence of stromatoporoids in these facies was supported by their location in shallow, high-energy environments. This also explains the resistance of most of these taxa to remobilization or fragmentation, which are typical phenomena in platform-marginal environments (Leinfelder *et al.*, 2005). In the upper part of the Upper Jurassic–lowermost Cretaceous succession, all the typical terms of peritidal environments (subtidal, intertidal and supratidal – MFA4) were recorded. The subtidal, marine environment is indicated by the type of carbonate components (bioclasts, peloids, oncoids), embedded in a micritic matrix associated sometimes with bioturbation structures. The intertidal environment includes beach deposits, formed in areas with high energy (peloidal packstones with “keystone vugs”) and fine, laminated fenestral deposits, typical for a low-energy environment. The supratidal environment contains non-fossiliferous and fenestral, micritic limestones that might have been affected by dolomitization. These deposits are comparable to the peritidal cycles of the lagoonal facies association (F1A, F1D and F1E), documented by Morsilli and Bosellini (1997) in Upper Jurassic–Lower Cretaceous platform-margin limestones in Southern Italy. Peritidal limestones of the Buila-Vânturarița Massif show a decrease of accommodation space in the carbonate platform. The Lower Cretaceous deposits assigned to the MFA5 (Bar-

remian) transgressively overlie the Jurassic–lowermost Cretaceous succession (Fig. 6). Towards the top of the Cretaceous succession the authors observed a more internal evolutionary trend of the depositional environment, as indicated by fracture fillings, bioclast micritization and oncoids.

The general distribution and features of the main facies associations, together with the associated biota, indicate a barrier-type development of the Buila-Vânturarița carbonate platform (Turnšek *et al.*, 1981; Schlagintweit and Gawlick, 2008; Rusciadelli *et al.*, 2011). Furthermore, the Buila-Vânturarița platform slope shows the classical geometry of many intra-Tethyan platform margins (Morsilli and Bosellini, 1997; Pleş *et al.*, 2013). The gradual transition from deeper to shallower sedimentary settings can be strongly correlated with the prograding character of the carbonate platform during the Upper Jurassic.

DIAGENETIC HISTORY

Early marine diagenesis in subtidal environments

The features identified in the eight profiles studied that are related to early marine diagenesis consist of: (i) successive generations of cements with fibrous fabrics, (ii) recrystallization, and (iii) micritization. Fibrous-radiaxial, microcrystalline and scalenohedral cements were frequently noticed in reef deposits (MFA2 and MFA3). Fibrous radiaxial cements may form crusts inside interparticle spaces or vugs, in most cases being associated with biogenic encrustations. Sometimes these cements can be seen as replacements for fast, early cementation products. This mechanism is supported by the presence of relic, prismatic crystals (possibly of aragonitic nature) within the crusts (Fig. 7A, C; Reitner, 1986). Morphological traits of the relic crystals are preserved (Kendall and Tucker, 1973; Henrich and Zankl, 1986; Koch and Schorr, 1986), or alternatively the replacement process may be traced by the presence of obvious discontinuities within the cements crusts formed in the cavities of the reef deposits (MFA2 and MFA3). The syndimentary nature of these cement crusts is reflected in the encrustation by micro-encrusters (Fig. 8A; Schlagintweit and Gawlick, 2008; Kołodziej, 2015b). The scalenohedral cements are mainly composed of denticulated, prismatic cements, developed as infilling crusts within intra-reef cavi-

Fig. 8. Microfacies and diagenetic features. **A.** Crusts of radiaxial-fibrous cement (RFC) strengthening a stromatopoid-microencruster framework with *Cylicopsis verticalis* (C), *Tubuliella fluegeli* Turnšek (T) and *Perturbatacrusta leini* (P) (sample 737 – Albu Mountain – P5). **B.** Scalenohedral cement (SC) formed in a reef cavity; note the presence of a boring structure (BS) inside the zoantharian fragment (sample 1109 – Arnota Mountain – P2). **C.** Stromatolitic-clotted meso-structure supporting the stromatopoid boundstone (sample 554 – Costești Gorges – P3). **D.** Dissolution cavity (DC) filled with fine silty or marly sediment; the external margins are bordered by a clear layer of non-isopachous dog-tooth crystals (NIDC) associated with syntaxial-overgrowth cement (SOC) (sample 1118 – Arnota Mountain – P2). **E.** Successive microcrystalline crusts probably induced by cyanobacterial growth; note the radial displacement of crystals associated with fine, micritic lens; in the upper part of the picture, there is a small cavity filled with geopetal sediment (arrow); in the lower part of the cavity a second generation of non-isopachous dog-tooth cement developed on an early generation of isopachous marine cement (sample 1130 – Arnota Mountain – P2). **F.** Multiple generations of cements in a closed pore; the chronological order of cementation is evidenced by cathodoluminescence; the first cementation stage is represented by non-luminescent, zoned crystal rims, followed by dull-luminescent much larger crystals; the final filling of the pore is marked by bright orange, blocky cementation probably formed during shallow burial (sample 5 – Bistriței Gorges – P1).

ties (Fig. 8B). In the peritidal facies types (MFA4), some crusts of microcrystalline cements are represented by very fine, micritic laminae intercalated with radial, microsparitic cements (Fig. 8E). The relatively small size of crystals in the microcrystalline filaments indicates relatively rapid cementation processes, probably induced by changes in the chemistry of the environments, where cyanobacteria were present. Heinrich and Zankl (1986) have described association of cyanobacteria with sparitic, microcrystalline cements as organic-diagenetic-type cements. Microbial micritization occurs in sponge-coral bioconstructions (MFA3) and particularly in peritidal facies (MFA4). This characteristic may be used as a proxy for incipient diagenetic processes (Qureshi *et al.*, 2008; Lazăr *et al.*, 2013). These processes consist of the formation of micritic microcrystalline cortices around some bioclasts (partial micritization); *in extremis*, the bioclasts can be completely replaced by micrite (complete micritization; Flügel, 2004). In the Jurassic reef deposits studied (MFA2 and MFA3), the authors have identified neomorphic processes represented by selective (partial or total) dissolution of the coral structure, or of some stromatoporoid skeletons. This was followed by low-Mg calcite recrystallization (precipitation) within the newly-created inter-skeletal space (Fig. 7B).

Diagenesis in meteoric environments

Selective dissolution, gravitational isopachous and syntaxial cements and microkarst associated with marly-fine siliciclastic fillings, are all proof of meteoric diagenesis in the limestones in the Buila-Vânturarița Massif. In the reef environments (MFA2 and MFA3), as well as in the intertidal facies (MFA4), gravitational cementation is represented by microstalactitic cements, formed on bioclasts or intraclasts (Fig. 7G; Scholle and Ulmer-Scholle, 2003). The most common cements that the authors have associated with processes in the phreatic zone are the bladed and dog-tooth types. They consist of crystalline bands, bordering cavities in reef deposits (Figs 7E, 8B). The crystals are oriented sub-perpendicular to their substrate. Another meteoric diagenesis-related product is the overgrowth of syntaxial cement formed on the echinoderm radioles and/or plates (Figs 7E, 8D), identified in reef detrital deposits (Pomoni-Papaioannou *et al.*, 1989; Flügel and Koch, 1995; Qureshi *et al.*, 2008). In the case of the samples in the present study, the syntaxial cements are often associated with the bladed or dog-tooth cements along the borders of cavities formed as a result of meteoric dissolution. Koch and Schorr (1986) considered the association of these cements as possible evidence for subaerial episodes. According to Buddemeier and Oberdorfer (1986), dolomitization is favoured by a certain degree of mixing between marine and meteoric fluids. The present authors consider that this may explain the fact that dolomitization was observed in their samples only in the peritidal facies (MFA4), on the top of the Jurassic succession. The results are rhombohedral crystals dispersed in the microcrystalline matrix, associated with late-generation cements (Fig. 7H). In the same peritidal facies the authors also have observed partial dissolution of dolomite crystals (de-dolomitization). De-dolomitization may point to a migration

of the mixing zone towards the basin, in connection with falls in sea level (Frykman, 1986).

Shallow burial diagenesis

In their material, the authors noticed a low compaction of sediments, given the fact that early cementation took place during marine diagenesis. The fragmentation of some macrofossils (corals, bivalves or gastropods), and the presence of fissures and cavities filled with sparitic calcite are the most typical features of burial diagenesis in the limestones of the Buila-Vânturarița Massif. In the reef deposits (MFA1, MFA2 and MFA3), the burial diagenesis-related cements show typical morphological features for burial environments. The crystals are mainly equigranular (mosaic), but in some places there is an increase in size towards the centre of the pores (druse-type cement) (Fig. 8F). Such late stages may be followed by a final, blocky-type cementation. In the peritidal facies (MFA4), the final cavity-filling products are represented by mosaic-type granular cements.

Porosity

The intense microbial activity and the encrusting character of most of the biota identified in the reef limestones of the Buila-Vânturarița Massif (MFA3) resulted in a significant decrease of primary porosity and in the development of morphologies, typical for bioconstructions. In the reef-slope deposits (MFA1 and MFA2) showing coarser fabric as a result of reef detrital flows, the primary porosity is relatively higher. Secondary porosity was clearly noticed in some samples, in spite of the fact that most of the cavities had been rapidly filled with early marine cements. The authors assume that, following the early burial, selective dissolution by meteoric waters led to increased porosity in the form of fissures and cavities (Heinrich and Zankl, 1986). These voids were subsequently filled in part by sparitic microcrystalline crusts or bands of inequigranular crystals (Fig. 7E). Kerans *et al.* (1986) stated that porosity shows particular features on the basis of the depositional environments. The highest values are registered in areas with constant syndimentary activity, such as reef-slope deposits (MFA1 and MFA2), or peritidal facies (MFA4) at the top of the carbonate succession.

CONCLUSIONS

1. Carbonate deposits of the Buila-Vânturarița Massif, studied here in eight profiles, mainly consist of massive, white carbonates, assigned to the Upper Jurassic–Lower Cretaceous interval. The lower and middle part of the succession is characterized by thick levels of reef debris inter-layered with coral-microbial bioconstructions with encrusting micro-organisms and stromatoporoids. Towards the top of the succession, peritidal limestones are assigned to the lowermost Cretaceous. The Barremian–?Lower Aptian deposits, which cover relatively small areas, spread over the massif, transgressively overlie the white limestones of Kimmeridgian–Valanginian age.

2. The assemblages of calcareous algae and benthic foraminifera allowed the authors to specify the age of the deposits studied. Reef limestones were assigned to the Kimmeridgian–Tithonian; peritidal facies to the Berriasian–Valanginian (*Montsalevia salevensis* and *Protopenneroplis ultragranulata* being the main supporting evidence), while limestones at the top of the succession, which transgressively overlie the Upper Jurassic–lowermost Cretaceous deposits, are referable to the Barremian–?Lower Aptian (with *Paracoskinolina? jourdanensis*).

3. The facies analysis allowed us to discriminate five major facies associations: MFA1 – fine bioclastic grainstone/packstone, MFA2 – coarse bioclastic-intraclastic grainstone/rudstone, MFA3 – coral-stromatoporoid-microencruster boundstone, MFA4 – peritidal bioclastic-oncoidal-fenestral wackestone, and MFA5 – bioclastic packstone/wackestone. The first three associations are typical for the Upper Jurassic, the latter being more frequent at the top of the Jurassic, or in the Lower Cretaceous part of the sections.

4. The distribution of the facies associations within the carbonate successions studied shows the evolution of the Late Jurassic–Early Cretaceous depositional environments: from slope and reef -front to internal-platform sedimentary settings, including peritidal environments in the lowermost Cretaceous (Berriasian–Valanginian) deposits. This succession is typical for platform progradation, as recently was documented in other areas of development for Upper Jurassic–Lower Cretaceous limestones in the Getic domain (e.g., Piatra Craiului-Dâmbovicioara area).

5. Investigation of diagenetic features of the Buila-Vânturarița Massif reveals three stages. The first stage is represented by syndepositional, marine cementation that strongly influenced the features of the reef deposits, either during their formation or during early burial. Successive laminae of radiaxial and microcrystalline fibrous cements have consolidated the reef framework and have strengthened the reef-slope and peritidal deposits. The second diagenetic stage is represented by dissolution, recrystallization and precipitation of specific cement types, as a result of the infiltration of meteoric waters. Processes of dolomitization and de-dolomitization took place sporadically, triggered by the mixing of marine and meteoric waters, probably connected with transgression episodes. Given the frequent marine and meteoric cementation stages, the deposits are poorly compacted. The third stage was the formation of fractures and late-generation cements, which took place in the burial environments.

Acknowledgements

This work was made possible by the financial support of the Sectoral Operational Programme for Human Resources Development 2007–2013, co-financed by the European Social Fund, under project number POSDRU/159/1.5/ S/132400 with the title “Successful young researchers – Professional development in interdisciplinary and international context”. We would like to thank Cristian Victor Mircescu for assistance in the field. We also thank the two reviewers, Felix Schlagintweit and Bogusław Kołodziej for their remarks, which helped to improve the paper. Iuliana Lazăr and Mihaela Grădinaru (University of Bucharest) are thanked for their support with cathodoluminescence analysis. We also thank

Stephen Kershaw for correcting the English of the manuscript. Last but not least, the first author gratefully acknowledge the suggestions of Michał Gradziński (Kraków).

REFERENCES

- Bodrogi, I., Bona J. & Lobitzer, H., 1994. Vergleichende Untersuchung der Foraminiferen- und Kalkalgen-Assoziationen der Urgon-Entwicklung des Schrattekalks in Voralberg (Österreich) und der Nagyarsány Kalkstein Formation des Villány-Gebirges (Ungarn). In: Lobitzer, H. & Csaszar, G. (eds), *Jubiläumsschrift 20 Jahre Geologische Zusammenarbeit. Österreich-Ungarn*. Geologische Bundesanstalt, Wien, pp. 225–283.
- Boldur, C., Iavorschi, M. & Braboveanu, D., 1968. Câteva date noi privind stratigrafia Jurasicului din culmea Vânturarița (Carpații Meridionali). *Dări de seamă ale Institutului Geologic al României*, 54: 217–221. [In Romanian.]
- Boldur, C., Stillă, A., Iavorschi, M. & Dumitru, I., 1970. Date noi în cunoașterea stratigrafiei și tectonicii sedimentarului mezozoic de la nord de Olănești (Carpații Meridionali). *Dări de seamă ale Institutului Geologic al României*, 55: 5–23. [In Romanian.]
- Bruchental, C., Bucur, I. I., Cociuba, I. & Hebrîștean, A. M., 2014. Sea-level changes and sedimentary response across the Barremian-Early Aptian in Pădurea Craiului Mountains, Romania. *Acta Palaeontologica Romaniae*, 9: 33–46.
- Bucur, I. I., 1978. Microfaciesurile calcarelor albe din partea nordică a masivului Piatra Craiului. Considerații biostratigrafice. *Dări de seamă ale Institutului Geologic al României*, 64: 89–105. [In Romanian.]
- Bucur, I. I., 1993. Les représentants du genre *Protopenneroplis* Weynschenk dans les dépôts du Crétacé inférieur de la zone de Reșița-Moldova Nouă (Carpathes Méridionales, Roumanie). *Revue de Micropaléontologie*, 36: 213–223.
- Bucur, I. I., Beles, D., Săsăran, E. & Balica, C., 2010a. New data on facies development and micropaleontology of the eastern margin of the Getic Carbonate Platform (South Carpathians, Romania): case study of Mateiaș Limestone, *Studia Universitatis Babeș-Bolyai Geologia*, 55: 33–41.
- Bucur, I. I., Cociuba, I. & Cociuba, M., 1993. Microfacies and microfossils in the Upper Jurassic-Lower Cretaceous limestone in the southern part of the Pădurea Craiului Mountains. *Romanian Journal of Stratigraphy*, 75: 33–40.
- Bucur, I. I., Conrad, M. A. & Radoičić, R., 1995. Foraminifers and calcareous algae from the Valanginian limestones in the Jerma River Canyon, Eastern Serbia. *Revue de Paléobiologie*, 14: 349–377.
- Bucur, I. I., Grădinaru, E., Lazar, I. & Grădinaru, M., 2014b. Early Cretaceous micropaleontological assemblages from a condensed section of the Codlea Area (Southern Carpathians, Romania). *Acta Palaeontologica Romaniae*, 9: 65–82.
- Bucur, I. I., Granier, B. & Krajewski, M., 2014a. Calcareous algae, microbial structures and micropaleontological assemblages from Upper Jurassic-Lowermost Cretaceous limestones of southern Crimea. *Acta Palaeontologica Romaniae*, 10: 61–86.
- Bucur, I. I., Păcurariu, A., Săsăran, E., Filipescu, S. & Filipescu, R., 2014c. First record of lowermost Cretaceous shallow-water limestones in the basement of the Transylvanian Depression (Romania). *Carnets de Géologie*, 14: 199–210.
- Bucur, I. I. & Săsăran, E., 2005. Micropaleontological assemblages from the Upper Jurassic-Lower Cretaceous deposits of Trascău Mountains and their biostratigraphic significance. *Acta Palaeontologica Romaniae*, 5: 27–38.

- Bucur, I. I. & Săsăran, E., 2012. Large dasycladalean algae from Upper Jurassic limestone deposits of the Apuseni Mountains (Romania) – habitat and depositional environment. *Geodiversitas* 34: 219–239.
- Bucur, I. I., Săsăran, E., Balica, C., Beleş, D., Bruchental, C., Chendeș, C., Chendeș, O., Hosu, A., Lazăr, D. F., Lăpădat, A., Marian, A. V., Mircescu, C., Turi, V. & Ungureanu, R., 2010b. *Mesozoic Carbonate Deposits from Some Areas of the Romanian Carpathians – Case Studies*. Presa Universitară Clujeană, Cluj Napoca, 203 pp.
- Buddemeier, R. W. & Oberdorfer, J. A., 1986. Internal hydrology and geochemistry of coral reefs and atoll islands: key to diagenetic variations. In: Schroeder, J. H. & Purser, B. H. (eds), *Reef Diagenesis*. Springer, Berlin, pp. 91–111.
- Bulot, L. G., Masse, J. P., Moutier, L. & Virgone, A., 1997. Organisation stratigraphique et dynamique du Valanginien au passage plate-forme/bassin en Basse-Provence (Sud-Est de la France). *Bulletin de la Société Géologique de France*, 168: 171–179.
- Carras, N. & Georgala, D., 1998. Upper Jurassic to Lower Cretaceous Carbonate Facies of African Affinities in a Peri-European Area: Chalkidiki Peninsula, Greece. *Facies*, 38: 153–164.
- Catincuț, C., Michetiuc, M. & Bucur, I. I., 2010. Microfacies and microfossils of the Upper Tithonian-Lower Berriasian calcareous klippe from Ampoiței Valley (West of Alba Iulia, Romania). *Acta Palaeontologica Romaniae*, 7: 77–86.
- Chiocchini, M., Farinacci, A., Mancinelli, A., Molinari, V. & Potetti, M., 1994. Biostratigrafia a foraminiferi, dasicladali e calpionelle delle successioni carbonatiche mesozoiche dell'Appennino centrale (Italia). In: Mancinelli, A. (ed.), *Biostratigrafia dell'Italia centrale*. Studi Geologici Camerti, pp. 9–129.
- Dragastan, O., 1975. Upper Jurassic and Lower Cretaceous micro-facies from the Bicaz Valley basin (East Carpathians). *Memoriile Institutului de Geologie și Geofizică*, 21: 1–87.
- Dragastan, O., 1980. *Alge calcareoase din Mezozoic și Terțiarul din România*. Editura Academiei R. S. R., București, 167 pp. [In Romanian.]
- Dragastan, O., 2010. *Platforma carbonatică Getică. Stratigrafia Jurasicului și Cretacicului inferior. Reconstituiri, paleogeografie, provincii și biodiversitate*. Editura Universității din București, București, 622 pp. [In Romanian.]
- Flügel, E., 2004. *Microfacies of Carbonate Rocks. Analysis Interpretation and Application*. Springer, Berlin, 984 pp.
- Flügel, E. & Koch, R., 1995. Controls on the diagenesis of Triassic carbonate ramp sediments: Steinplatte, Northern Alps, Austria. *Geologisch-Paläontologische Mitteilungen Innsbruck*, 20: 283–311.
- Frykman, D. K., 1986. Diagenesis of Silurian bioherms in the Klinteberg Formation; Gotland, Sweden. In: Schroeder, J. H. & Purser, B. H. (eds), *Reef Diagenesis*. Springer, Berlin, pp. 399–423.
- Hardie, L. A. & Shinn, E. A., 1986. *Carbonate Depositional Environments, Modern and Ancient. Part 3: Tidal flats*. Colorado School of Mines Quarterly, Golden, Colorado, 81: 1–74.
- Henrich, R. & Zankl, H., 1986. Diagenesis of Upper Triassic Wetterstein Reefs of the Bavarian Alps. In: Schroeder, J. H. & Purser, B. H. (eds), *Reef Diagenesis*. Springer, Berlin, pp. 245–268.
- Ivanova, D., 1999. Middle Callovian to Valanginian microfossil biostratigraphy in the West Balkan Mountain, Bulgaria (SE Europe). *Acta Palaeontologica Romaniae*, 2: 231–236.
- Ivanova, D. & Kołodziej, B., 2010. Late Jurassic–Early Cretaceous foraminifera from Štramberg-type limestones, Polish Outer Carpathians. *Studia Universitatis Babeș-Bolyai Geologia*, 55: 3–31.
- Ivanova, D., Kołodziej, B., Koleva-Rekalova, E. & Roniewicz, E., 2008. Oxfordian to Valanginian palaeoenvironmental evolution on the Western Moesian Carbonate Platform: A case study from SW Bulgaria. *Annales Societatis Geologorum Poloniae*, 78: 65–90.
- Kendall, A. C. & Tucker, M. E., 1973. Radial-fibrous calcite: a replacement after acicular carbonate. *Sedimentology*, 20: 365–389.
- Kerans, C., Hurley, N. F. & Playford, P. E., 1986. Marine diagenesis in Devonian reef complexes of the Canning Basin, Western Australia. In: Schroeder, J. H. & Purser, B. H. (eds), *Reef Diagenesis*. Springer, Berlin, pp. 357–380.
- Koch, R. & Schorr, M., 1986. Diagenesis of Upper Jurassic sponge-algal reefs in SW Germany. In: Schroeder, J. H. & Purser, B. H. (eds), *Reef Diagenesis*. Springer, Berlin, pp. 224–244.
- Kołodziej, B., 2015a. Geological context and age of the Štramberg-type limestones from the Polish Outer Carpathians: an overview. *Neues Jahrbuch für Geologie und Paläontologie – Abhandlungen*, 276: 173–179.
- Kołodziej, B., 2015b. Corals of the Štramberg-type limestones from Poland: taxonomic and palaeoecological aspects. *Neues Jahrbuch für Geologie und Paläontologie – Abhandlungen*, 276: 181–199.
- Kołodziej, B. & Decrouez, D., 1997. Foraminifera and incertae sedis from the Stramberk-type limestone (Tithonian–Berriasian). Preliminary results. *Revue de Paléobiologie*, 16: 145–150.
- Krajewski, M., 2000. Lithology and morphology of Upper Jurassic carbonate buildups in the Będkowska Valley, Kraków region, southern Poland. *Annales Societatis Geologorum Poloniae*, 70: 151–163.
- Lazăr, D. F., Bucur, I. I., Cociuba, I. & Săsăran, E., 2012. Sedimentary succession of the Lower Cretaceous deposits from the north-western part of Pădurea Craiului (Apuseni Mountains, Romania). *Studia Universitatis Babeș-Bolyai Geologia*, 57: 33–51.
- Lazăr, I., Grădinaru, M. & Petrescu, L., 2013. Ferruginous microstromatolites related to Middle Jurassic condensed sequences and hardgrounds (Bucegi Mountains, Southern Carpathians, Romania). *Facies*, 59: 359–390.
- Leinfelder, R., Nose, M., Schmid, U. D. & Werner, W., 1993. Microbial crusts of the Late Jurassic: composition, palaeoecological significance and importance in reef construction. *Facies*, 29: 195–230.
- Leinfelder, R. R., Schlagintweit, F., Werner, W., Ebli, O., Nose, M., Schmid, D. U. & Hughes, G. W., 2005. Significance of stromatoporoids in Jurassic reefs and carbonate platforms – concepts and implications. *Facies*, 51: 288–326.
- Lupu, M., Popescu, B., Szasz, L., Hann, H., Gheuca, I., Dumitrica, P. & Popescu, G., 1978. *Geological Map of Romania, 1:50 000, Sheet 126a, Vânturarița (Olănești)*. Institutul de Geologie și Geofizică, București.
- Masse, J.-P., Fenerci-Masse, M., Özer, S., Güngör, T. & Akal, C., 2015. Berriasian rudist faunas and micropalaeontology of Stramberk type carbonate exotics from the Lycian nappes, Bodrum Peninsula, southwest Turkey. *Cretaceous Research*, 56: 76–92.
- Michetiuc, M., Catincuț, C. & Bucur, I. I., 2012. An Upper Jurassic-Lower Cretaceous carbonate platform from the Vâlcan Mountains (Southern Carpathians, Romania): palaeoenvironmental interpretation. *Geologica Carpathica*, 63: 33–48.
- Mircescu, C. V., Bucur, I. I. & Săsăran, E., 2014. Dasycladalean

- algae from Upper Jurassic-Lower Cretaceous limestones of Piatra Craiului Massif (South Carpathians, Romania) and their relationship to paleoenvironment. *Studia Universitatis Babeş-Bolyai Geologia*, 59: 5–27.
- Morsilli, M. & Bosellini, A., 1997. Carbonate facies zonation of the upper Jurassic-lower Cretaceous Apulia platform margin (Gargano Promontory, southern Italy). *Rivista Italiana di Paleontologia e Stratigrafia*, 103: 193–206.
- Moshhammer, B. & Schlagintweit, F., 1999. The Ernstbrunn Limestone (Lower Austria): New data on biostratigraphy and applied geology. *Abhandlungen der Geologischen Bundesanstalt*, 56: 553–565.
- Mullins, H. T. & Cook, H. E., 1986. Carbonate apron models: alternatives to the submarine fan model for paleoenvironmental analysis and hydrocarbon exploration. *Sedimentary Geology*, 48: 37–79.
- Olivier, N., Carpentier, C., Bertrand, M. G., Lathuilière, M., Gaillard, C., Ferry, S., Hantzpergue, P. & Geister, J., 2004. Coral-microbialite reefs in pure carbonate versus mixed carbonate-siliciclastic depositional environments: the example of the Pagny-sur-Meuse section (Upper Jurassic, Northern France). *Facies*, 50: 229–255.
- Pleş, G. & Schlagintweit, F., 2014. New data on *Perturbatacrusta leini* Schlagintweit & Gawlick, 2011 (Calcareous sponge?) from the Late Jurassic (Tithonian) of the Southern Carpathians, Romania. *Acta Palaeontologica Romaniae*, 9: 3–9.
- Pleş, G., Mircescu, C. V., Bucur, I. I. & Săsăran, E., 2013. Encrusting micro-organisms and microbial structures in the Upper Jurassic limestones from Southern Carpathians. *Facies*, 59: 18–48.
- Pomoni-Papaioannou, P. F., Flügel, E. & Koch, R., 1989. Depositional environments and diagenesis of Upper Jurassic sub-surface sponge-and Tubiphytes reef limestones: Altensteig 1 well, western mollase basin, southern Germany. *Facies*, 21: 263–284.
- Pop, G. & Bucur, I. I., 2001. Upper Jurassic and Lower Cretaceous sedimentary formations from the Vâlcan Mountains (south Carpathians). *Studia Universitatis Babeş-Bolyai Geologia*, 46: 77–94.
- Popescu, G., 1954. Geological studies between Govora – Rm. Vâlcea – Olanesti regions (Getic Depression). *Dări de seamă ale şedinţelor Comitetului Geologic*, 38: 118–136.
- Qureshi, M. K. A., Butt, A. A. & Ghazi, S., 2008. Shallow shelf sedimentation of the Jurassic Samana Suk Limestone, Kala Chitta Range, Lesser Himalayas, Pakistan. *Geological Bulletin Punjab University*, 43: 1–14.
- Radoičić, R., 2005. New Dasycladales and microbiota from the lowermost Valanginian of the Mirdita zone. *Annales Géologiques de la Péninsule Balkanique*, 66: 27–53.
- Reitner, J., 1986. A comparative study of the diagenesis in diapir-influenced reef atolls and a fault block reef platform in the Late Albian of the Vasco-Cantabrian Basin (northern Spain). In: Schroeder, J. H. & Purser, B. H. (eds), *Reef Diagenesis*. Springer, Berlin, pp. 186–209.
- Rusciadelli, G., Ricci, C. & Lathuilière, B., 2011. The *Ellipsactinia* Limestones of the Marsica area (Central Apennines): a reference zonation model for Upper Jurassic Intra-Tethys reef complexes. *Sedimentary Geology*, 233: 69–87.
- Săndulescu, M., 1984. *Geotectonica României*. Editura Tehnică, Bucureşti, 336 pp. [In Romanian.]
- Săsăran, E., 2006. *Calcarele Jurasicului superior-Cretacicului inferior din Munţii Trascău*. Presa Universitară Clujeană, Cluj Napoca, 249 pp. [In Romanian.]
- Săsăran, E., Bucur, I. I., Pleş, G. & Riding, R., 2014. Late Jurassic Epiphyton-like cyanobacteria: indicators of long-term episodic variation in marine bioinduced microbial calcification? *Palaeogeography, Palaeoclimatology, Palaeoecology*, 401: 122–131.
- Schlagintweit, F. & Ebli, O., 1999. New results on microfacies, biostratigraphy and sedimentology of Late Jurassic–Early Cretaceous platform carbonates of the Northern Calcareous Alps. Part 1: Tressenstein Limestone, Plassen Formation. *Abhandlungen der Geologischen Bundesanstalt*, 56: 379–418.
- Schlagintweit, F. & Gawlick, H.-J., 2008. The occurrence and role of microencruster frameworks in Late Jurassic to Early Cretaceous platform margin deposits of the Northern Calcareous Alps (Austria). *Facies*, 54: 207–231.
- Schlagintweit F., Gawlick H.-J. & Lein, R., 2005. Mikropaläontologie und Biostratigraphie der Plassen-Karbonatplattform der Typlokalität (Ober-Jura bis Unter-Kreide, Salzkammergut, Österreich). *Journal of Alpine Geology*, 47: 11–102.
- Scholle, P. A. & Ulmer-Scholle, D. S., 2003. *A Color Guide to the Petrography of Carbonate Rocks*. American Association of Petroleum Geologists Memoir, 77: 1–474.
- Schroeder, R., Clavel, B., Conrad, M.-A., Zaninetti, L., Busnardo, R., Charollais, J. & Cherchi, A., 2000. Correlations biostratigraphiques entre la coupe d’Organya (Pyrénées Catalanes, NE de l’Espagne) et le Sud–Est de la France pour l’intervalle Valanginien–Aptien. *Treballs del Museu de Geologia de Barcelona*, 9: 5–41.
- Shinn, E. A., 1983. Tidal flat. In: Scholle, A. P., Bebout, D. G. & Moore, C. H. (eds), *Carbonate depositional environments* American Association of Petroleum Geologists Memoir, 33: 171–210.
- Şerban, D., Bucur, I. I. & Săsăran, E., 2004. Micropaleontological assemblages and microfacies characteristics of the Upper Jurassic limestones from Căprioara-Pojoga (Mureş Trough). *Acta Palaeontologica Romaniae*, 4: 475–484.
- Todiriţă-Mihăilescu, V., 1973. Contributions to the study of Cretaceous deposits on the northeastern flank of the Vânturariţa ridge. *Analele Universităţii Bucureşti*, 22: 89–98. [In Romanian.]
- Turi, V., Sasaran, E. & Bucur, I. I., 2011. New data on the Upper Jurassic–Lower Cretaceous limestones from Bihor Mountains: case study of Gârda Seacă-Hodobana region, Romania. *Studia Universitatis Babeş-Bolyai Geologia*, 56: 11–27.
- Turnšek, D., Buser, S. & Ogorolec, B., 1981. An upper Jurassic reef complex from Slovenia, Yugoslavia. In: Toomey, D. F. (ed), *European Fossil Reef Models*. Society of Economic Paleontologists and Mineralogists Special Publication, Tulsa, Oklahoma, 30: 361–369.
- Uţă, A. & Bucur, I. I., 2003. Microbial structures and microencrusters in the Upper Jurassic-Lower Cretaceous deposits from Buila-Vânturariţa Massif (South Carpathians). *Studia Universitatis Babeş-Bolyai Geologia*, 48: 3–14.