

Article ID: 192461
DOI: 10.5586/am/192461

Publication History
Received: 2024-05-24
Accepted: 2024-08-19
Published: 2024-10-03

Handling Editor
Piotr Zaniewski; Warsaw
University of Life Sciences –
SGGW, Warsaw, Poland;
<https://orcid.org/0000-0002-0792-9854>

Authors' Contributions
KS, MK, EAO: Research concept and design; Collection and/or assembly of data; Data analysis and interpretation; Writing the article; Critical revision of the article; Final approval of the article




Funding
The research was self-financed by the authors.

Competing Interests
No competing interests have been declared.

Copyright Notice
© The Author(s) 2024. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits redistribution, commercial and noncommercial, provided that the article is properly cited.

ORIGINAL RESEARCH

Aspicilia fluviatilis and *Circinaria leproscens*, two lichen species of the family Megasporaceae new to Poland

Katarzyna Szczepańska ^{1*}, Martin Kukwa ²,
Emilia Anna Ossowska ²

¹Department of Botany and Plant Ecology, Wrocław University of Environmental and Life Sciences, pl. Grunwaldzki 24a, 50-363 Wrocław, Poland

²Department of Plant Taxonomy and Nature Conservation, University of Gdańsk, Wita Stwosza 59, 80-308 Gdańsk, Poland

* To whom correspondence should be addressed. Email: katarzyna.szczepanska@upwr.edu.pl

Abstract

The paper presents two saxicolous species of the family Megasporaceae, *Aspicilia fluviatilis* and *Circinaria leproscens*, discovered as new to Poland. *Aspicilia fluviatilis* is a rare, arctic-alpine species with elongated marginal areoles, and *Circinaria leproscens*, occurs only in marine and strongly nitrophilous habitats and is often found sterile with granular to isidiolate upper cortex. The identity of *C. leproscens* was confirmed by molecular data, BLAST searches, and haplotype network analysis based on nucITS rDNA sequences. *Aspicilia fluviatilis* is also reported as new to Italy based on historical material from the Alps. Detailed information on morphology, chemistry, distribution, and ecology are given for both species.

Keywords

barcoding; biodiversity; haplotype network; lichen-forming fungi; taxonomy

Dedicated to Professor Lucyna Śliwa

1. Introduction

In recent years, the development of molecular methods has led to a significant increase in the number of taxa described as new to science (e.g., Guzow-Krzemińska et al., 2019; Kukwa et al., 2023; Nascimbene et al., 2023; Ossowska et al., 2022; van den Boom & Magain, 2020). At the same time, in many places around the world, including Poland, previously unrecorded species are still being discovered (e.g., Frisch et al., 2020; Khodosovtsev, 2023; Kossowska et al., 2022; Kukwa & Ossowska, 2023; Malíček et al., 2023; Matura et al., 2017; Ossowska et al., 2021; Ptach-Styn et al., 2024). Consequently, local lichen checklists must be constantly updated (Borgato et al., 2020; John et al., 2020; Tumur et al., 2021). According to the latest available data, the lichen biota of Poland comprises 1687 species (Fałtynowicz et al., 2024). However, there are still several lichen groups that need revisions, and that number will change.

The Megasporaceae is a lichen-forming fungal family, which includes eight genera: *Aspicilia* A. Massal., *Circinaria* Link, *Lobothallia* (Clauzade & Cl. Roux) Hafellner, *Megaspora* (Clauzade & Cl. Roux) Hafellner & V. Wirth, *Sagedia* Ach. (Nordin et al., 2010), *Teuvoa* Sohrabi & S. Leavitt (Sohrabi

et al., 2013), *Aspiciliella* M. Choisy (Zakeri et al., 2017) and *Oxneriaria* S.Y. Kondr. & Lőkös (Haji Moniri et al., 2017). Most of the family representatives are saxicolous or terricolous species and are often widely distributed. Although taxonomic revisions continue resulting in the recognition of new species (Fryday et al., 2021; Iqbal et al., 2023; Lee et al., 2022; Sohrabi et al., 2023), Megasporaceae is still perceived as taxonomically complicated, with species difficult in the identification due to the inconspicuous, crustose thalli with limited morphological and chemical characters (Zakeri et al., 2019). The most important features of the family are crater-like, urceolate or lecanorine apothecia, asci with a non-amyloid tholus, the 'Caesiocinerea-green' pigment (Meyer & Printzen, 2000) present in the epihymenium, simple and hyaline ascospores, and branched, anastomosing and swollen at tips (moniliform or submoniliform) paraphyses (Lumbsch et al., 1994; Nordin et al., 2010). Recognition of species within the family must be additionally based on the ascospore size, the number of ascospores in the ascus, length of conidia, secondary lichen metabolites, and habitat preferences (Szczepańska et al., 2023).

The occurrence and distribution of representatives of the Megasporaceae in Poland are still insufficiently known, as well

as the presumed number of species present. In addition to these listed by Fałtynowicz & Kossowska in (2016), four new species (*Aspicilia verrucigera* Hue, *Oxneriaria supertegens* (Arnold) S.Y. Kondr. & Lőkös, *Sagedia mastrucata* (Wahlenb.) A. Nordin, Savić & Tibell and *S. zonata* Ach.) occurring in the country have been recently recorded (Szczepeńska et al., 2023) and supplemented in the latest checklist of Fałtynowicz et al. (2024). Nevertheless, here we report two additional taxa of the family Megasporaceae new to Poland, which were discovered during the herbarium revision and the fieldwork.

2. Material and methods

The study material is deposited in UGDA (University of Gdańsk), KRAM (Polish Academy of Sciences, Kraków), and WRS� (University of Wrocław) herbaria. Descriptions of the species are based on our own observations. The morphology and anatomy of the species were studied with dissecting and light microscopes, following routine techniques; for light microscopy, hand sections were made with a razor blade and mounted in water. In the case of apothecia, hymenium measurements were made in water, and ascospore measurements were made in 10% KOH. The presence of the epihymenium pigment “Caesiocinerea-green” was detected by color reaction after the application of 50% HNO₃ (Meyer & Printzen, 2000). The TLC analyses were performed in solvents A and C using the standardized method of Culberson (1972) and following Orange et al. (2001).

Additionally, in both species, attempts were made to isolate DNA to a barcoding procedure and to confirm taxa identification. For *Circinaria leproscens*, sequences of ITS rDNA were obtained; however, in the case of *Aspicilia fluviatilis*, due to the small amount of material and the age of the specimen, DNA isolation was unsuccessful. The genomic DNA was extracted using a CTAB method according to the standard protocol of isolation (Doyle & Doyle, 1987; Guzow-Krzemińska & Węgrzyn, 2000). The primers used for DNA amplification were ITS1F (Gardes & Bruns, 1993) and ITS4 (White et al., 1990). PCR reactions were carried out using the following program: initial denaturation at 94 °C for 3 min and 33 cycles of 94 °C for 30 s; annealing at 52 °C for 45 s; extension at 72 °C for 1 min and a final extension at 72 °C for 10 min. PCR was performed in a volume of 25 µl using StartWarm HS-PCR Mix (A&A Biotechnology, Poland). PCR products were purified using Clean-Up (A&A Biotechnology, Poland) according to the manufacturer’s instructions. The purified DNA was sequenced by the MacroGen sequencing service (<http://www.macrogen.com>). For both sequences, we have obtained two F and R strands, which we have aligned in Auto Assembler v. 1.4.0 (Parker, 1997). The newly obtained nucITS rDNA sequences of *C. leproscens* were subjected to a BLAST search (Altschul et al., 1997) and then deposited in GenBank (<http://www.ncbi.nlm.nih.gov/genbank>). For haplotype network analysis, the only one *C. leproscens* sequence available in GenBank was downloaded and aligned in Seaview (Galtier et al., 1996; Gouy et al., 2010), and the terminal ends were cut. The TCS network, including three nucITS rDNA sequences (Clement et al., 2002) was created using PopArt software (<http://popart.otago.ac.nz>) and modified in Inkscape (<http://inkscape.org>).

3. The species

3.1. *Aspicilia fluviatilis* A. Nordin & Owe-Larss.

Lichenologist 43(1): 30. 2011.

Figure 1A–C.

Morphology: Thallus lichenized, crustose, thin, radiating, rounded, light creamy yellow in the center to brownish yellow and grey on the edge, matt. Areoles flat to slightly convex, angular, smooth, 0.2–0.4 mm in diam. in center, elongated and branched in the outer part, 0.5–1 mm long. Prothallus invisible. Apothecia immersed, 1–2 per areole, 0.2–0.3 mm in diam., thalline margin conspicuous, thick, same color as the thallus or darker, brighter and radially incised on the inside edge, disc rounded to irregular, not pruinose or with white pruina. Hymenium colorless, 50–60 µm tall, with submoniliform paraphyses (3–5 globose apical cells), epihymenium olive-brown, N+ clearly green, K+ orange-brown (Caesiocinerea-green), hypothecium colorless. Asci 8-spored, ascospores hyaline, simple, narrowly ellipsoid, 15–20 × 5–10 µm. Pycnidia not seen in the examined material.

Chemistry: thallus K+ yellow turning red, P+ yellow, C–, KC–. Norstictic acid in the cortex, as well as in the apothecium discs pruina was detected, which is in accordance with literature data (Nordin et al., 2011). However, probably due to very low content, the red needles and color change were not observed in the cross-section of epihymenium.

Distribution and ecology: *Aspicilia fluviatilis* is a very rare species occurring mainly in subalpine and alpine habitats. It grows on siliceous rocks near running water or on the rocky scree under mountainsides. Single records of the taxon come only from Norway, Sweden and Russia (Nordin et al., 2011). What is interesting, during the revision of materials stored in KRAM and WRS� herbaria, two specimens collected in the Italian Alps in the 19th by Arnold were found. They were attributed to *Aspicilia cinerea* f. *alpina* Arnold. Perhaps the name is a synonym of *Aspicilia fluviatilis*, but this needs the revision of the type material. The single Polish specimen of this species was also found in KRAM herbarium under *Aspicilia polychroma* Anzi var. *rubrireagens* Asta & Roux. The species was collected on mylonite rock in the subnival belt in the Polish High Tatra Mountains.

Comments: *Aspicilia fluviatilis* belongs to the group of taxa with elongated, marginal areoles, occurring mainly in arctic and alpine zones (Nordin et al., 2011). Because of norstictic acid present in the cortex, it can be confused with *A. granulosa* A. Nordin and *A. subradialis* (Nyl.) Hue, however these taxa have not been reported from Poland yet. Unlike *A. fluviatilis*, the main characteristic feature of *A. granulosa* is the granulose to subsiliate or verrucose central part of thallus (Nordin et al., 2011), whereas the typical feature of *A. subradialis* is the very dark, almost black color of the upper surface and usually indistinct marginal areoles (Thomson, 1984). Additionally, *A. fluviatilis* can be distinguished from both species by pruinose discs with pruina containing norstictic acid and thalline margin with a white inside edge.

The specimen described herein was found in a single locality in the High Tatra Mountains, determined as *A. polychroma* var. *rubrireagens*, and reported as a taxon new to Poland (Flakus, 2007). Despite similarities in secondary chemistry,




Figure 1 Specimens treated. (A–C) *Aspicilia fluviatilis* (KRAM L-50841), scales 1 mm; (D–F) *Circinaria leproscens* (UGDA L-62011), (D) – scale 2 mm, (E–F) – scales 1 mm.

the two species are different in some characters. *Aspicilia polychroma*, including var. *rubriregens* always has areolate or verrucose-areolate, not radiating thallus without any visible, elongate marginal areoles, and not pruinose, black disc. What is more, it usually occurs on slightly calciferous rocks in neutrophilic habitats (Szczeńska et al., 2023).

Specimen examined: Poland, West Carpathians, The High Tatra Mts, Szpiglasowa Przełęcz Pass, 49.198055°N, 20.042777°E, ATPOL grid square Ge-60, on mylonite rock, N aspect, slope 90°, mylonite area; the subnival belt, alt. 2107 m, 27 July 2003, leg. A. Flakus 971 (KRAM L-50841).

Additional material examined:

Italy, Alps, Trentino province, Lusia pass, Paneveggio-Pale di San Martino natural park, on porphyry rocks, 20 July 1888, leg. Arnold (KRAM L-996; WRS L-7949).

3.2. *Circinaria leproscens* (Sandst.) A. Nordin, Savić & Tibell

Mycologia 102 (6): 1346. 2010 ≡ *Aspicilia leproscens* (Sandst.) Hue, *Nouv. Arch. Mus. Hist. Nat. Paris* 5 sér. 2(1): 113. 1910.

Figure 1D–F.

Morphology: Thallus lichenized, crustose, 0.3–0.5 mm thick, areolate, smooth to irregularly cracked, light grey. Thallus surface granular with distinct, rounded, convex, brownish-grey granules resembling soredia or short isidia. Prothallus usually absent. Photobiont chlorococcoid. Apothecia and pycnidia not seen in the analyzed material.

Chemistry: thallus K–, P–, C–, KC–, medulla I–. Spicilin was detected in Polish samples, as reported by Smith et al. (2009) and Wirth et al. (2013).




Figure 2 Haplotype network showing relationships among ITS haplotypes of *Circinaria leproscens*. The new sequences from Poland are in red.

Distribution and ecology: *Circinaria leproscens* is a typical maritime and strongly nitrophilous species (Sheard, 1965), associated with siliceous seashore rocks, especially manured by birds in xeric (lightly sprayed by sea water) or supralittoral (rarely submerged in the lower regions but subject to heavy spray) zones (<http://www.lichensmaritimes.org>). The species is widely distributed and locally frequent. It has been reported mainly from Europe, including Denmark (Søchting & Alstrup, 2008), Finland, Norway and Sweden (Westberg et al., 2021), France (Roux, 2012), Germany (Wirth et al., 2013), Great Britain and Ireland (Smith et al., 2009), Italy (Nimis & Martellos, 2024), Netherlands (Aptroot et al., 1999) and Ukraine (Kondratyuk et al., 2021). In Poland, the species was found on stones at the shore of Pucka Bay.

Comments – *Circinaria leproscens* had not been previously reported from Poland; however, it was mentioned as probably occurring in Western Pomerania (Nowak & Tobolewski, 1975). During the revision of Megasperaceae specimens stored in Polish herbaria, only one specimen of the species was found, but originating from Sweden. Due to very characteristic ecology and habitat requirements connected with salted water, this species is rather easily recognized. Nevertheless, it seems that the thallus appearance of *C. leproscens* is not well defined. According to literature data, thallus surface is variously described as squamulate, papillate, granular, isidiate (Sheard, 1965; Smith et al., 2009), sorediate (Nowak & Tobolewski, 1975; Wirth et al., 2013) or forming globulose or flattened schizidia (Nimis & Martellos, 2024). Due to the morphological similarity (especially bright color and soredia-like granules), *C. leproscens* may be confused with some species of the genus *Lepraria* Ach., although *Lepraria* thallus is completely ecorticate, leprose and different in the secondary chemistry (Smith et al., 2009). In some cases, *C. leproscens* may resemble *Lepra corallina* (L.) Hafellner, because of the presence of granules, which possess pale grey thallus and short, cylindrical isidia on the thallus surface. Both taxa are, however, different in habitat preferences, apothecia and ascospores appearance, as well as secondary metabolites (thamnolic acid present in *L. corallina*) (Smith et al., 2009). Within the family Megasperaceae, *C. leproscens* may be confused mainly with *Aspicilia aquatica* (Fr.) Körb. and *Oxneriaria supertegens* (Arnold) S.Y. Kondr. & Lökös, occurring in humid habitats in the mountains and with *Circinaria caesocinerea* (Nyl. ex Malbr.) A. Nordin, Savić & Tibell, a common, saxicolous, lowland lichen (Smith et al.,

2009; Wirth et al., 2013). However, in each case, careful analysis of morphological and anatomical features of the specimens, connected with habitat data, will lead to the correct identification of species.

Two new nucITS rDNA sequences were obtained from our material. They show >98% similarity to the sequence of *C. leproscens* from Sweden (EU057911), which was deposited by Nordin et al. (2010). On the haplotype network, species shows little variation, with only two haplotypes that differ by one nucleotide position (Figure 2).

Specimen examined: Poland, Pobrzeże Kaszubskie, E of Puck, 54.716182°N, 18.429469°E, stones rinsed with seawater, ATPOL grid square Ac-49, on stone, 19 March 2023, leg. M. Kukwa 24551 & 24552 (UGDA L-61207 & 61208); ibidem, 08 Aug. 2023, leg. M. Kukwa 25122 & 25124 (UGDA L-63133 & 63135, GenBank PP768118); NE of Osłonino, by the cliff on the shore of Pucka Bay, 54.670731°N, 18.466133°E, ATPOL grid square Ac-59, on stones rinsed with seawater, 30 May 2023, leg. M. Kukwa 24810 (UGDA L-62011, Hb. K. Szczepańska 1431, GenBank PP768117).

Additional material examined: Sweden, Blekinge archipelago, Karlskrona city, 1875, leg. Svanlund (WRS�-5789).

Acknowledgments

We are deeply indebted to Dr hab. Adam Flakus, curator of KRAM herbarium, for a loan of the specimen that was analyzed and cited in this study.

References

- Altschul, S. F., Madden, T. L., Schäffer, A. A., Zhang, J., Zhang, Z., Miller, W., & Lipman, J. (1997). Gapped BLAST and PSI-BLAST: A new generation of protein database search programs. *Nucleic Acids Research*, 25, 3389–3402. <https://doi.org/10.1093/nar/25.17.3389>
- Aptroot, A., van Herk, K., Sparrius, L., & van den Boom, P. P. G. (1999). Checklist van de Nederlandse lichenen en lichenicole fungi. *Buxbaumia*, 50, 4–64.
- Borgato, L., Fryday, A. M., & Ertz, D. (2020). Preliminary checklist of the lichens and lichenicolous fungi of Martinique, with 144 new records. *Herzogia*, 33(1), 139–178. <https://doi.org/10.13158/hea.33.1.2020.139>
- Clement, M., Snell, Q., Walker, P., Posada, D., & Crandall, K. (2002). TCS: Estimating gene genealogies. In *Parallel and Distributed Processing Symposium (IPDPS '02)* (Vol. 2, p. 184). IEEE Computer Society. <https://doi.org/10.1109/IPDPS.2002.1016585>
- Culberson, C. F. (1972). Improved conditions and new data for identification of lichen products by standardized thin-layer chromatographic method. *Journal of Chromatography*, 72, 113–125. [https://doi.org/10.1016/0021-9673\(72\)80013-X](https://doi.org/10.1016/0021-9673(72)80013-X).
- Doyle, J. J., & Doyle, J. L. (1987). A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochemical Bulletin*, 19, 11–15.
- Fałtynowicz, W., Czarnota, P., Krzewicka, B., Wilk, K., Jabłońska, A., Oset, M., Ossowska, E. A., Śliwa, L., & Kukwa, M. (2024). *Lichens of Poland. A fifth annotated checklist*.

- Instytut Botaniki im. W. Szafera Polskiej Akademii Nauk.
<https://doi.org/10.35535/978-83-62975-47-1>
- Fałtynowicz, W., & Kossowska, M. (2016). The lichens of Poland. A fourth checklist. *Acta Botanica Silesiaca Monographiae*, 8, 3–122.
- Flakus, A. (2007). Lichenized and lichenicolous fungi from mylonitized areas of the subnival belt in the Tatra Mountains (Western Carpathians). *Annales Botanici Fennici*, 44(6), 427–449.
<https://www.jstor.org/stable/23727663>
- Frisch, A., Klepsland, J., Palice, Z., Bendiksby, M., Tønsberg, T., & Holien, H. (2020). New and noteworthy lichens and lichenicolous fungi from Norway. *Graphis Scripta*, 32(1), 1–47. <https://hdl.handle.net/11250/2731898>
- Fryday, A. M., Wheeler, T. B., & Etayo, J. (2021). A new species of *Aspicilia* (Megasporeaceae), with a new lichenicolous *Sagediopsis* (Adelococcaceae), from the Falkland Islands. *The Lichenologist*, 53(4), 307–315.
<https://doi.org/10.1017/S0024282921000244>
- Galtier, N., Gouy, M., & Gautier, C. (1996). SEAVIEW and PHYLO_WIN: Two graphic tools for sequence alignment and molecular phylogeny. *Computational Applied Biosciences*, 12, 543–548.
<https://doi.org/10.1093/bioinformatics/12.6.543>
- Gardes, M., & Bruns, T. D. (1993). ITS primers with enhanced specificity for basidiomycetes – Application to the identification of mycorrhizae and rusts. *Molecular Ecology*, 2, 113–118.
<https://doi.org/10.1111/j.1365-294X.1993.tb00005.x>
- Gouy, M., Guindon, S., & Gascuel, O. (2010). SeaView version 4: A multiplatform graphical user interface for sequence alignment and phylogenetic tree building. *Molecular Biology and Evolution*, 27, 221–224.
<https://doi.org/10.1093/molbev/msp259>
- Guzow-Krzemińska, B., Sérusiaux, E., van den Boom, P. P. G., Brand, A. M., Launis, A., Łubek, A., & Kukwa, M. (2019). Understanding the evolution of phenotypical characters in the *Micarea prasina* group (Pilocarpaceae) and descriptions of six new species within the group. *Mycologia*, 57, 1–30.
<https://doi.org/10.3897/mycokeys.57.33267>
- Guzow-Krzemińska, B., & Węgrzyn, G. (2000). Potential use of restriction analysis of PCR-amplified DNA fragments in taxonomy of lichens. *Mycotaxon*, 76, 305–313.
- Haji Moniri, M., Gromakova, A. B., Lőkös, L., & Kondratyuk, S. Y. (2017). New members of the Megasporeaceae (Pertusariales, lichen-forming Ascomycota): *Megaspora iranica* spec. nova and *Oxneriaria* gen. nova. *Acta Botanica Hungarica*, 59, 343–370.
<https://doi.org/10.1556/034.59.2017.3-4.5>
- Iqbal, M. S., Usman, M., Habib, K., & Khalid, A. N. (2023). *Oxneriaria pakistanica* sp. nov. (Megasporeaceae, Pertusariales, Ascomycota) from Darel Valley, Gilgit Baltistan, Pakistan. *Phytotaxa*, 579(2), 125–131.
<https://doi.org/10.11646/phytotaxa.579.2.6>
- John, V., Güvenç, Ş., & Türk, A. (2020). Additions to the checklist and bibliography of the lichens and lichenicolous fungi of Turkey II. *Archive for Lichenology*, 34, 1–49.
- Khodostovtsev, O. (2023). The first records of the lichen-forming, lichenicolous and allied fungi from Ukraine. *Folia Cryptogamica Estonica*, 60, 31–39.
<https://doi.org/10.12697/fce.2023.60.05>
- Kondratyuk, S., Popova, L. P., Khodosovtsev, O. Y., Lőkös, L., Fedorenko, N. M., & Kapets, N. V. (2021). The fourth checklist of Ukrainian lichen-forming and lichenicolous fungi with analysis of current additions. *Acta Botanica Hungarica*, 63(1–2), 97–163.
<https://doi.org/10.1556/034.63.2021.1-2.8>
- Kossowska, M., Kubiak, D., Kowalewska, A., Fałtynowicz, W., & Kukwa, M. (2022). Five lichen species new to Poland. *Folia Cryptogamica Estonica*, 59, 3–10.
<https://doi.org/10.12697/fce.2022.59.02>
- Kukwa, M., Kosecka, M., Jabłońska, A., Flakus, A., Rodriguez-Flakus, P., & Guzow-Krzemińska, B. (2023). *Pseudolepraria*, a new leprose genus revealed in Ramalinaceae (Ascomycota, Lecanoromycetes, Lecanorales) to accommodate *Lepraria stephaniana*. *Mycologia*, 96, 97–112.
<https://doi.org/10.3897/mycokeys.96.98029>
- Kukwa, M., & Ossowska, E. A. (2023). Additions to the biota of lichenicolous and lichenized fungi of Poland. *Folia Cryptogamica Estonica*, 60, 41–45.
<https://doi.org/10.12697/fce.2023.60.06>
- Lee, B. G., Shin, H. T., & Hur, J. S. (2022). A new lichen-forming fungus, *Aspicilia humida*, from a forested wetland in South Korea, with a taxonomic key for aspicilioid species of Korea. *Mycobiology*, 50(1), 20–29.
<https://doi.org/10.1080/12298093.2021.2021642>
- Lumbsch, H. T., Feige, G. B., & Schmitz, K. E. (1994). Systematic studies in the Pertusariales I. Megasporeaceae, a new family of lichenized Ascomycetes. *The Journal of the Hattori Botanical Laboratory*, 75, 295–304.
https://doi.org/10.18968/jhbl.75.0_295
- Malíček, J., Konečná, E., & Steinová, J. (2023). Contribution to the lichen biota of Romania. *Herzogia*, 36(2), 409–427.
<https://doi.org/10.13158/heia.36.2.2023.409>
- Matura, N., Krzewicka, B., & Flakus, A. (2017). Seven species of freshwater lichen-forming fungi newly recorded from Poland. *Polish Botanical Journal*, 62(2), 273–278.
<https://doi.org/10.1515/pbj-2017-0029>
- Meyer, B., & Printzen, C. (2000). Proposal for a standardized nomenclature and characterization of insoluble lichen pigments. *Lichenologist*, 32, 571–583.
<https://doi.org/10.1006/lich.2000.0294>
- Nascimbene, J., Nimis, P. L., Klüßendorf, J., & Thüs, H. (2023). Freshwater lichens, including new species in the genera *Verrucaria*, *Placopyrenium* and *Circinaria*, associated with *Lobothallia hydrocharis* (Poelt & Nimis) Sohrabi & Nimis from watercourses of Sardinia. *Journal of Fungi*, 9(3), Article 380. <https://doi.org/10.3390/jof9030380>
- Nimis, P. L., & Martellos, S. (2024). *ITALIC - The Information System on Italian Lichens. Version 7.0*. University of Trieste, Department of Biology. Retrieved March 14, 2024 from <https://dryades.units.it/italic>
- Nordin, A., Owe-Larsson, B., & Tibell, L. (2011). Two new *Aspicilia* species from Fennoscandia and Russia. *Lichenologist*, 43(1), 27–37.
<https://doi.org/10.1017/S0024282910000629>
- Nordin, A., Savić, S., & Tibell, L. (2010). Phylogeny and taxonomy of *Aspicilia* and Megasporeaceae. *Mycologia*, 102, 1339–1349. <https://doi.org/10.3852/09-266>

- Nowak, J., & Tobolewski, Z. (1975). *Porosty polskie* [Polish lichens]. PWN.
- Orange, A., James, P. W., & White, F. J. (2001). *Microchemical methods for the identification of lichens*. British Lichen Society.
- Ossowska, E. A., Guzow-Krzemińska, B., Szymczyk, R., & Kukwa, M. (2021). A molecular re-evaluation of *Parmelia encryptata* with notes on its distribution. *Lichenologist*, 53(4), 341–345. <https://doi.org/10.1017/S0024282921000219>
- Ossowska, E. A., Moncada, B., Kukwa, M., Flakus, A., Rodriguez-Flakus, P., Olszewska, S., & Lücking, R. (2022). New species of *Sticta* (lichenised Ascomycota, lobaroid Peltigeraceae) from Bolivia suggest a high level of endemism in the Central Andes. *MycKeys*, 92, 131–160. <https://doi.org/10.3897/mycokeys.92.89960>
- Parker, S. R. (1997). AutoAssembler sequence assembly software. *Methods in Molecular Biology*, 70, 107–117. <https://doi.org/10.1385/0-89603-358-9:107>
- Ptach-Styn, Ł., Guzow-Krzemińska, B., Lendemmer, J. C., Tønsberg, T., & Kukwa, M. (2024). Phylogeny of the genus *Loxospora* s.l. (Sarrameanales, Lecanoromycetes, Ascomycota), with *Chicitaeta* gen. nov. and five new combinations in *Chicitaeta* and *Loxospora*. *MycKeys*, 102, 155–181. <https://doi.org/10.3897/mycokeys.102.116196>
- Roux, C. (2012). Liste des lichens et champignons lichénicoles de France. *Bulletin de la Société linnéenne de Provence*, 16, 1–220.
- Sheard, J. W. (1965). *Lecanora* (Sect. *Aspicilia*) *leproscens* Sandst: New to the British Isles. *Lichenologist*, 3(1), 93–94. <https://doi.org/10.1017/S0024282965000129>
- Smith, C. W., Aptroot, A., Coppins, B. J., Fletcher, A., Gilbert, O. L., James, P. W., & Wolseley, P. A. (2009). *The lichens of Great Britain and Ireland*. British Lichen Society.
- Søchting, U., & Alstrup, V. (2008). *Danish Lichen Checklist. Version 2*. Faculty of Science, University of Copenhagen.
- Sohrabi, M., Leavitt, S. D., Obermayer, W., & Mayrhofer, H. (2023). *Circinaria nimisii* (Megasporaceae, lichenized Ascomycota), a new manna lichen from Greece. *Lichenologist*, 55(5), 367–376. <https://doi.org/10.1017/S0024282923000336>
- Sohrabi, M., Leavitt, S. D., Rico, V. J., Halici, M. G., Shrestha, G., & Stenroos, S. (2013). *Teuvoa*, a new lichen genus in Megasporaceae (Ascomycota: Pertusariales), including *Teuvoa junipericola* sp. nov. *Lichenologist*, 45, 347–360. <https://doi.org/10.1017/S0024282913000108>
- Szczepańska, K., Kukwa, M., Guzow-Krzemińska, B., & Urbaniak, J. (2023). New and rare lichens of the family Megasporaceae discovered in Poland. *Phytotaxa*, 598(2), 133–144. <https://doi.org/10.11646/phytotaxa.598.2.3>
- Thomson, J. W. (1984). *American arctic lichens: The microlichens* (Vol. 2). Univ of Wisconsin Press.
- Tumur, A., Mamut, R., & Seaward, M. R. (2021). An updated checklist of lichens of Xinjiang Province, China. *Herzogia*, 34(1), 62–92. <https://doi.org/10.13158/hea.34.1.2021.62>
- van den Boom, P., & Magain, N. (2020). Three new lichen species from Macaronesia belonging in Ramalinaceae, with the description of a new genus. *Plant and Fungal Systematics*, 65(1), 167–175. <https://hdl.handle.net/2268/247933>
- Westberg, M., Moberg, R., Myrdal, M., Nordin, A., & Ekman, S. (2021). *Santesson's Checklist of Fennoscandian Lichen-Forming and Lichenicolous Fungi*. Museum of Evolution, Uppsala University.
- White, T. J., Bruns, T., Lee, S., & Taylor, J. W. (1990). Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In M. A. Innes, D. H. Gelfand, J. J. Sninsky, & T. J. White (Eds.), *PCR Protocols: A guide to methods and applications* (pp. 315–322). Academic Press. <https://doi.org/10.1016/B978-0-12-372180-8.50042-1>
- Wirth, V., Hauck, M., & Schultz, M. (2013). *Die Flechten Deutschlands* [The lichens of Germany]. Ulmer.
- Zakeri, Z., Divakar, P. K., & Otte, V. (2017). Taxonomy and phylogeny of *Aspiciliella*, a resurrected genus of Megasporaceae, including the new species *A. portosantana*. *Herzogia*, 30, 166–176. <https://doi.org/10.13158/hea.30.1.2017.166>
- Zakeri, Z., Otte, V., Sipman, H., Malíček, J., Cubas, P., Rico, V. J., Lenzová, V., Svoboda, D., & Divakar, P. K. (2019). Discovering cryptic species in the *Aspiciliella intermutans* complex (Megasporaceae, Ascomycota) – First results using gene concatenation and coalescent-based species tree approaches. *Plos One*, 14, Article e0216675. <https://doi.org/10.1371/journal.pone.0216675>