

## Review article

---

S. SCHMITT-SCHILLIG, S. SCHAFFER, C.C. WEBER, G.P. ECKERT, W.E. MÜLLER

### FLAVONOIDS AND THE AGING BRAIN

Institute of Pharmacology (ZAFES), Biocenter Niederursel, University of Frankfurt,  
Frankfurt am Main, Germany

Like in all other organs, the functional capacity of the human brain deteriorates over time. Pathological events such as oxidative stress, due to the elevated release of free radicals and reactive oxygen or nitrogen species, the subsequently enhanced oxidative modification of lipids, protein, and nucleic acids, and the modulation of apoptotic signaling pathways contribute to loss of brain function. The identification of neuroprotective food components is one strategy to facilitate healthy brain aging. Flavonoids were shown to activate key enzymes in mitochondrial respiration and to protect neuronal cells by acting as antioxidants, thus breaking the vicious cycle of oxidative stress and tissue damage. Furthermore, recent data indicate a favorable effect of flavonoids on neuro-inflammatory events. Whereas most of these effects have been shown in vitro, limited data in vivo are available, suggesting a rather low penetration of flavonoids into the brain. Nevertheless, several reports support the concept that flavonoid intake inhibits certain biochemical processes of brain aging, and might thus prevent to some extent the decline of cognitive functions with aging as well as the development or the course of neurodegenerative diseases. However, more data are needed to assess the true impact of flavonoids on brain aging.

*Key words: brain aging, ROS, oxidative stress, flavonoids*

#### INTRODUCTION

As the elderly population increases, the prevalence of aging-related brain diseases, for example Alzheimer's disease (AD) or Parkinson's disease (PD), is likely to increase. Even if the cause of both neurodegenerative diseases is not yet finally known, enhanced oxidative stress seems to have an important contribution to brain aging in general and neurodegenerative diseases in specific. Therefore it becomes more and more important to develop drugs or nutraceuticals that

possibly are neuroprotective. The brain is the most susceptible organ to oxidative damage due to its high oxygen demand (1,2). Elevated oxygen consumption may lead to oxidative stress. The pathology of aging-associated neurodegeneration is increasingly linked to oxidative and nitrosative stress mediated by free radicals (3,4). Within the cell, free radicals, such as superoxide ( $O_2^-$ ) and hydroxyl ( $OH^\cdot$ ), are normal products of cellular oxygen metabolism. Moreover, other molecules like  $H_2O_2$  and peroxynitrite ( $ONOO^-$ ), although not themselves free radicals, can lead to generation of free radicals. Together, these highly reactive metabolites are referred to as reactive oxygen (ROS) and nitrogen species (RNS). Oxidative stress arises from an imbalance between cellular ROS/RNS production and the ability of cells to defend themselves against this stress. The attack of free radicals, ROS and RNS causes cellular damage by oxidizing proteins, membrane lipids, and DNA.

Normal cells have various mechanisms to protect themselves against oxidative and nitrosative attacks. Beside glutathione (GSH) and the vitamins C and E, all powerful antioxidant scavengers, the major cellular defense consists of antioxidative enzymes and bioactives such as carotenoids and flavonoids. Antioxidant enzymes in the brain include Cu/Zn superoxide dismutase (SOD-1) and Mn superoxide dismutase (SOD-2), both catalyzing the reaction of  $O_2^-$  to  $H_2O_2$ . Catalase and glutathione peroxidase (GPx) subsequently catalyze the conversion of  $H_2O_2$  to  $H_2O$ . The brain contains the highest concentrations of ascorbic acid of all body tissues (5). Vitamin C is able to scavenge a broad spectrum of ROS and is therefore one of the most important exogenous antioxidants in the body. Vitamin E is a lipid-soluble molecule with chain-breaking properties, which by restoring ascorbate reduced form in a regenerative cycle reaction, enhances its antioxidant activity. Hence, the ratio of both vitamins determines their antioxidant efficacy. However, both vitamin E and C recently failed to exert protective effects on cognitive function in humans (6). Recent data from epidemiological studies and animal experiments indicate that flavonoid-rich diets may positively influence brain aging and lower the incidence of neurodegenerative disorders such as AD and PD (1). The present article reviews not only recent data from *in vitro* studies pointing out possible mechanisms responsible for the observed neuroprotective effect of flavonoids, but also the outcome of *in vivo* studies using flavonoid supplements.

#### BRAIN AGING

Changes in brain function, for example the slowing down of information processing, can already be detected in middle-aged humans (35-65 year old) and rodents (12-24 month old) without fulfilling the criteria of dementia. Whereas neurodegenerative diseases, such as AD, are accompanied by a devastating loss of neurons, certain features of normal brain aging often proceed without

significant brain atrophy (7). However, impairment of neuronal activity in different brain regions represents a common feature of normal brain aging (8). One of the most popular theories for explaining the aging process is the free radical theory, initially proposed by Denham Harman (9). As outlined before, the brain is especially vulnerable towards the accumulation of oxidative stress-induced damage (2). Although the brain accounts for less than 2% of the bodyweight it consumes 20% of the basal oxygen uptake. High oxygen consumption is linked to leakage of electrons along the respiratory chain with subsequent radical formation. A second reason is the high amount of polyunsaturated fatty acids (PUFAs) present in neuronal membranes (10). These PUFAs are especially prone to undergo lipid peroxidation reactions resulting in the formation of cytotoxic aldehydes such as malondialdehyde (MDA) and 4-hydroxynonenal (HNE) (11). Furthermore, the insufficient supply of nutrients, caused by an impaired cerebrovascular function, contributes to the diminished neuronal capacity in the aging brain (12). Oxidative stress, deprivation of essential nutrient as well as age related and oxidative-stress induced damage hampers the activity of mitochondria, the key organelles for ATP production. Reduced energy supply will lead to reduced neuronal and finally brain function including impaired cognition.

#### FLAVONOID: BASIC PHARMACOKINETICS

Flavonoids belong to the broad group of polyphenols. Plants synthesize these secondary metabolites from the aromatic amino acids phenylalanine or tyrosin, or from malonate as defense against radiation and animals (13). Flavonoids contribute to the blue, orange or purple colors of fruits, leaves and flowers (14) and are mainly found in fruits, vegetables, nuts, grains, spices, seeds, wine, tea, and beer. By now, over 6000 different flavonoids have been identified (13) that can be divided in 6 classes due to their different structures: flavonoles, flavones, isoflavones, flavanoles (catechine), flavanones and anthocyanidins. Flavonoles, like quercetin and kaempferol, are most ubiquitous in food plants. The general flavonoid structure is a flavan nucleus, 15 carbon atoms arranged in three rings (A, B, C). The various classes differ in their level of oxidation and in their pattern of substitution of the c-ring (*Fig. 1*).

Flavonoids are almost exclusively present in human diet as glycosylated derivatives ( $\beta$ -glucosides) and not as aglycones, with the exception of catechins (flavanols). Flavonoid glycosides (FG), fairly large and highly polar molecules, are too hydrophilic to diffuse passively across biological membranes and the absorption of FGs was long believed negligible (15). Recent data, however, indicate that the bioavailability of FGs is determined by a complex network of absorption, metabolism and efflux processes (16-19). Despite rather high absorption rates, the bioavailability of flavonoids is comparatively low.





Figure 1: Main groups of flavonoids

Quercetin, for example, was found to reach 0.1 - 10  $\mu\text{mol/L}$  in the circulation. The concentration of quercetin was mainly due to the presence of quercetin metabolites rather than its aglycone, as recently reviewed by Murota and Terao (20). Consequently, the action of flavonoid metabolites is attracting increasing scientific interest (17,21).

#### *Effects on the blood brain barrier*

In spite of the evidence for flavonoid-dependent effects on brain function including neuroprotection, there is little information on their ability to cross the blood brain barrier (BBB) in order to reach the central nervous system (CNS). The BBB is formed by brain capillary endothelial cells tightly controlling the access of most small polar molecules and macromolecules into the brain. Youdim et al. (22) reported the uptake of several flavonoids and their relevant conjugated metabolites in a model of brain endothelial cells. Furthermore, the authors found that most of the tested flavonoids and respective metabolites are able to cross the BBB. The potential for BBB permeation is dependent on the compound's lipophilicity and the activity of efflux transporters, such as P-glycoprotein (Pgp), in the BBB (23). Inhibitory and stimulatory effects of quercetin on Pgp activity have been found *in vitro* and *in vivo*, recently (24,25). The exact mechanism, however, by which flavonoids modulate P-gp activity, is still unclear. Nonetheless, the ability of flavonoids to enter the brain has been confirmed by their identification in brain materials after p.o. and i.v. administration. Abd El Mohsen et al. (26) reported the presence of picomolar concentrations of epicatechin glucuronide and 3'-O-methyl epicatechin glucuronide in the brain of rats after oral supplementation with epicatechin (100 mg/kg bodyweight) for 1, 5 or 10 days. Similarly, naringenin and hesperetin have been detected in brain tissue after intravenous administration (27,28). These findings are supported by recent results from our laboratory, showing a biphasic modulatory effect of quercetin on the transport activity of Pgp in porcine brain capillary endothelial cells (PBCEC) and a human lymphocytic leukemia cell line expressing no Pgp (CEM) and their multidrug resistant variant (VLB) which express Pgp. At low concentrations (10 nM) quercetin activated the

transport of calcein-AM by Pgp and at high concentrations (1-3  $\mu\text{M}$ ) it significantly inhibited the transport in PBCEC cells (*Fig 2*). A similar, but not significant effect was observed in VLB cells. 30 nM quercetin activate the calcein transport by Pgp and 1-3  $\mu\text{M}$  inhibit this transport (25).



*Figure 2:* Influence of quercetin on the Calcein-AM uptake in VLB cells expressing Pgp (upper panel) and PBCEC cells expressing no Pgp (lower panel). Data are given as percent of calcein fluorescence under control condition and represent means  $\pm$  SEM for  $n = 6$  (PBCEC) /  $n = 5-6$  (VLB) experiments performed in triplicates. \*  $p < 0,05$ ; \*\*  $p < 0,01$ ; \*\*\*  $p < 0,001$ . Positive percentages indicate an inhibition of Pgp and negative values may be explained by activation of Pgp.

## NEUROPROTECTION

### *Direct antioxidant activities*

Flavonoids possess a highly reactive hydroxyl group that gets oxidized by electron-donation, thus stabilizing the radical to a less reactive molecule. One way of reaction is the direct scavenging of free radicals, for example superoxide anions, singlet oxygen and lipid peroxyl radicals. Structurally important features defining the reducing potential of flavonoids are the hydroxylation pattern, a 3',4'-dihydroxy catechol structure in the B-ring, the planarity of the molecule and the presence of 2,3-unsaturation in conjugation with a 4-oxo function in the C-ring. There is considerable evidence that flavonoids efficiently attenuate the deleterious effect of free radicals and ROS/RNS. Quercetin and some structurally related flavonoids, for example, showed a marked cytoprotective effect in PC12 cells exposed to  $\text{H}_2\text{O}_2$  (29). Also in PC12 cells, the addition of flavonoids or flavonoid-rich extracts suppressed the hydroperoxide-induced increase in ROS, thus improving cell survival (30,31). Matsuoka et al. (32) showed that tea catechins protect cultured newborn-mouse cerebral nerve cells from ROS-induced cell death after addition of glucose oxidase. The authors further examined learning ability of mice by a step-down-type passive avoidance test and memory impairment of mice by intracisternal injection of glucose oxidase or induced cerebral ischemia. Glucose oxidase-induced memory impairment was improved by intracisternal injection of epicatechin and i.v.

injection of catechin or epicatechin attenuated memory impairment induced by cerebral ischemia. These results suggest that tea catechins ameliorate ROS-induced cytotoxicity through intracellular scavenging. Another beneficial effect resulting of this scavenging activity is to decrease oxidation of membrane lipids. For example, gossypin dramatically inhibited lipid peroxidation initiated by Fe<sup>2+</sup> and ascorbic acid in rat brain homogenates. In addition, gossypin significantly attenuated the neurotoxicity induced by A $\beta$  (25-35) (33). Anthocyanidins inhibited H<sub>2</sub>O<sub>2</sub>-induced lipid peroxidation in rat brain homogenates, too. The ID<sub>50</sub> values of delphinidin, cyanidin, and pelargonidin for them were 0.7, 3.5, and 85  $\mu$ M, respectively (34). Antioxidant effects of flavonoids have furthermore been detected in vivo. The volume of the lipid peroxidation marker lipofuscin was significantly reduced in the brain of adult rats chronically exposed to ethanol (35). These findings are supported by data from Zhang et al. (16,36) who reported a decrease of ROS production, thiobarbituric acid reactive substances content and nitrite/nitrate concentration in brain homogenates in a Mongolian gerbil stroke model after Crataegus flavonoid supplementation. Furthermore, pretreatment of the animals with green catechins increased the antioxidative level and amount of biologically available NO by scavenging of superoxide anion produced during reperfusion (36,37). Total infarction volume in the ipsilateral hemisphere of ischemia/reperfusion rats was significantly lowered by treatment with the flavonoid balicalcin (50 mg/kg) given 2 h apart, promptly prior to and 2 h after reperfusion (38). Also significant ROS reduction in striatum of aged rats was found by Joseph et al. after 8 weeks of supplementing diet with either blueberry, strawberry or spinach extracts (39).

#### *Indirect antioxidant activities*

NADH-oxidase is a key enzyme in mitochondrial respiration. In a recent examination Hodnick et al. (40), compared hydroxylation and methoxylation patterns of flavonoids referring to their NADH-oxidase inhibition ability. The order of potency was robinetin, rhamnetin, eupatorin, baicalcin, 7,8-dihydroxyflavone and norwogonin. Cos et al. (41) reported that flavones have higher xanthine oxidase inhibitory activity than flavonoles. Xanthine oxidase catalyses the oxidation of xanthine and hypoxanthine to uric acid. During the re-oxidation of the enzyme superoxide radicals as well as H<sub>2</sub>O<sub>2</sub> are produced. The authors further showed that hydroxyl groups at C-3 and C-3' are essential for high superoxide scavenging. Recently, Suzuki et al. (37) administered catechin extract in drinking water to rats for five days before and during middle cerebral artery occlusion (MCAO) and reperfusion to examine their protective effects on various deteriorative processes following stroke. Catechins significantly ameliorated neurological deficits observed after reperfusion by the inhibition of iNOS expression and infiltration of neutrophils. In addition, the formation of peroxynitrite was found to be decreased due to the potent radical scavenging

Table 1: Neuroprotective effects of flavonoids and flavonoid-rich extracts

| Source  | Treatment                                | Beneficial properties   | Ref. |
|---|--|---|------|
| Strawberry extract,<br>Spinach extracts,<br>Vitamin E | 9.5 g/kg<br>6.4 g/kg<br>500 IU/kg        | Reduction of striatal ROS level in 6 month old rats after two weeks of supplementation  | (62) |
| Crataegus flavonoid extract                           | 0.5 mg/mL<br>2.5mg/mL                    | Protection against ischemia/reperfusion injury-caused delayed cell death in gerbils after 15 days of extract application through drinking water | (36) |
| Grape polyphenol extract                              | 5 mg/dl                                  | Prevention of ethanol-induced dopamine uptake activity decrease in young rats after grape polyphenol supplementation for 2 month                | (63) |
| Catechin  | 3.4 $\mu$ M<br>34 $\mu$ M<br>340 $\mu$ M | Reduction of lipid peroxidation in rat embryonic mesencephalic cell cultures after 6-OHDA-induced neuronal cell death                           | (64) |
| Resveratrol   | 30 mg/kg BW                              | Reduction of kainic acid-induced cell death in adult rat brains after i.p. administration for 5 days  | (65) |
| (-)-epigallocatechine gallate                         | 10 $\mu$ M<br>25 or 50 m/kg<br>i.p.      | Attenuation of MDA formation in rat embryonic hippocampal neurons<br>Reduction of postischemic brain edema in gerbils                           | (66) |

properties of catechins. Komatsu et al. recently found increased SOD activity in the mitochondrial fraction of striatums and midbrains of aged rats after one-month administration of beta-catechin solution (42).

#### *Anti-inflammatory activities*

Many flavonoids are known to possess anti-inflammatory properties and suppression of the inflammatory response in chronic diseases may beneficially affect disease outcome. Therefore flavonoids are getting more and more in the focus as a possibility to help preventing these diseases. Several mechanisms for the anti-inflammatory effect of flavonoids have been proposed. Wogonin, for example, a flavonoid from medicinal herb, was recently found to be a potent neuroprotectant from natural source. In co-cultured PC12 cells as well as *in vivo*, wogonin showed inhibition of inflammatory activation and reduction in microglial cytotoxicity (43). Wogonin diminished lipopolysaccharide-induced tumor necrosis factor-alpha (TNF-alpha), interleukin-1beta, and nitric oxide (NO) production *in vitro*. Latter effect of wogonin has been suggested to be facilitated by suppression of inducible NO synthase (iNOS) induction and NF-kappaB activation. The neuroprotective effect of wogonin was further demonstrated *in vivo* using two experimental brain injury models. In both animal models, wogonin conferred neuroprotection by attenuating the death of hippocampal neurons and the neuroprotective effect was associated with inhibition of the inflammatory activation of microglia. Also baicalein, a major flavonoid present in a traditional Chinese herb, showed potent anti-inflammatory properties on dopaminergic




Figure 3: ROS formation and detoxification

neurons and primary midbrain neuron-glia cultures from rat. The maximum protective effect on lipopolysaccharide-induced damage to dopaminergic neurons evaluated by uptake capacity for [(3)H]dopamine and tyrosine hydroxylase (TH)-immunocytochemistry was observed at a concentration of 5  $\mu$ M (44). Furthermore, baicalein prevented cell death in primary midbrain cultures, possibly due to its activity as a LOX-inhibitor (45). Quercetin and other flavonoids (e.g. catechin, chrysin, puerarin, naringenin, and genestein) also protected mesencephalic cultures from injury by MPP(+), which was shown by DNA fragmentation studies and tyrosine hydroxylase (TH) immunocytochemistry of mesencephalic dopamine (DA) neurons to occur by apoptosis. Similarly, catechin reduced cellular injury induced by hydrogen peroxide, 4-hydroxynonenal, rotenone, and 6-hydroxydopamine as shown by increases in cellular viability and [(3)H]DA uptake (46). Another polyphenol with anti-inflammatory effects is silymarin. This flavonoid recently demonstrated to exert neuroprotective effects against lipopolysaccharide (LPS)-induced neurotoxicity in mesencephalic mixed neuron-glia cultures (47).

#### Protection against $A\beta$ neurotoxicity

Aging is the most important risk factor for Alzheimer's disease (AD). Its neuronal pathology is characterized by the abundance of intracellular neurofilament tangles and the extracellular deposition of  $\beta$ -amyloid peptide ( $A\beta$ ) as senile plaques (48). The pathological effects of  $A\beta$  are significantly determined by its oxidative stress-inducing properties (49). As mentioned before, certain flavonoids are able to protect cells by acting as scavengers of ROS and RNS. Wang et al. (50) investigated in cortical neurons the effect of several flavonoids on  $A\beta$ -(25-35)-induced cytotoxicity. Pre-incubation with 50  $\mu$ M of kaempferol, apigenin, luteolin, and quercetin significantly lowered the  $A\beta$ -mediated increase



in ROS production. Furthermore, apigenin and kaempferol - but not quercetin - decreased A $\beta$ -induced neuronal death by alleviating the release of cytochrome c and the activation of the caspase cascade. Moreover, quercetin potentized the protective effect of apigenin. Neuroprotective effects of green tea and soy flavonoids due to their ability to scavenge A $\beta$ -induced ROS have also been shown by Choi et al. (51) and Zeng et al. (52), respectively. An additional explanation for the neuroprotective activity of flavonoids has been presented, recently. Myricetin, morin, and quercetin dose-dependently not only inhibited the formation and extension of A $\beta$  fibrils, but also destabilized preformed A $\beta$  fibrils with effective concentrations (EC<sub>50</sub>) in the order of 0.1-1  $\mu$ M. The underlying mechanisms responsible for the observed effects, however, are still unclear (53). Bastianetto et al. showed that flavonoids-rich *Ginkgo biloba* and red wine extracts are able to significantly protect hippocampal cells against A $\beta$  peptides- and/or oxidative stress-induced toxicities (37,54).

#### *Improvement of cognition in aging and dementia*

In a substantial number of studies about the effect of flavonoids or flavonoids-rich extracts on cognitive function in aged animals have been published (Table 2). Patil et al. (55) reported for aged but not young mice significant effects of quercetin supplementation on the step-through and transfer latency in the passive avoidance and elevated plus-maze task, respectively. Similarly, the learning time of aged mice supplied with catechin-enriched tap water was improved (56). Latter results are supported by Shirai and Suzuki (57) feeding adult and old mice with diets containing catechin. Time needed and distance travelled to reach the maze exit and the number of times strayed into blind alleys were significantly improved in the verum group. An extract of

Table 2: Cognition improvement properties of flavonoids and flavonoid-rich extracts

| Source   | Treatment                               | Beneficial properties   | Ref. |
|--|---|---|------|
| Apigenin-7-glucoside<br>Quercetin                          | 5-20 mg/kg i.p.<br>25-100 mg/kg<br>i.p. | Reduction of aging-induced impairment of<br>Passive Avoidance Performance in aged mice<br>after chronic administration for 7 days i.p.  | (67) |
| DHA and<br>Catechins                                       | 1.5 %<br>0,5 %                          | Enhancement of Maze behaviour in adult and<br>aged mice after 3.5 month intake of DHA and<br>catechin supplemented 5 % lard diet  | (68) |
| Green tea catechins  | 0,02 % in tap<br>water                  | Suppressive effect on cognitive dysfunction<br>markers (passive avoidance and Y maze test)<br>in accelerated senescence mice  | (69) |
| Strawberry extract<br>Spinach extract<br>Blueberry extract | 14.8 g/kg<br>9.1 g/kg<br>18.6 g/kg      | Reversal of age-related declines in several<br>cognitive and motor behavioural deficits in<br>aged male rats after 8 weeks of extract intake  | (39) |
| Quercetin  | 10-50 mg/kg                             | Amelioration of cognitive performance in<br>aged mice after chronic quercetin intake for<br>24 days measured as retention performance in<br>passive avoidance and elevated plus-maze test | (70) |

purple sweet potato anthocyanin markedly enhanced cognitive performance, assessed by passive avoidance test in ethanol-treated mice (58). Water-maze performance of aged rats was significantly improved after strawberry, spinach and, blueberry supplementation for 8 month (39).

Up to now there are few studies showing significant effects of flavonoids on cognitive deficits in aged humans or demented patients. One of the few reports (*Table 3*) indicates that the development of dementia might be slowed down by flavonoid rich diet. The Paquid Study was done in 1991-1996 in south-west France. The authors concluded that the relative risk of incident dementia was inversely related to the mean antioxidant flavonoid intake during the last 5 years as determined by a questionair. After adjustment to age, gender, education, weight and vitamin C intake, the relative risk of dementia for the highest flavonoid intake percentile was 0.49, respectively (1).

*Table 3: Effects of flavonoid intake on Dementia (Paquid Study, France) (1)*

| Study design  | Flavonoid consumption<br>(mg/day) | Age-adjusted model |             |         |
|---------------|-----------------------------------|--------------------|-------------|---------|
|               |                                   | RR                 | 95 % CI     | p-value |
| 1367 subjects | < 11.5 mg                         | 1.00               | -           | -       |
|               | 11.5, 16.2                        | 0.50               | 0.27 – 0.94 | 0.03    |
|               | > 16.2 mg                         | 0.59               | 0.33 – 1.06 | 0.08    |

Authors investigated whether flavonoid intake could be associated with a lower incidence of dementia in a cohort study of over 1300 subjects above 65 years of age. A questionnaire was used to evaluate their intake of flavonoids and subjects were followed-up for 5 years. Relative risk (RR) of dementia according to tertiles of flavonoid intake was estimated using a Cox model.

Reprinted with kind permission of Springer Science and Business Media.

## CONCLUSIONS

The constantly increasing number of people affected by neurodegenerative diseases is one of the major medical and social challenges of the 21<sup>st</sup> century. Although it is currently unknown whether oxidative stress is the primary cause of aging-related cognitive decline and dementia, the elevated release of ROS and RNS significantly contributes to the pathological changes observed in neurodegeneration (2). Flavonoids are ubiquitous in plant foods and beverages and can therefore be consumed with the daily diet (59). In numerous cell-free and cell-based experiments flavonoids demonstrated not only to possess excellent antioxidant activity but also to attenuate the accumulation of dementia-promoting metabolites, such as A $\beta$ , and to suppress inflammation by interacting with pro-inflammatory enzymes. Whereas detailed data on flavonoid consumption are available for humans, unfortunately only a few studies investigated a possible activity of flavonoid intake as neuroprotectants (1). However, as summarized in *Table 1* and *2*, results obtained in animal studies support health-beneficial effects

of flavonoids in the aging brain on both, the reduction of ROS-induced damage to biomolecules and the improvement of cognitive performance. Discrepancies in the magnitude of effect due to flavonoids supplementation are due to the diversity of bioactives used, to the different modality of administration, to the diversity of utilized methods, and to the timing of treatment (60). Furthermore, the effect of metabolic changes in the structure of flavonoids and their interaction with the blood brain barrier have not been adequately studied, yet, to draw a clear picture of the possible neuroprotective mechanism linked to flavonoids intake.

Despite of all these findings there is not enough data on the special effects of flavonoids or flavonoid-rich extracts *in vivo* to come to a final conclusion. The majority of data which are considered as evidence for this neuroprotection, however, come from studies of complex mixtures of compound with high flavonoid contents, such as green tea and herbal medicines, which have been recently reviewed by Youdim et al (61). Considering the health benefiting effects of flavonoid supplements or nutraceuticals, the absorption and brain permeability of the containing flavonoids have to be taken into account. Some studies only worked with the flavonoid aglycones and not their naturally occurring glycosides or metabolites. Nevertheless available data suggest a positive role of supplementation with flavonoids to slow down neurodegeneration in pathological processes and aging.

#### REFERENCES

1. Commenges D, Scotet V, Renaud S et al. Intake of flavonoids and risk of dementia. *Eur J Epidemiol* 2000; 16: 357-363
2. Halliwell B. Reactive oxygen species and the central nervous system. *J Neurochem* 1992; 59: 1609-1623
3. Eckert A, Keil U, Marques CA et al. Mitochondrial dysfunction, apoptotic cell death, and Alzheimer's disease. *Biochem Pharmacol* 2003; 66: 1627-1634
4. Barja G. Free radicals and aging. *Trends Neurosci* 2004; 27: 595-600
5. Rice ME. Ascorbate regulation and its neuroprotective role in the brain. *Trends Neurosci* 2000; 23: 209-216
6. Luchsinger JA, Tang MX, Shea S, Mayeux R. Antioxidant vitamin intake and risk of Alzheimer disease. *Arch Neurol* 2003; 60: 203-208
7. Finch CE. Neurons, glia, and plasticity in normal brain aging. *Neurobiol Aging* 2003; 24 Suppl 1: S123-S127
8. Rutten BP, Korr H, Steinbusch HW, Schmitz C. The aging brain: less neurons could be better. *Mech Ageing Dev* 2003; 124: 349-355
9. Harman D. Free radical theory of aging: effect of free radical reaction inhibitors on the mortality rate of male LAF mice. *J Gerontol* 1968; 23: 476-482
10. Floyd RA. Antioxidants, oxidative stress, and degenerative neurological disorders. *Proc Soc Exp Biol Med* 1999; 222: 236-245
11. Smith KJ, Kapoor R, Felts PA. Demyelination: the role of reactive oxygen and nitrogen species. *Brain Pathol* 1999; 9: 69-92
12. Zarchin N, Meilin S, Rifkind J, Mayevsky A. Effect of aging on brain energy-metabolism. *Comp Biochem Physiol A Mol Integr Physiol* 2002; 132: 117-120

13. Harborne JB ,Williams CA. Advances in flavonoid research since 1992. *Phytochemistry* 2000; 55: 481-504
14. Timberlake CF, Henry BS. Plant pigments as natural food colours. *Endeavour* 1986; 10: 31-36
15. Ross JA ,Kasum CM. Dietary flavonoids: bioavailability, metabolic effects, and safety. *Annu Rev Nutr* 2002; 22: 19-34
16. Nemeth K, Plumb GW, Berrin JG et al. Deglycosylation by small intestinal epithelial cell beta-glucosidases is a critical step in the absorption and metabolism of dietary flavonoid glycosides in humans. *Eur J Nutr* 2003; 42: 29-42
17. Sesink AL, Arts IC, Faassen-Peters M, Hollman PC. Intestinal uptake of quercetin-3-glucoside in rats involves hydrolysis by lactase phlorizin hydrolase. *J Nutr* 2003; 133: 773-776
18. Vaidyanathan JB, Walle T. Cellular uptake and efflux of the tea flavonoid (-)epicatechin-3-gallate in the human intestinal cell line Caco-2. *J Pharmacol Exp Ther* 2003; 307: 745-752
19. Walgren RA, Karnaky KJ, Jr., Lindenmayer GE, Walle T. Efflux of dietary flavonoid quercetin 4'-beta-glucoside across human intestinal Caco-2 cell monolayers by apical multidrug resistance-associated protein-2. *J Pharmacol Exp Ther* 2000; 294: 830-836
20. Murota K, Terao J. Antioxidative flavonoid quercetin: implication of its intestinal absorption and metabolism. *Arch Biochem Biophys* 2003; 417: 12-17
21. Spencer JP, Abd-el-Mohsen MM, Rice-Evans C. Cellular uptake and metabolism of flavonoids and their metabolites: implications for their bioactivity. *Arch Biochem Biophys* 2004; 423: 148-161
22. Youdim KA, Dobbie MS, Kuhnle G et al. Interaction between flavonoids and the blood-brain barrier: in vitro studies. *J Neurochem* 2003; 85: 180-192
23. Youdim KA, Shukitt-Hale B, Joseph JA. Flavonoids and the brain: interactions at the blood-brain barrier and their physiological effects on the central nervous system. *Free Radic Biol Med* 2004; 37: 1683-1693
24. Mitsunaga Y, Takanaga H, Matsuo H et al. Effect of bioflavonoids on vincristine transport across blood-brain barrier. *Eur J Pharmacol* 2000; 395: 193-201
25. Weber CC, Kressmann S, Fricker G, Muller WE. Modulation of P-glycoprotein function by St John's wort extract and its major constituents. *Pharmacopsychiatry* 2004; 37: 292-298
26. Abd El Mohsen MM, Kuhnle G, Rechner AR et al. Uptake and metabolism of epicatechin and its access to the brain after oral ingestion. *Free Radic Biol Med* 2002; 33: 1693-1702
27. Peng HW, Cheng FC, Huang YT, Chen CF, Tsai TH. Determination of naringinin and its glucuronide conjugate in rat plasma and brain tissue by high-performance liquid chromatography. *J Chromatogr* 1998; 714: 369-374
28. Tsai TH ,Chen YF. Determination of unbound hesperetin in rat blood and brain by microdialysis coupled to microbore liquid chromatography. *J Food Drug Anal* 2000; 8: 331-336
29. Dajas F, Rivera-Megret F, Blasina F et al. Neuroprotection by flavonoids. *Braz J Med Biol Res* 2003; 36: 1613-1620
30. Sasaki N, Toda T, Kaneko T, Baba N, Matsuo M. Protective effects of flavonoids on the cytotoxicity of linoleic acid hydroperoxide toward rat pheochromocytoma PC12 cells. *Chem Biol Interact* 2003; 145: 101-116
31. Horakova L, Licht A, Sandig G et al. Standardized extracts of flavonoids increase the viability of PC12 cells treated with hydrogen peroxide: effects on oxidative injury. *Arch Toxicol* 2003; 77: 22-29
32. Matsuoka Y, Hasegawa H, Okuda S et al. Ameliorative effects of tea catechins on active oxygen-related nerve cell injuries. *J Pharmacol Exp Ther* 1995; 274: 602-608
33. Yoon I, Lee KH, Cho J. Gossypin protects primary cultured rat cortical cells from oxidative stress- and beta-amyloid-induced toxicity. *Arch Pharm Res* 2004; 27: 454-459

34. Noda Y, Kaneyuki T, Mori A, Packer L. Antioxidant activities of pomegranate fruit extract and its anthocyanidins: delphinidin, cyanidin, and pelargonidin. *J Agric Food Chem* 2002; 50: 166-171
35. de F, V, da Silva PP, Assuncao M et al. Flavonoids from grape seeds prevent increased alcohol-induced neuronal lipofuscin formation. *Alcohol Alcohol* 2004; 39: 303-311
36. Zhang DL, Zhang YT, Yin JJ, Zhao BL. Oral administration of Crataegus flavonoids protects against ischemia/reperfusion brain damage in gerbils. *J Neurochem* 2004; 90: 211-219
37. Suzuki M, Tabuchi M, Ikeda M, Umegaki K, Tomita T. Protective effects of green tea catechins on cerebral ischemic damage. *Med Sci Monit* 2004; 10: BR166-BR174
38. Hwang YS, Shin CY, Huh Y, Ryu JH. Hwangryun-Hae-Dok-tang (Huanglian-Jie-Du-Tang) extract and its constituents reduce ischemia-reperfusion brain injury and neutrophil infiltration in rats. *Life Sci* 2002; 71: 2105-2117
39. Joseph JA, Shukitt-Hale B, Denisova NA et al. Reversals of age-related declines in neuronal signal transduction, cognitive, and motor behavioral deficits with blueberry, spinach, or strawberry dietary supplementation. *J Neurosci* 1999; 19: 8114-8121
40. Hodnick WF, Duval DL, Pardini RS. Inhibition of mitochondrial respiration and cyanide-stimulated generation of reactive oxygen species by selected flavonoids. *Biochem Pharmacol* 1994; 47: 573-580
41. Cos P, Ying L, Calomme M et al. Structure-activity relationship and classification of flavonoids as inhibitors of xanthine oxidase and superoxide scavengers. *J Nat Prod* 1998; 61: 71-76
42. Komatsu M, Hiramatsu M. The efficacy of an antioxidant cocktail on lipid peroxide level and superoxide dismutase activity in aged rat brain and DNA damage in iron-induced epileptogenic foci. *Toxicology* 2000; 148: 143-148
43. Lee H, Kim YO, Kim H et al. Flavonoid wogonin from medicinal herb is neuroprotective by inhibiting inflammatory activation of microglia. *FASEB J* 2003; 17: 1943-1944
44. Li FQ, Wang T, Pei Z, Liu B, Hong JS. Inhibition of microglial activation by the herbal flavonoid baicalein attenuates inflammation-mediated degeneration of dopaminergic neurons. *J Neural Transm* 2004;
45. Canals S, Casarejos MJ, de Bernardo S, Rodriguez-Martin E, Mena MA. Nitric oxide triggers the toxicity due to glutathione depletion in midbrain cultures through 12-lipoxygenase. *J Biol Chem* 2003; 278: 21542-21549
46. Mercer LD, Kelly BL, Horne MK, Beart PM. Dietary polyphenols protect dopamine neurons from oxidative insults and apoptosis: investigations in primary rat mesencephalic cultures. *Biochem Pharmacol* 2005; 69: 339-345
47. Wang MJ, Lin WW, Chen HL et al. Silymarin protects dopaminergic neurons against lipopolysaccharide-induced neurotoxicity by inhibiting microglia activation. *Eur J Neurosci* 2002; 16: 2103-2112
48. Selkoe DJ. Alzheimer's disease: genes, proteins, and therapy. *Physiol Rev* 2001; 81: 741-766
49. Varadarajan S, Yatin S, Aksenova M, Butterfield DA. Review: Alzheimer's amyloid beta-peptide-associated free radical oxidative stress and neurotoxicity. *J Struct Biol* 2000; 130: 184-208
50. Wang CN, Chi CW, Lin YL, Chen CF, Shiao YJ. The neuroprotective effects of phytoestrogens on amyloid beta protein-induced toxicity are mediated by abrogating the activation of caspase cascade in rat cortical neurons. *J Biol Chem* 2001; 276: 5287-5295
51. Choi YT, Jung CH, Lee SR et al. The green tea polyphenol (-)-epigallocatechin gallate attenuates beta-amyloid-induced neurotoxicity in cultured hippocampal neurons. *Life Sci* 2001; 70: 603-614
52. Zeng H, Chen Q, Zhao B. Genistein ameliorates beta-amyloid peptide (25-35)-induced hippocampal neuronal apoptosis. *Free Radic Biol Med* 2004; 36: 180-188
53. Ono K, Yoshiike Y, Takashima A et al. Potent anti-amyloidogenic and fibril-destabilizing effects of polyphenols in vitro: implications for the prevention and therapeutics of Alzheimer's disease. *J Neurochem* 2003; 87: 172-181

54. Bastianetto S, Quirion R. Natural extracts as possible protective agents of brain aging. *Neurobiol Aging* 2002; 23: 891-
55. Patil CS, Singh VP, Satyanarayan PS et al. Protective effect of flavonoids against aging- and lipopolysaccharide-induced cognitive impairment in mice. *Pharmacology* 2003; 69: 59-67
56. Unno K, Takabayashi F, Kishido T, Oku N. Suppressive effect of green tea catechins on morphologic and functional regression of the brain in aged mice with accelerated senescence (SAMP10). *Exp Gerontol* 2004; 39: 1027-1034
57. Shirai N, Suzuki H. Effects of Western, Vegetarian, and Japanese dietary fat model diets with or without green tea extract on the plasma lipids and glucose, and liver lipids in mice. A long-term feeding experiment. *Ann Nutr Metab* 2004; 48: 95-102
58. Cho J, Kang JS, Long PH et al. Antioxidant and memory enhancing effects of purple sweet potato anthocyanin and cordyceps mushroom extract. *Arch Pharm Res* 2003; 26: 821-825
59. Bravo L. Polyphenols: chemistry, dietary sources, metabolism, and nutritional significance. *Nutr Rev* 1998; 56: 317-333
60. Cantuti-Castelvetri I, Shukitt-Hale B, Joseph JA. Neurobehavioral aspects of antioxidants in aging. *Int J Dev Neurosci* 2000; 18: 367-381
61. Youdim KA, Spencer JP, Schroeter H, Rice-Evans C. Dietary flavonoids as potential neuroprotectants. *Biol Chem* 2002; 383: 503-519
62. Joseph JA, Shukitt-Hale B, Denisova NA et al. Long-term dietary strawberry, spinach, or vitamin E supplementation retards the onset of age-related neuronal signal-transduction and cognitive behavioral deficits. *J Neurosci* 1998; 18: 8047-8055
63. Sun GY, Xia J, Draczynska-Lusiak B, Simonyi A, Sun AY. Grape polyphenols protect neurodegenerative changes induced by chronic ethanol administration. *Neuroreport* 1999; 10: 93-96
64. Nobre Junior HV, Cunha GM, Maia FD et al. Catechin attenuates 6-hydroxydopamine (6-OHDA)-induced cell death in primary cultures of mesencephalic cells. *Comp Biochem Physiol C Toxicol Pharmacol* 2003; 136: 175-180
65. Wang Q, Yu S, Simonyi A et al. Resveratrol protects against neurotoxicity induced by kainic acid. *Neurochem Res* 2004; 29: 2105-2112
66. Lee H, Bae JH, Lee SR. Protective effect of green tea polyphenol EGCG against neuronal damage and brain edema after unilateral cerebral ischemia in gerbils. *J Neurosci Res* 2004; 77: 892-900
67. Patil CS, Singh VP, Satyanarayan PS et al. Protective effect of flavonoids against aging- and lipopolysaccharide-induced cognitive impairment in mice. *Pharmacology* 2003; 69: 59-67
68. Shirai N, Suzuki H. Effect of dietary docosahexaenoic acid and catechins on maze behavior in mice. *Ann Nutr Metab* 2004; 48: 51-58
69. Unno K, Takabayashi F, Kishido T, Oku N. Suppressive effect of green tea catechins on morphologic and functional regression of the brain in aged mice with accelerated senescence (SAMP10). *Exp Gerontol* 2004; 39: 1027-1034
70. Singh A, Naidu PS, Kulkarni SK. Reversal of aging and chronic ethanol-induced cognitive dysfunction by quercetin a bioflavonoid. *Free Radic Res* 2003; 37: 1245-1252

Received: January 31, 2005

Accepted: February 15, 2005

Author's address: Walter E. Müller, Institute of Pharmacology, Biocenter Niederursel, Marie-Curie-Strasse 9, 60439 Frankfurt am Main, Germany.  
E-mail: PharmacolNat@em.uni-frankfurt.de