



World Scientific News

WSN 16 (2015) 96-124

EISSN 2392-2192

Occurrence, ecotoxicology, removal of diclofenac by adsorption on activated carbon and biodegradation and its effect on bacterial community: A review

D. Cherik, M. Benali, K. Louhab*

Research Laboratory of Alimentary Technology, Faculty of Engineering Sciences,
University of Boumerdes, Boumerdes 35000, Algeria

*E-mail address: louhab_ka@yahoo.fr

ABSTRACT

This paper reviewed the occurrence of diclofenac in the ecosystem, its ecotoxicology and its removal. Treatment technologies were based on two methods. Firstly, the physicochemical process namely adsorption on activated carbon thanks to its economic benefits. At the other end, the second method is biological process as biodegradation by activated sludge and membrane bioreactor. In this latter process diclofenac can affect the bacterial community in sludge and influence the efficiency of treatment or biodegradation.

Keywords: diclofenac, adsorption, membrane bioreactor, conventional activated sludge, bacterial community

Table of Contents

1. Introduction
2. Occurrence of diclofenac
3. Ecotoxicology of diclofenac
4. Conventional wastewater treatment of anti-inflammatory diclofenac
 - 4.1. Adsorption on activated carbon
 - 4.2. Biodegradation of diclofenac
 - 4.2.1. Conventional activated sludge
 - 4.2.2. Membranes bioreactors
5. Effect of diclofenac on bacterial community
6. Conclusion
5. References

1. INTRODUCTION

Today, the occurrence of drugs in the aquatic environment has become a worldwide issue of increasing environmental concern [107]. Pharmaceutical products are consumed in very big quantity and present at very low concentrations ($\mu\text{g/L}$ to ng/L range) [1-4,69]. They are presented in the environment via various human activities, including direct discarding of unused medication, liberates from pharmaceutical manufacturing plants [4], human and veterinary drug use [5,6]. Furthermore, PCs are also introduced to the environment as they are excreted from humans as unchanged compounds, metabolites or conjugates via urine and feces and enter the municipal sewage system [7,8,65,79,93] and are not eliminated in wastewater treatment plants [9,78,109]. They represent a threat to both public health and the environment [10-13] due to their physicochemical properties, particularly polarity, water solubility, persistence, microbial resistance [14,15] and bioaccumulation in the food chain [16]. Sewage treatment plays an important part in removing contaminants from reclaimed water but it's known that conventional wastewater treatment plants (WWTP) do not remove all the pollutants [98], especially the persistent polar pollutants due to their physicochemical properties [25,90,91].


Most treatment methods of pharmaceuticals wastewater are physicochemical and conventional biological processes [17]. Activated carbon adsorption is frequent examples of physicochemical mechanisms [18-21]. Currently there are many adsorbents developed and reported for the removal of drugs pollutants from aqueous solutions and industrial effluents [22]. Biological wastewater treatment is one of the main important biotechnological processes [23,24] in which microorganisms are applied for removal of organic contaminants [4] and of which activated sludge Process [26,66] and membrane bioreactors are the most frequently used [83,85,89]. Most WWTPs use activated sludge processes where in microorganisms are utilized to mineralize the pollutants, or degrade them to acceptable forms. Pollutants can also be eliminated from water by denudation into air or by sorption onto sludge that is regularly discharged. Membrane bioreactors (MBR) combine the biological degradation of waste products with membrane filtration [84]. The use of MBR in wastewater treatment is becoming increasingly important, since their various benefits; high biodegradation efficiency, smaller footprint and less sludge production [87].

In these biological processes (activated sludge and membrane bioreactor) using microorganisms [145,147] such as ammonia-oxidizing bacteria for the removal of ammonium [151], *Escherichia coli* [155], Methanogens [156] and nitrite-oxidizing bacteria [140] and as micropollutants may affect the ecosystem and human health [10-13], they can also affect these bacterial communities as well as their microbial diversity [142-144,155] and thus this impact affects the proper functioning of the wastewater treatment plant [147], which explains the occurrence of micropollutants such as pharmaceutical residues in effluents of these treatment plants [9, 78 and 109], but unfortunately, there are few studies concerning the impact of micropollutants on the bacterial community.

Diclofenac (DCF), a non-steroidal anti-inflammatory drug [27,81] is one of the most extensively studied pharmaceuticals, for his potential toxic effects on no-target organisms [28]. Diclofenac was detected in wastewater [94,86] in a number of countries. According to published documents, the concentrations of this drug varied in a range from 0.01 to 8.5 $\mu\text{g/l}$ [38,16] the rate of his elimination in sewage treatment plants is low (up to 40%) [29,123]. Table 1 present physicochemical properties and molecular structure of Diclofenac.

In this context, the objective of this study is to write a review of the literature by recent publications and studies in full and small scale between 2008s and 2015 on the presence of diclofenac in water, his toxicity, removal and effects on bacterial community. We inspect the performance of different water treatment systems in the elimination of this drug, particularly technologies that use adsorption on activated carbons and biodegradation with biological processes as well as activated sludge and membrane bioreactor plus an analysis of the influence of pharmaceutical residues and diclofenac on bacterial diversity in the sludge during its removal.

Table 1. Main characteristics of diclofenac.

Structure ^a	Molecular weight (g/mol) ^a	Log Kow ^{a, b}	pKa ^a
	296.14	0.7-4.51	4.15

^a; [30]

^b; [156]

2. OCCURRENCE OF DICLOFENAC

The non-steroidal anti-inflammatory; diclofenac is the most habitually identified in the aquatic environment [31,80], including drinking water [32] and that is usually used to treat rheumatoid arthritis [31,33]. The occurrence of Diclofenac in wastewaters was observed in Asia countries such as Korea and Taiwan with concentration of up to 1 µg l⁻¹ [54-57]. [34] Investigated the spatial distributions and seasonal variations of diclofenac (DFC) in the Huangpu River system (a drinking water source for Shanghai in China) over a period of practically two years and the concentration was 13.6 ng/l. [101] also investigated the occurrence of diclofenac in surface water (Beiyun River) of Beijing in three sampling events representing different seasonal flow conditions, diclofenac was detected with detection frequencies 100%, this high detection of diclofenac reflect his extensive use in Beijing.

The levels were the same in two seasons (March and June) with median concentrations 67 ng/l and 67.8 ng/l, respectively and 57.3 ng/l in September. This difference due to the period of seasons, where period from March to June is an arid season and the flow was low causal a higher levels and in September the season is moist and the flow was high causal a lower levels.

Table 2. Diclofenac concentrations detected in environment of various countries.

Country	Samples	Concentration (ng/l)	References
USA	drinking water	0.25	[99]
	effluent	<10	[96]
Taiwan	Drug production facilities	20733	[54]
	Hospitals	286	
	effluent	61	
	Regional discharges	84	
	Animal husbandries	4	
	Aquacultures	4	
	Hospital	10	[55]
	Influent	58-367	[105]
	Effluent	25-182	
China	River system (drinking water)	13.6	[34]
	Pearl Rivers water	1.1	[100]
Korea	Influent	131	[57]
	Effluent	24	
Spain	Urban wastewater	570	[59]
	Industrial wastewater	140	
	Effluent	<15	[60]
Serbia	Municipal waste water	1338	[64]
	Surface water	<324	
Greece	Influent wastewaters		[62]
	- Ioannina City	100.8	
	- Ioannina hospital	180.7	
	- Arta	106.9	
	- Preveza	1346.0	
	- Agrinio	98.8	
	- Grevena	78.3	
	- Kozani	83.1	
	- Veroia	-	
	Effluent wastewaters		
	- Ioannina City	98	
	- Ioannina hospital	63.2	
	- Arta	162.5	
	- Preveza	56.1	
	- Agrinio	154.9	
- Grevena	-		
- Kozani	97.1		
- Veroia	145.6		
Portugal	Seawater	30.6	[58]

These results confirm the results found by [119] where they observed that concentrations average of diclofenac is bigger in the dry season than the wet. In Western Balkan Region such as Serbia [64] studied the occurrence of 81 pharmaceuticals drugs using solid-phase extraction and the presence of this pharmaceuticals compounds was analyzed by Ultra-High Performance Liquid chromatography coupled to mass spectrometry with hybrid triple quadrupole-linear ion trap (UPLC-QqLIT-MS/MS), Results shown the presence of Diclofenac with concentration of 1.34 µg/l in Danube River with frequency of occurrence 11.11%. [96] Were surveyed monthly for 12 months the presence of this drug after different steps of treatment in an advanced wastewater reclamation plant in Gwinnett County, GA, U.S.A. The results show that diclofenac was detected in primary effluent with 220 ng/l of average concentration, 99 ng/l in membrane effluent, granular active carbon effluent and final effluent with average concentrations below 10 ng/l. Diclofenac was also detected in Spain [59,60,73], in Greece [62,110], in USA [97], in Sweden [63], Portugal [58,67], Rio de Janeiro in Brazil [88] and in South Wales region of the United Kingdom (UK) [61,95]. In the African continent diclofenac was detected in the concentration range of 1.1-15.6 µg/l in South Africa [152]. Table 2 shows the concentrations of diclofenac detected in WWTPS of various countries. Note that concentrations in WWTPS effluents in Asia countries such as Taiwan [54,105] (from 25 ng/l to 182 ng/l) are more than in Europe [60] (<10 ng/l) and America [96] (<15 ng/l) countries.

3. ECOTOXICOLOGY OF DICLOFENAC IN

The removal percentage of DCF during wastewater treatment processes typically ranges from 21% to 40% [29], which justifies their occurrence in surface water, groundwater and even in drinking water at very low concentration [34, 35] and adverse effects in different organisms were observed with high BOD, COD [106]. It has considered being a very toxic anti-inflammatory due to the recorded death of birds shortly after scavenging on contaminated livestock in India and Pakistan [35, 36]; it has showed a potential endocrine disrupting influence by delaying and reducing hatching success of fish eggs and low concentrations damage digestive organs [36]. [92] Suggested that the anti-inflammatory drug diclofenac caused oxidative stress and reduced testosterone level that can has a negative impact in aquatic organisms. [111] explained the effects of three drugs, diclofenac, ibuprofen and propanol on Baltic Sea blue mussels using *Mytilus edulis trossolus* as a model organism.

The uptake and stress engendered in *Mytilus edulis trossolus* from exposure to diclofenac was studied using the two physiological indicators; strength and abundance of byssus threads, and scope for growth (SFG). Results shows that exposure to diclofenac would lead to bio concentration in the mussels, and negatively affect the regeneration of the byssus threads (weaker and lower abundance of threads) and decrease the scope for growth. [112] studied the effect of diclofenac exposure over 15 days (250ng/l) in mussel *Mytilus galloprovincialis* and evaluated the responses of biomarkers in the mussel tissues; this study show that mussel were affected during initial days.

The antioxidant system was rapidly triggered, mostly in the gills; evidence of damage was observed in digestive glands. DCF made a direct relationship between the acetylcholinesterase activity and the vitellogenin-like protein levels in females. [113] also studied the effect of diclofenac exposure on the marine mussel (*Mytilus* spp.) and results

shows a lower response in the biomarker approach but with an induction of lipid peroxidation (LPO) after 96 h diclofenac potential in tissue damaging has been seen. [114] investigated the Toxicological effects of clofibric acid and diclofenac on plasma thyroid hormones of an Indian major carp, *Cirrhinus mrigala* during short (96 h) and long-term (35 days) exposures to different concentrations (1, 10 and 100 µg/l), this study demonstrated that diclofenac has a profound effect on plasma thyroid hormones an Indian major carp, *Cirrhinus mrigala* in short and long term. [118] evaluated also the effects of diclofenac on early life stages of common carp (*Cyprinus carpio*) during 30 days and the progress of carp performed to be delayed at the beginning of the test and any damaging impact was observed on the hatching and viability of carp embryos. [115] studied the consequences of exposure to diclofenac up to 3 months on two freshwater cladocerans (*Daphnia magna* and *Moina macrocopa*) and Japanese medaka (*Oryzias latipes*). Clear diminution of reproduction was observed at 25 mg/l for *Daphnia magna*, and at 50 mg/l for *Moina macrocopa*.

For fish (*Oryzias latipes*), the chronic exposure for 3 months of diclofenac at 0,001-10 mg/l causes an important reduction in hatching success and delay in hatch and at 10mg/l the hatching of eggs was completely interfered and Gonadosomatic index (GSI) of female fish was affected also, whereas fertility of the parent generation was not affected. [116] also evaluated the Toxicity screening of Diclofenac, Propranolol, Sertraline and Simvastatin using zebrafish (*Danio rerio*) and sea-urchin (*Paracentrotus lividus*) embryo bioassays, results show that effects of diclofenac has a delicate endpoints on hatching, 75% epiboly-stage, abnormal cellular growth and yolk-sac abnormalities these results were observed in [117], this consequence contributed to a significant rise of total abnormalities and high sensitivity of sea urchin embryos to diclofenac.

Diclofenac also has effects on the soil [120] studied his time dissipation and the consequences of exposure on *Folsomia candida*. The results indicated that diclofenac can exert severe effects on the survival of adult *F. candida*, so limiting the reproductive potential of the population and the impact of diclofenac on the neural system may be different in soil invertebrates compared to vertebrates.

4. CONVENTIONAL WASTEWATER TREATMENT OF ANTI-INFLAMMATORY DICLOFENAC

4. 1. Adsorption on activated carbon

The adsorption process on activated carbon is most commonly used to remove diclofenac present in the aquatic environment thanks to its economic benefits to have an effective treatment of polluted waters [37,39,42,82].

The activated carbon is most collectively used thanks to his high surface area and its superficial chemical properties. The surface of the activated carbon is hydrophobic can contain functional groups. The latter contains heterosexuals-atoms to know l-hydrogen, the oxygen, nitrogen, chlorine and sulfur [40]. They depend on the nature of the activation and contribute to the acid or basic character of the surface [41]. By-products such as olive nuts [44], carbon nanotube [46], mesoporous silica functionalized with different organic moieties [74] and zeolite [51] can be valued and testified for the removal of diclofenac from aqueous solutions.

The activated carbon can be used in two forms; powder (PAC) and granular (GAC), activated carbon's ability to remove the pollutant depends on its properties as pH of point of

zero charge (pH_{pzc}) and the properties of diclofenac as its solubility in water and also the reaction parameters namely the contact time, the dose of activated carbon, reaction temperature and the pH of solution [40]. Several studies have demonstrated the efficiency of the elimination of diclofenac on the activated carbon powder (PAC) or granular (GAC) (Table 3). [1] investigated the efficiency removal of diclofenac by two commercially available activated carbon; Clarimex 061 CAE and Epibon YM 12X40.

This study showed that the increase in concentration of diclofenac from 10 mg/ l to 50 mg/ l promotes the elimination rate of 68% to 99%, respectively for Clarimex 061 CAE, by against in the case of the use of Epibon the elimination rate the rate of elimination decreased from 82.70 % to 70.20 %. [157] examined the correlation between the coefficient of partition octanol-water ($\log K_{ow}$) and the adsorption, they found that by increasing the concentration of diclofenac which is a hydrophobic compound, the adsorption was better presented by the Freundlich isotherm ($R^2 > 0.92$) which is positively correlated with $\log K_{ow}$ ($R^2 = 0.8397$).




Fig. 1. Structures of diclofenac and the degradation products identified by NMR [70].

Table 3. Key publications on diclofenac adsorption on activated carbons.

Reference	Adsorbent	Experimental conditions	Observations	Adsorption capacity
[43]	Isabel grape bagasse	[C ₀] = 5-30 mg l ⁻¹ T = 22°C V = 25ml Carbone dose = 5mg pH = 5 time contact = 72h	Isabel grape bagasse before being crushed and sieved to a particle size less than 150 µm. After this procedure, the adsorbent was used without any physical or chemical pretreatment.	q _e = 5.12-23.77mg g ⁻¹
[44]	GAC (from exhausted olive-waste cake)	[C ₀] = 14.8 mg l ⁻¹ T = 25°C V = 300ml Carbone dose = 100-450 mg pH = 4.1 time contact = 26h	Activated carbon was produced via chemical activation using phosphoric acid	q _e = 13.776 mg g ⁻¹
[45]	functionalized silica-based porous materials	[C ₀] = 100 µg l ⁻¹ T = 25°C Carbone dose = 2 g l ⁻¹ time contact = 4 h pH = 7	Modification of HMS with either aminopropyltriethoxy or 3 mercaptopropyltrimethoxy (for amine- or mercapto- modification, respectively) was performed by a co- condensation method following the procedure described by Lee et al. [20]. SBA-15 was prepared under acidic conditions according to the synthesis procedure described by Zhao et al. [27], whilst MCM-41 was synthesized under basic conditions following the procedure described by Kim et al. [28]. The procedures for synthesis of the adsorbents used in this study were described in Supplementary information, pages 10–11.	q _{eHMS} = 31.93 mg g ⁻¹ q _{eM-HMS} = 35.59 mg g ⁻¹ q _{eA-HMS} = 5.89 mg g ⁻¹ q _{eSBA-15} = 34.18 mg g ⁻¹ q _{eMCM-41} = 32.64 mg g ⁻¹ q _{ePAC} = 40.55 mg g ⁻¹
[46]	granular carbon nanotubes/alumina hybrid	[C ₀] = 15.5 µmol l ⁻¹ T = 25°C Carbone dose = 25mg time contact = 72 h pH = 6	The granular CNTs/Al ₂ O ₃ hybrid adsorbents with the CNTs/Al ₂ O ₃ mass ratios of 1:1, 1:3, 1:5, 1:7 and 1:9 were prepared by a solegel method (granular adsorbent cannot be prepared at less alumina content than 1:1). A certain amount of pseudo boehmite predetermined according to the mass ratio of CNTs/Al ₂ O ₃ was added. In to 10 mL of diluted nitric acid solution containing 0.04 g Brij 35.	Q _{max(CNTs/Al₂O₃)} = 106.5µmol/g

Reference	Adsorbent	Experimental conditions	Observations	Adsorption capacity
[47]	activated biochars	[C ₀]= 20 μM T= 25°C Carbone dose= 2g l ⁻¹ time contact= 72 h pH= 7	After stirring for 10 min, 0.2 g MWCNTs was slowly added into the mixture with vigorous 3 stirring for 10 min, and then five drops of diluted aqueous ammonia were added into the mixture to form the gel. The gel was dried for about 15 min at 80°C to remove the extra ammonia on the surface and then was granulated in the size of about 3 mm. The granules were aged at 25°C for 8 h, followed by drying at 50°C for 12 h. Finally, the dried granules were calcined at 450°C for 4 h with a heating rate of 2°C/min.	q _{N-biochar} = 372 mg g ⁻¹
[53]	GAC and PAC	[C ₀]= 20 μM T= 25°C Carbone dose= 2g l ⁻¹ time contact= 72 h	Biochars produced by pyrolyzing torrefied loblolly pine chip (15*6 mm) at 300°C for 15 min were classified as either N-biochar or O-biochar in terms of pyrolysis with pure nitrogen or 7% oxygen with 93% nitrogen gas, respectively. Both biochars were activated by NaOH to increase the surface area and pore volume of the biochars. Biochars were milled and passed through a 200-mesh (74lm) sieve, and then stored with ultrapure water (2000 mg/L) for 24 h as a stock solution to hydrate prior to use; the designed dosages of biochar were confirmed by measuring control samples. Three NSAIDs One granular activated carbon and two powdered activated carbons	q _{O-biochar} = 214 mg g ⁻¹ q _e = 87.4 μg g ⁻¹
[49]	trimethylsilylated mesoporous SBA-15	[C ₀] = 100 μg/l T = 25°C Carbone dose = 1g l ⁻¹ time contact = 24 h pH = 5.5	Calcined SBA-15 (1.0 g) was Heated at 150°C in vacuo for 6 h and then slurried in anhydrous toluene (100 mL) for 1 h. Next, a five-fold excess amount of HMDS (3.0ml) in anhydrous toluene (50 ml) was added and stirring was continued at room temperature [21] for 24 h. The resulting product was filtered and washed with different solvents in the following order: toluene, acetone, ethanol, ethanol/water (50:50, v/v), water, ethanol, and acetone. The sample was then dried in vacuo at 150°C for 4 h and stored in a desiccator.	q _e = 87.40 μg/g
Reference	Adsorbent	Experimental conditions	Observations	Adsorption capacity
[48, 50]	Carbone Xerogels	[C ₀] = 100 mg/l T = 31°C Carbone dose = 60 mg	A Carbon Xerogel (CX) was prepared by polycondensation of resorcinol with formaldehyde (with a molar ratio of 1:2). The original CX material was modified by	q _{eCX} = 32.4 mg g ⁻¹ q _{eCXN} = 33.8 mg g ⁻¹

[51]	magnetic nanoparticles coated zeolite (MNCZ)	<p>V = 25ml</p> <p>[C₀] = 100 µg/l T = 31°C Carbone dose= 1g l-1 pH = 7 contact time=300 min</p>	<p>liquid phase, thermal and hydrothermal resulting in this production of 3 additional materials:</p> <ul style="list-style-type: none"> - CX with Sulfuric acid resulting in the CXS material -CX with Nitric acid resulting in the CXN material -CXN with Urea solution (1mol l-1) resulting in the CXNUT material <p>0.5 g of zeolite was mixed with a 6.1 g FeCl₃.6H₂O and 4.2 g FeSO₄.7H₂O and then dissolved in 100 mL using ultrasonication, as well as, the pH of the mixture was adjusted to 10.0 using 0.1 M NaOH. The mixture was agitated in a rotary shaker at 160 rpm for 24 h. Thereafter 25 mL of the 6.5 M NaOH was then slowly added and mixed with the above solution. The system was mixed for an hour following the addition of NaOH. The formed black precipitates were washed with ultrapure water several times; an external magnetic field was used to enhance the washing process. This procedure leads to form α-Fe₃O₄-Zeolite anoparticles with a size of around 10–20 nm. The α-Fe₃O₄-Zeolite was then oxidized at 300°C for 3 h to obtain γ-Fe₂O₃-Zeolite (MNCZ).</p>	<p>q_{eCXNUT}= 38.7 mg g⁻¹ q_{eCXS}= 40.2 mg g⁻¹</p> <p>q_e= 95.56 mg g⁻¹</p>
[52]	the tissues of Cyclamen persicum tubers	<p>[C₀] = 70 mg/l T = 25°C Carbone dose = 0.7g pH = 4 contact time=120 min</p>	<p>the tissues of Cyclamen persicum tubers was prepared by the physical and chemical activation with ZnCl₂, H₃PO₄ and KOH</p>	<p>q_{max}= 22.22 mg/g</p>

[43] studied the adsorption of diclofenac sodium on Isabel grape bagasse as adsorbent; the results showed that the percentage of elimination was from 16.4 % to 22.8 % and independent from the initial concentration of diclofenac sodium. The kinetics of adsorption have follows the model of pseudo second order and isotherms were presented by the model of freundlich, study thermodynamics showed that the process of adsorption was produced by an exothermic process accompanied with a reduction in the randomness at the solid/solution interface. [44] Also studied adsorption of diclofenac onto a low-cost activated carbon prepared from olive-waste cakes and the influence of reaction parameters such as pH and temperature. In this study the adsorption process also followed the pseudo-second-order kinetic and the increase of the pH of the solution influenced inversely on the rate of adsorption, on the other hand the temperature had no effect on the process. [53] Investigated the adsorption of diclofenac sodium onto three different commercial activated carbons (one granular carbon denoted C1 and two powdered activated carbon (denoted C2 and C3) possessing ~651, 1470 and 913 m²g⁻¹ surface areas, respectively at 303 K using batch adsorption experiments.

The characteristics of these materials were estimated; their textural, surface features and points of zero charge and the obtained results enabled to estimate the possibility to use the

activated carbons in the removal of diclofenac by adsorption. [45] Estimated the mechanism of adsorption of diclofenac on various types of adsorbents with mesoporous silicates and the commercial activated carbon, the adsorption process followed the pseudo-second-order kinetic and the adsorption isotherm was best described by a linear isotherm model. The modification of the hexagonal mesoporous silicates adsorbent (HMS) with mercapto functional caused an increase of the adsorption capacity and they explained this increase through improved hydrogen bonding between the hydrophobic interactions. [74] studied the adsorption of many pharmaceutical residues among them the anionic compound; diclofenac on mesoporous silica SBA-15 and its postfunctionalized counterparts with hydroxymethyl (HM-SBA-15), aminopropyl (AP-SBA-15) and trimethylsilyl (TMS-SBA-15) and found that its adsorption rate was not significant at pH = 5.5 on SBA-15, these results are similar to those of AP-SBA-15. For HM-SBA-15 the adsorption rate is improving slightly and with TMS-SBA-15 the rate exceeded 70%.

4. 2. Biodegradation of diclofenac

Biodegradation is the most commonly used to eliminate persistent pharmaceutical residues process. This process breaks down and mineralizes these pollutants by either bacteria or fungi in simpler compounds [149,150].

[68] investigated the transformation of diclofenac by pellets of *Phanerochaete chrysosporium* in fed-batch bioreactors operating under continuous air supply or periodic pulsation of oxygen. The performance of the fungal reactors was steady over a 30-day treatment and the effect of oxygen pulses on the pellet morphology was evidenced. Complete elimination of diclofenac was achieved in the aerated and the oxygenated reactors, even with a fast oxidation rate in the presence of oxygen (77% after 2 h), reaching a total removal after 23 h. [70] Assessed the degradation of diclofenac sodium using the white-rot fungus *Trametes versicolor*. The use of *T. versicolor* pellets to degrade diclofenac led to an unusual fast degradation rate at concentrations in the range of $\mu\text{g/l}$ to mg/l in a defined liquid medium. In vivo and in vitro experiments using the cytochrome P450 inhibitor 1-aminobenzotriazole and purified laccase, respectively, suggested at least two different mechanisms employed by *T. versicolor* to initiate diclofenac degradation.

Two hydroxylated metabolites, 4-hydroxydiclofenac and 5-hydroxydiclofenac, were structurally elucidated by nuclear magnetic resonance as degradation intermediates in fungal cultures spiked with diclofenac. Both parent compound and intermediates disappeared after 24 h leading to a decrease in ecotoxicity calculated by the Microtox test. Laccase-catalyzed transformation of diclofenac led to the formation of 4-(2,6-dichlorophenylamino)-1,3-benzenedimethanol, which was not detected in in vivo experiments probably due to the low laccase activity levels observed through the first hours of incubation. Figure 1 shows the structure of diclofenac and its metabolites.

Biodegradation is considered the most important process for eliminating the majority of pharmaceuticals [68]. Sewage sludge is divided into two types; primary sludge after primary treatment [153] and secondary sludge after biological secondary treatments [23,24,76] in which microorganisms play a key role in removal of organic contaminants [4] and of which activated sludge Process (CAS) [26, 66] and membrane bioreactors (MBR) are the most frequently used [77] under aerobic or anaerobic condition [151].

Diclofenac is a polar pharmaceutical compound mostly used as the sodium salt diclofenac-Na in human and veterinary medicine to reduce inflammation and pain

[88,94,102]. Many studies have evaluated the biodegradation of diclofenac by conventional activated sludge and membrane bioreactors.

Diclofenac biodegradability in activated sludge and the membrane bioreactor process is influenced by various factors [149]; namely physicochemical properties of diclofenac and the operational parameters of wastewater treatment process including temperature, pH, concentration of biomass, hydraulic retention time (HRT), sludge retention time (SRT), biodegradation kinetics (K_{biol}) and redox conditions [79,130,132] and hydrophobic interactions [146]

Sorption of pharmaceutical residues on solid depends on the hydrophobicity of the molecule to be eliminated according to its coefficient octanol-water K_{ow} where when $\log K_{\text{ow}} < 2.5$ sorption is considered low and when $\log K_{\text{ow}}$ is between 2.5 and 4 is estimated an average sorption and a high sorption when $\log K_{\text{ow}} > 4$ [131,154]. Certainly, a long retention time (HRT) allows a very good interaction between diclofenac and polluted water [133].

4. 2. 1. Conventional activated sludge

In conventional activated sludge (CAS) the aeration tank and the final clarifier form one process unit. The separation of treated sewage and sludge happens in the clarifier via sedimentation. Therefore the capability to sediment is an important selection criterion. The biomass concentration in the mixed liquor is limited by the capacity of the clarifier [88].

Microbial degradation of diclofenac by activated sludge does not lead to its complete elimination but produces other metabolites which are also considered such as pollutants [139], these authors investigated the repartition of diclofenac in activated sludge, they detected 7 diclofenac transformation products among them 1-(2,6-dichlorophenyl)-1,3-dihydro-2H-indol-2-one and 2,3-dichloro-N-(phenyl)aniline.

Activated sludge is the most used in the biological treatment and the rate of elimination of diclofenac by this process was insignificant and does not exceed 50 % [122, 129 and 130]. [86] Confirmed this results when they investigated the performance of activated sludge to remove diclofenac and the results showed that this drug was not completely eliminated ($21.8 \pm 28.5\%$). These results are in accordance with [108] who demonstrated that the removal of diclofenac was 48%. [24] studied the elimination of diclofenac by activated sludge in three WWTP and they showed that the removal rate was 60% for activated sludge presented by tertiary treatment in WWTP1, while for the other two stations the removal rate was $< 25\%$.

[75] Studied the biodegradability of the diclofenac on two types of sludge; sterilized and activated. Results of batch adsorption experiments via sterilized sludge showed that the removal efficiency was 40.1% at 6 hours and 19.7% for activated sludge where the contributions of sludge adsorption and biodegradation were 14.9% at 6 hours and 4.8%, respectively, The authors suggested that this difference in removal efficiency by sterilized and activated sludge is due probably by the sterilizing effect of the activated sludge which can be caused changes in the characteristics of the sludge and improves the elimination of diclofenac and also suggested that the structure of diclofenac (the electronic withdrawing groups) namely its functional groups (amine, halogen, and carboxylic groups) causes the resistance of diclofenac in its biodegradation.

[124] found that the biodegradation kinetics of diclofenac with activated sludge follows the pseudo first order model with a 0.87% correlation coefficient (R^2). These authors also confirmed the persistence of diclofenac to biodegradation. [125] investigated the biodegradation of diclofenac by nitrifying activated sludge and established the no removal of

diclofenac with this process, these results are consistent with those of [126] who have studied the removal of diclofenac by nitrifying activated sludge in large scale (WWTP) and small scale (laboratory level) at a temperature of 12 °C, the authors found that diclofenac is not biodegradable in large scale and the reaction rate constant (k') and biodegradation constant (k_{biol}) were 1.7 d⁻¹ and 0.6 l g_{ss}⁻¹d⁻¹, respectively in WWTP. In laboratory scale the biodegradation rate showed a difference between the reactors where with 10 µg / l of diclofenac the biodegradation rate was 86% in the reactor 1, whereas in the other reactors was 90%. The authors explained this difference by the amount of active nitrifying bacteria in the reactors which was influenced by temperature.

4. 2. 2. Membranes bioreactors

In membrane bioreactor the clarifier used in activated sludge is replaced by a membrane [85]. So, the plant can be operated at higher biomass concentrations resulting in smaller plant sizes [87,104]. The most essential benefit of MBR is the complete retention of suspended solids, thus decreasing emissions to the dissolved fractions [103]. The sorption can be estimated by the K_d value which is the ratio between the concentration of compound in the solid and in the aqueous phase in equilibrium conditions [121], this parameter indicates the sorption efficiency of the sludge, wherein when the K_d value is less than 500 L/kg, the efficiency of the sorption is considered negligible [104].

Recently, the membrane bioreactor process has become the most effective process for the removal of diclofenac compared to activated sludge [86], where the authors have demonstrated a low eliminating for diclofenac by the activated sludge process with a rate of (21.8 ±28.5%), as against the disposal of this waste was 65% for the membrane bioreactor process. On the other hand anaerobic MBR system has not shown an effective elimination rate for diclofenac (<10%) [127], these results are consistent with those of [71] these authors worked on a process of anaerobic / anoxic / aerobic membrane bioreactor in a large scale and observed removal rate below 20% for diclofenac.

In contrast in another study [90] where they used a anaerobic system by anaerobic fluidized membrane bioreactor (AFMBR) using granular activated carbon (GAC) as carrier medium, 78% was the removal rate of diclofenac and the elimination rate of COD was 95% at total HRT of 5 h. It can be concluded that the use of GAC helps eliminate diclofenac in the anaerobic system.

[72] investigated the degradation of diclofenac by a white-rot fungus-augmented membrane bioreactor (MBR) with and without addition of a redox mediator (1-hydroxy benzotriazole, HBT); the results showed that the addition of a redox mediator (1-hydroxy benzotriazole, HBT) improve the removal efficiency of diclofenac from 70% to 95%. In non-sterile conditions studied by [138] the removal rate of diclofenac was 55% in an MBR system with a hydraulic retention time of two days. [134] have shown that increasing the biomass concentration in the MBR system due to the long retention time favors the elimination of diclofenac and can be influenced by microbial activity according to [135]. These authors found that the reduction of microbial activity in the MBR system induces a decrease in the rate of elimination of diclofenac which indicates that the latter is eliminated by biological degradation and not by the sorption process only. [136] investigated the elimination of diclofenac by MBR under different temperature (10-45 °C) in laboratory-scale and demonstrated that the temperature has an effect on the microbial activity. The results showed that the maximum and minimum removal rate were at 10 °C and 20 °C, respectively.

The full-scale MBR showed significantly better removal of hydrophilic compound ($\log D < 3$) [141]; they explained the greater removal by the existence of preand post-anoxic tanks and the combination of aerobic zones with different levels of DO (Dissolved Oxygen) relative to a pre-anoxic and one aerobic tank in the pilot MBR.

The molecular features of diclofenac can affect the rate of his elimination by membrane bioreactor on the scale of laboratory [77]. In this study, the authors observed a very low rate of elimination (17 %) and they justified this behavior of diclofenac in the MBR system by the presence of chlorine group and also the electron withdrawing functional groups generates an electron deficiency and thus renders the drug less susceptible to oxidative catabolism.

As regards the influence of pH, [137] explain the relationship between physicochemical properties of diclofenac and his efficiency removal by a laboratory scale MBR at mixed liquor pH values of 5, 6, 7, 8, and 9. At pH 5 the rate of removal was highest; this result is due to the physicochemical properties of this ionisable compound under acidic conditions wherein the pK_a value is 4.15 [30] so at pH 5 diclofenac is neutral which makes it a hydrophobic compound. The change in pH causes a variation in the hydrophobicity of diclofenac what gives us a $\log D$. The increase of the made pH decreases $\log D$ and consequently the hydrophobicity of diclofenac what makes decrease also the efficiency elimination.

5. EFFECT OF DICLOFENAC ON BACTERIAL COMMUNITY

Sludge from sewage treatment plants is rich in microorganisms responsible for removing contaminants. Consequently, these micro pollutants can have an impact on these bacterial communities and affect the efficiency of biological treatment, namely activated sludge and membrane bioreactor process [142-145 and 147]. These effects are studied by tests of inhibition of development of the bacterial culture [155].

[152] studied the effect of four pharmaceuticals residues (Ketoprofen, naproxen, carbamazepine and gemfibrozil) on the ammonia oxidizing bacterium *Nitrosomonas europaea* at concentrations of 1 and 10 μM . They suggested that the presence of these pharmaceutical residues can affect the activity of the bacterial community and the elimination of nitrogen. [148] used the molecular-based method, fluorescent in situ hybridisation (FISH) to study the effect of organic loads on the structure of the bacterial community and estimated the stability of the anaerobic reactor by measuring the rate of elimination of DCO, the results showed that the rate of elimination of DCO decrease with the increase of the organic load in the reactor and a change was observed in the populations microbial.

Only a few studies evaluated effects of diclofenac on the bacterial community in activated sludge or membrane bioreactor system. Diclofenac can affect bacterial community used for its removal in the activated sludge process [140]; these authors demonstrated that diclofenac affects the structure of the sensitive nitrite-oxidizing bacterial community. [129] found that exposure of microbial community used in the activated sludge treatment causes a strong influence on this community after two months of exposure. This influence on the microbial diversity may affect the wastewater treatment plants. [128] studied this effect in laboratory scale and found that the increasing of concentration of pharmaceutical residues among them diclofenac caused a larger structural divergence on bacterial population and reduced bacterial diversity may affect the essential functions of activated sludge wastewater

treatment systems compared to the reactor which was operated without the addition of pharmaceutical residues.

The abundance of the methanogenic populations on anaerobic digestion (AD) process with regard to the pharmaceutical residue; diclofenac was studied by [156]. The authors compared the effect of diclofenac on biomass using a mesophilic wastewater treatment plant sludge-based (SI) and a thermophilic manure-based inoculum (MI) and they found that, SI inoculum were more resistant to the effect of diclofenac compared to the MI inoculum as regards the production of methane then the increase in biomass caused a decrease of this production until 31% and provoke an increase of the effect of diclofenac. This study demonstrated that hydrogenotrophic methanogens are more resistant than methanogens acetoclastic to the inhibitory effect of diclofenac via the method city by [148].

6. CONCLUSIONS

Diclofenac is the most pharmaceutical residue detected in the environment following its extensive human and veterinary use; it may be outcome of drug manufacturing plants and discharges from hospitals and domestic. This anti-inflammatory drug residue pollutes the aquatic environment and cause harmful effects on the human being, flora and fauna of the ecosystem.

The elimination of diclofenac was demonstrated to large and small scale with two process treatment of polluted water; physicochemical process such as activated carbon adsorption thanks to its advantages economic and biological process or biodegradation using conventional activated sludge (CAS) and membrane bioreactor (MBR) where the microorganisms are responsible for its degradation. Discussion of studies and publications used in this review demonstrated that biodegradation by membrane bioreactor is the most effective process for the removal of a resistant and persistent contaminant such as diclofenac with complete retention of suspended solids, thus reducing emissions to the dissolved fractions, but this removal may be influenced by factors such as the HRT where the increase increases the time of the contact of water with biomass and improves elimination of diclofenac and even an increase in biomass and even the increase in biomass can promote this removal. To see the efficiency of the biological treatment we studied the relationship between diclofenac and the bacterial community of sludge. After this review we find that there is not a lot of research concerned with the effect of diclofenac on biomass but little found in the literature allows us to conclude that the DCF has an influence on the growth of the bacteria sludge and consequently on their diversity.

In summary, to improve biodegradation diclofenac we need to deepen and develop methods to study the impact of pharmaceutical residues as diclofenac on bacterial community of sludge to understand the process of biodegradation and therefore improvement of its removal.

References

[1] Comparison of different removal techniques for selected pharmaceuticals. Andrea Vona, Francesco di Martino, Jorge Garcia-Ivars, Yolanda Picó, José-Antonio Mendoza-Roca, Maria-Isabel Iborra-Clar. *Journal of Water Process Engineering* 5 (2015) 48-57

- [2] Adsorption thermodynamics and kinetics of ranitidine hydrochloride onto superheated steam activated carbon derived from mung bean husk. Sandip Mondal, Keka Sinha, Kaustav Aikat, Gopinath Halder. *Journal of Environmental Chemical Engineering* 3 (2015) 187-195
- [3] Selective removal of diclofenac from contaminated water using molecularly imprinted polymer microspheres. Chao-meng Dai, Sven-Uwe Geissen, Ya-lei Zhang, Yong-jun Zhang, Xue-fei Zhou. *Environmental Pollution* 159 (2011) 1660-1666
- [4] Pharmaceuticals as emerging contaminants and their removal from water. A review José Rivera-Utrilla, Manuel Sánchez-Polo, María Ángeles Ferro-García, Gonzalo Prados-Joya, Raúl Ocampo-Pérez. *Chemosphere* 93 (2013) 1268-1287
- [5] Degradation of drugs in water with advanced oxidation processes and ozone. Maria Quero-Pastor, Alvaro Valenzuela, Jose M. Quiroga, Asuncion Acevedo. *Journal of Environmental Management* 137 (2014) 197-203
- [6] Removal of ibuprofen and its transformation products: Experimental and simulation studies. N. Collado, G. Buttiglieri, L. Ferrando-Climent, S. Rodriguez-Mozaz, J. Comas, I. Rodriguez-Roda, D. Barceló. *Science of the Total Environment* 433 (2012) 296-301
- [7] Production of adsorbents by pyrolysis of paper mill sludge and application on the removal of citalopram from water. Vânia Calisto, Catarina I.A. Ferreira, Sérgio M. Santos, Maria Victoria Gil, Valdemar I. Esteves, Marta Otero. *Bioresource Technology* 166 (2014) 335-344
- [8] Occurrence, fate and ecotoxicological assessment of pharmaceutically active compounds in wastewater and sludge from wastewater treatment plants in Chongqing, the Three Gorges Reservoir Area. Qing Yan, Xu Gao, You-Peng Chen, Xu-Ya Peng, Yi-Xin Zhang, Xiu-Mei Gan, Cheng-Fang Zi, Jin-Song Guo. *Science of the Total Environment* 470-471 (2014) 618-630
- [9] Selective removal of acidic pharmaceuticals from contaminated lake water using multi-templates molecularly imprinted polymer. Chao-Meng Dai, Juan Zhang, Ya-Lei Zhang, Xue-Fei Zhou, Yan-Ping Duan, Shu-Guang Liu. *Chemical Engineering Journal* 211-212 (2012) 302-309
- [10] Degradation of paracetamol by catalytic wet air oxidation and sequential adsorption – Catalytic wet air oxidation on activated carbons. I. Quesada-Penate, C. Julcour-Lebigue, U.J. Jáuregui-Haza, A., M. Wilhelm, H. Delmas. *Journal of Hazardous Materials* 221-222 (2014) 131-138
- [11] Occurrence of acidic pharmaceuticals and personal care products in Turia River Basin: From waste to drinking water. Eric Carmona, Vicente Andreu, Yolanda Picó. *Science of the Total Environment* 484 (2014) 53-63.
- [12] Rapid detection of multiple class pharmaceuticals in both municipal wastewater and sludge with ultra high performance liquid chromatography tandem mass spectrometry. Xiangjuan Yuan, Zhimin Qiang, Weiwei Ben, Bing Zhu, Junxin Liu. *Journal of Environmental Sciences* 26 (2014) 1949-1959
- [13] Mechanism of paracetamol removal by vegetable wastes: The contribution of π - π interactions, hydrogen bonding and hydrophobic effect. Isabel Villaescusa, Núria Fiol, Jordi Poch, Antonio Bianchi, Carla Bazzicalupi. *Desalination* 270 (2011) 135-142

- [14] Occurrence and risk assessment of acidic pharmaceuticals in the Yellow River, Hai River and Liao River of north China. Li Wang, Guang-Guo Ying, Jian-Liang Zhao, Xiao-Bing Yang, Feng Chen, Ran Tao, Shan Liu, Li-Jun Zhou. *Science of the Total Environment* 408 (2010) 3139-3147
- [15] Estimating the toxicities of organic chemicals in activated sludge process. Alan R. Katritzky, Kalev Kasemets, Svetoslav Slavov, Maksim Radzvilovits, Kaido Tamm, Mati Karelson. *Water research* 44 (2010) 2451-2460
- [16] Ecotoxicological aspects related to the presence of pharmaceuticals in the aquatic environment. Lúcia H.M.L.M. Santos, A.N. Araújo, Adriano Fachini, A. Pena, C. Delerue-Matos, M.C.B.S.M. Montenegro. *Journal of Hazardous Materials* 175 (2010) 45-95
- [17] Biological removal of pharmaceutical and personal care products by a mixed microbial culture: Sorption, desorption and biodegradation. Ioanna A. Vasiliadou, Raul Molina, Fernando Martínez, Juan A. Melero. *Biochemical Engineering Journal* 81 (2013) 108-119
- [18] Aqueous adsorption and removal of organic contaminants by carbon nanotubes. Review. Jin-Gang Yu, Xiu-Hui Zhao, Hua Yang, Xiao-Hong Chen, Qiaoqin Yang, Lin-Yan Yu, Jian-Hui Jiang, Xiao-Qing Chen. *Science of the Total Environment* 482-483 (2014) 241-251
- [19] Environmental friendly technology for the removal of pharmaceutical contaminants from wastewaters using modified chitosan adsorbents. George Z. Kyzas, Margaritis Kostoglou, Nikolaos K. Lazaridis, Dimitra A. Lambropoulou. *Chemical Engineering Journal* 222 (2013) 248-258
- [20] Competitive adsorption of ibuprofen and amoxicillin mixtures from aqueous solution on activated carbons. Hayet Mansouri, Rocio J. Carmona, Alicia Gomis-Berenguer, Souad Souissi-Najar. *Journal of Colloid and Interface Science* (2014).
- [21] Transition metal modified and partially calcined inorganic–organic pillared clays for the adsorption of salicylic acid, clofibric acid, carbamazepine, and caffeine from water. Wilman A. Cabrera-Lafaurie, Félix R. Román, Arturo J. Hernández-Maldonado. *Journal of Colloid and Interface Science* 386 (2012) 381-391
- [22] Removal of caffeine and diclofenac on activated carbon in fixed bed column. J.L. Sotelo, A. Rodríguez, S. Álvarez, J. García. *Chemical engineering research and design* 90 (2012) 967-974
- [23] Occurrence and removal of PPCPs in municipal and hospital wastewaters in Greece. Christina I. Kosma, Dimitra A. Lambropoulou, Triantafyllos A. Albanis. *Journal of Hazardous Materials* 179 (2010) 804-817
- [24] Occurrence, partition and removal of pharmaceuticals in sewage water and sludge during wastewater treatment. Aleksandra Jelic, Meritxell Gros, Antoni Ginebreda, Raquel Cespedes-Sanchez. *Water research* 45 (2011) 1165-1176
- [25] Occurrence of pharmaceutical compounds in urban wastewater: Removal, mass load and environmental risk after a secondary treatment-A review. P. Verlicchi, M. Al Aukidy, E. Zambello. *Science of the Total Environment* 429 (2012) 123-155

- [26] Adsorption of three pharmaceuticals on two magnetic ion-exchange resins. Miao Jiang, Weiben Yang, Ziwei Zhang, Zhen Yang, Yuping Wang. *Journal of Environmental Sciences* (2015)
- [27] Investigation of adsorption and release of diclofenac sodium by modified zeolites composites. Danina Krajišnik, Aleksandra Daković, Anđelija Malenović, Maja Milojević-Rakić, Vera Dondur, Željka Radulović, Jela Milić. *Applied Clay Science* 83-84 (2013) 322-326
- [28] The non-steroidal anti-inflammatory drug diclofenac is readily biodegradable in agricultural soils. Abdul Jabbar Al-Rajab, Lyne Sabourin, David R. Lapen, Edward Topp. *Science of the Total Environment* 409 (2010) 78–82
- [29] Degradation of the drug diclofenac in water by sonolysis in presence of catalysts. J. Hartmann, P. Bartels, U. Mau, M. Witter, W.v. Tumpling. *Chemosphere* 70 (2008) 453-461
- [30] A pilot-scale hybrid municipal wastewater reclamation system using combined coagulation and disk filtration, ultrafiltration, and reverse osmosis: Removal of nutrients and micropollutants, and characterization of membrane foulants. Kangmin Chon, Jaeweon Cho, Ho Kyong Shon. *Bioresource Technology* 141 (2013) 109-116
- [31] Degradation of diclofenac sodium using combined processes based on hydrodynamic cavitation and heterogeneous photocatalysis. Manisha V. Bagal, Parag R. Gogate. *Ultrasonics Sonochemistry* 21 (2014) 1035-1043
- [32] Carbamazepine and diclofenac: Removal in wastewater treatment plants and occurrence in water bodies. Yongjun Zhang, Sven-Uwe Geißen, Carmen Gal. *Chemosphere* 73 (2008) 1151-1161
- [33] Formation of diclofenac and sulfamethoxazole reversible transformation products in aquifer material under denitrifying conditions: Batch experiments. Manuela Barbieri, Jesús Carrera, Carlos Ayora, Xavier Sanchez-Vila, Tobias Licha, Karsten Nödler, Victoria Osorio, Sandra Pérez, Marianne Köck-Schulmeyer, Miren López de Alda, Damià Barceló. *Science of the Total Environment* 426 (2012) 256-263
- [34] Occurrence and human health risk of wastewater-derived pharmaceuticals in a drinking water source for Shanghai, East China. Zhi-Hao Wen, Ling Chen, Xiang-Zhou Meng, Yan-Ping Duan, Zeng-Sheng Zhang, Eddy Y. Zeng. *Science of the Total Environment* 490 (2014) 987–993
- [35] Pharmaceutical residues in wastewater treatment works effluents and their impact on receiving river water. J.L. Zhou, Z.L. Zhang, E. Banks, D. Grover, J.Q. Jiang. *Journal of Hazardous Materials* 166 (2009) 655-661
- [36] The occurrence and fate of anti-inflammatory and analgesic pharmaceuticals in sewage and fresh water: Treatability by conventional and non-conventional processes. Asu Ziylan, Nilsun H. Ince. *Journal of Hazardous Materials* 187 (2011) 24-36
- [37] Adsorption of basic dyes on activated carbon prepared from *Polygonum orientale* Linn: Equilibrium, kinetic and thermodynamic studies. Li Wang, Jian Zhang, Ran Zhao, Cong Li, Ye Li, Chenglu Zhang. *Desalination* 254 (2010) 68-74

- [38] Adsorption of Diclofenac Sodium from Aqueous Solutions on Expanded Graphite. M. D. Vedenyapina, D. A. Borisova, A. P. Simakova, L. P. Proshina, and A. A. Vedenyapin. ISSN 03615219, *Solid Fuel Chemistry*, 2013, Vol. 47, No. 1, pp. 59–63. © Allerton Press, Inc., 2013.
- [39] Adsorption characteristics of selected pharmaceuticals and an endocrine disrupting compound—Naproxen, carbamazepine and nonylphenol—on activated carbon. Zirui Yu, Sigrid Peldszus, Peter M. Huck. *Water Research* 42 (2008) 2873-882
- [40] The removal of endocrine disrupting compounds, pharmaceutically activated compounds and cyanobacterial toxins during drinking water preparation using activated carbon-A review. Luis F. Delgado, Philippe Charles, Karl Glucina, Catherine Morlay. *Science of the Total Environment* 435-436 (2012) 509-525
- [41] Waste-derived activated carbons for removal of ibuprofen from solution: Role of surface chemistry and pore structure. Ana S. Mestre, João Pires, José M.F. Nogueira, Jose B. Parra, Ana P. Carvalho, Conchi O. Ania. *Bioresource Technology* 100 (2009) 1720-1726
- [42] Activated carbons from sisal waste by chemical activation with K_2CO_3 : Kinetics of paracetamol and ibuprofen removal from aqueous solution. Ana S. Mestre, Ana S. Bexiga, Margarida Proença, Marta Andrade, Moisés L. Pinto, Inês Matos, Isabel M. Fonseca, Ana P. Carvalho. *Bioresource Technology* 102 (2011) 8253-8260
- [43] Removal of diclofenac sodium from aqueous solution by Isabel grape bagasse. Márjore Antunes, Valdemar I. Esteves, Régis Guégan, Janaina S. Crespo, Andreia N. Fernandes. Marcelo Giovanela. *Chemical Engineering Journal* 192 (2012) 114-121
- [44] Removal of pharmaceutical compounds by activated carbon prepared from agricultural by-product. R. Baccar, M. Sarrà, J. Bouzid, M. Feki, P. Blánquez. *Chemical Engineering Journal* 211-212 (2012) 310-317
- [45] Mechanistic study of diclofenac and carbamazepine adsorption on functionalized silica-based porous materials. Nakorn Suriyanon, Patiparn Punyapalakul, Chawalit Ngamcharussrivichai. *Chemical Engineering Journal* 214 (2013) 208–218
- [46] Regenerable granular carbon nanotubes/alumina hybrid adsorbents for diclofenac sodium and carbamazepine removal from aqueous solution. Haoran Wei, Shubo Deng*, Qian Huang, Yao Nie, Bin Wang, Jun Huang, Gang Yu. *Water research* (2013) 1-9
- [47] Competitive adsorption of selected non-steroidal anti-inflammatory drugs on activated biochars: Experimental and molecular modeling study. Chanil Jung, Linkel K. Boateng, Joseph R.V. Flora, Jeill Oh, Marcus C. Braswell, Ahjeong Son, Yeomin Yoon. *Chemical Engineering Journal* 264 (2015) 1-9
- [48] Synthesis of carbon xerogels and their application in adsorption studies of caffeine and diclofenac as emerging contaminants. S.Álvarez, R.S. Ribeiro, H.T. Gomes, J.L. Sotelo, J. García. *Chemical engineering research and design* 95 (2015) 229-238
- [49] Adsorption of pharmaceuticals onto trimethylsilylated mesoporous SBA-15. Tung Xuan Bui, Viet Hung Pham, Son Thanh Le, Heechul Choi. *Journal of Hazardous Materials* 254-255 (2013) 345-353

- [50] Synthesis of carbon xerogels and their application in adsorption studies of caffeine and diclofenac as emerging contaminants. S. Álvarez, R.S. Ribeiro, H.T. Gomes, J.L. Sotelo, J. García. *Chemical engineering research and design* (2014).
- [51] Synthesized magnetic nanoparticles coated zeolite for the adsorption of pharmaceutical compounds from aqueous solution using batch and column studies. Tamer Mohamed Salem Attia, Xia Lin Hu, Yin Da Qiang. *Chemosphere* 93 (2013) 2076-2085.
- [52] Adsorption of diclofenac from aqueous solution using Cyclamen persicum tubers based activated carbon (CTAC). Shehdeh Jodeh, Fatima Abdelwahab, Nidal Jaradat, Ismail Warad, Wade Jodeh. *Journal of the Association of Arab Universities for Basic and Applied Sciences* (2015)
- [53] The Adsorption of pharmaceutically active compounds from aqueous solutions onto activated carbons. Vesna Raki, Vladislav Rac, Marija Krmar, Otman Otman, Aline Auroux. *Journal of Hazardous Materials* 282 (2015) 141-149
- [54] Pharmaceutical contamination in residential, industrial, and agricultural waste streams: Risk to aqueous environments in Taiwan. Angela Yu-Chen Lin, Tsung-Hsien Yu, Cheng-Fang Lin. *Chemosphere* 74 (2008) 131-141
- [55] Occurrence of pharmaceuticals in Taiwan's surface waters: Impact of waste streams from hospitals and pharmaceutical production facilities. Angela Yu-Chen Lin, Yu-Ting Tsai. *Science of the Total Environment* 407 (2009) 3793-3802
- [56] Occurrence and distribution of pharmaceuticals in wastewater from households, livestock farms, hospitals and pharmaceutical manufactures. Won-Jin Sim, Ji-Woo Lee, Eung-Sun Lee, Sun-Kyoung Shin, Seung-Ryul Hwang, Jeong-Eun Oh. *Chemosphere* 82 (2011) 179-186
- [57] Occurrence and removal of antibiotics, hormones and several other pharmaceuticals in wastewater treatment plants of the largest industrial city of Korea. Shishir Kumar Behera, Hyeong Woo Kim, Jeong-Eun Oh, Hung-Suck Park. *Science of the Total Environment* 409 (2011) 4351-4360
- [58] Assessment of non-steroidal anti-inflammatory and analgesic pharmaceuticals in seawaters of North of Portugal: Occurrence and environmental risk. Aleksandar Lolić, Paula Paíga, Lúcia H.M.L.M. Santos, Sandra Ramos, Manuela Correia, Cristina Delerue-Matos. *Science of the Total Environment* 508 (2015) 240-250
- [59] Concentration evolution of pharmaceutically active compounds in raw urban and industrial wastewater. Dolores Camacho-Muñoz, Julia Martín, Juan Luis Santos, Irene Aparicio, Esteban Alonso. *Chemosphere* 111 (2014) 70-79
- [60] Presence of pharmaceutically active compounds in Donana Park (Spain) main watersheds. M.D. Camacho-Munoz, J.L. Santos, I. Aparicio, E. Alonso. *Journal of Hazardous Materials* 177 (2010) 1159-1162
- [61] The removal of pharmaceuticals, personal care products, endocrine disruptors and illicit drugs during wastewater treatment and its impact on the quality of receiving waters. Barbara Kasprzyk Hordern, Richard M. Dinsdale, Alan J. Guwy. *Water research* 43 (2009) 363-380

- [62] Investigation of PPCPs in wastewater treatment plants in Greece: Occurrence, removal and environmental risk assessment. Christina I. Kosma, Dimitra A. Lambropoulou, Triantafyllos A. Albanis. *Science of the Total Environment* 466-467 (2014) 421-438
- [63] Occurrence and removal of pharmaceuticals in a municipal sewage treatment system in the south of Sweden. Saioa Zorita, Lennart Mårtensson, Lennart Mathiasson. *Science of the Total Environment* 407 (2009) 2760-2770
- [64] Determination of 81 pharmaceutical drugs by high performance liquid chromatography coupled to mass spectrometry with hybrid triple quadrupole–linear ion trap in different types of water in Serbia. Mira Petrović, Biljana Škrbić, Jelena Živančev, Laura Ferrando-Climent, Damia Barcelo. *Science of the Total Environment* 468-469 (2014) 415-428
- [65] Occurrence and fate of pharmaceutically active compounds in the largest municipal wastewater treatment plant in Southwest China: Mass balance analysis and consumption back-calculated model. Qing Yan, Xu Gao, Lei Huang, Xiu-Mei Gan, Yi-Xin Zhang, You-Peng Chen, Xu-Ya Peng. *Chemosphere* 99 (2014) 160-170
- [66] Occurrence and fate of quinolone and fluoroquinolone antibiotics in a municipal sewage treatment plant. Ai Jia, Yi Wan, Yang Xiao, Jianying Hu. *Water research* 46 (2012) 387e394
- [67] Environmental impact of pharmaceuticals from Portuguese wastewaters: geographical and seasonal occurrence, removal and risk assessment. André M.P.T. Pereira, Liliana J.G. Silva, Leonor M. Meisel, Celeste M. Lino. *Environmental Research* 136 (2015) 108-119
- [68] Biotransformation of three pharmaceutical active compounds by the fungus *Phanerochaete chrysosporium* in a fed batch stirred reactor under air and oxygen supply. A. I. Rodarte-Morales, M. T. Moreira, G. Feijoo, J. M. Lema. *Biodegradation* (2012) 23: 145-156
- [69] L. Berthod, et al., A solid-phase extraction approach for the identification of pharmaceutical–sludge adsorption mechanisms, *J. Pharm. Anal.* (2013), <http://dx.doi.org/10.1016/j.jpha.2013.08.003>
- [70] Degradation of the drug sodium diclofenac by *Trametes versicolor* pellets and identification of some intermediates by NMR. Ernest Marco-Urrea, Miriam Pérez-Trujillo, Carles Cruz-Morató, Gloria Caminal, Teresa Vicent. *Journal of Hazardous Materials* 176 (2010) 836-842
- [71] Elimination and fate of selected micro-organic pollutants in a full-scale anaerobic/anoxic/aerobic process combined with membrane bioreactor for municipal wastewater reclamation. Wenchao Xue, Chunying Wu, Kang Xiao, Xia Huang, Haidong Zhou, Hiroshi Tsuno, Hiroaki Tanaka. *Water research* 44 (2010) 5999-6010
- [72] Removal of trace organic contaminants by an MBR comprising a mixed culture of bacteria and white-rot fungi. Luong N. Nguyen, Faisal I. Hai, Shufan Yang, Jinguo Kang, Frederic D.L. Leusch, Felicity Roddick, William E. Price, Long D. Nghiem. *Bioresour. Technology* 148 (2013) 234-241
- [73] Occurrence and removal of pharmaceuticals in wastewater treatment plants at the Spanish Mediterranean area of Valencia. Emma Gracia-Lor, Juan V. Sancho, Roque Serrano, Félix Hernández. *Chemosphere* 87 (2012) 453-462

- [74] Organically functionalized mesoporous SBA-15 as sorbents for removal of selected Pharmaceuticals from water. Tung Xuan Bui, Seo-Young Kang, Sang-Hyup Lee, Heechul Choi. *Journal of Hazardous Materials* 193 (2011) 156-163
- [75] Contribution of sludge adsorption and biodegradation to the removal of five pharmaceuticals in a submerged membrane bioreactor. Huiju Fan, Jing Li, Liqiu Zhang, Li Feng. *Biochemical Engineering Journal* 88 (2014) 101-107
- [76] Microbial community related to lysozyme digestion process for boosting waste activated sludge biodegradability. Xiao-Dong Xin, Jun-Guo He, Wei Qiu, Jian Tang, Tian-Tian Liu. *Bioresource Technology* 175 (2015) 112-119
- [77] Removal of trace organics by MBR treatment: The role of molecular properties. Nichanan Tadkaew, Faisal I. Hai, James A. McDonald, Stuart J. Khan, Long D. Nghiem. *Water Research* 45 (2011) 2439-2451
- [78] Occurrence and behaviour of 105 active pharmaceutical ingredients in sewage waters of a municipal sewer collection system. Richard H. Lindberg, Marcus Ostman, Ulrika Olofsson, Roman Grabic, Jerker Fick. *Water research* 58 (2014) 221-229
- [79] Removal of pharmaceuticals during wastewater treatment and environmental risk assessment using hazard indexes. Meritxell Gros, Mira Petrović, Antoni Ginebreda, Damià Barceló. *Environment International* 36 (2010) 15-26
- [80] Diclofenac photodegradation under simulated sunlight: Effect of different forms of nitrogen and Kinetics. Nan Zhang, Guoguang Liu, Haijin Liu, Yingling Wang, Zhanwei He, Gang Wang. *Journal of Hazardous Materials* 192 (2011) 411-418
- [81] Role of cetyltrimethyl ammonium bromide, crystal violet and humic acid in the degradation of diclofenac under simulated sunlight irradiation. Zhang Nan, Liu Hai Jin, Liu Guo Guang, Wang Ying Ling, Li Tong, Wang Gang. *Science China Chemistry* Vol. 55 No.12: 2610-2616
- [82] Adsorption characteristics of diclofenac and sulfamethoxazole to graphene oxide in aqueous solution. Seung-Woo Nam, Chanil Jung, Hang Li, Miao Yu, Joseph R.V. Flora, Linkel K. Boateng, Namguk Her, Kyung-Duk Zoh, Yeomin Yoon. *Chemosphere* 136 (2015) 20-26
- [83] Bio-entrapped membrane reactor and salt marsh sediment membrane bioreactor for the treatment of pharmaceutical wastewater: Treatment performance and microbial communities. Kok Kwang Ng, Xueqing Shi, Yinuo Yao, How Yong Ng. *Bioresource Technology* 171 (2014) 265-273
- [84] Effect of solid retention time on sludge filterability and biomass activity: Long-term experiment on a pilot-scale membrane bioreactor treating municipal wastewater. Gianpaolo Sabia, Marco Ferraris, Alessandro Spagni. *Chemical Engineering Journal* 221 (2013) 176-184
- [85] Membrane bioreactor: Applications and limitations in treating high strength industrial wastewater. Noor Sabrina Ahmad Mutamim, Zainura Zainon Noor, Mohd Ariffin Abu Hassan, Gustaf Olsson, Adhi Yuniarto. *Chemical Engineering Journal* 225 (2013) 109-119

- [86] Fate and distribution of pharmaceuticals in wastewater and sewage sludge of the conventional activated sludge (CAS) and advanced membrane bioreactor (MBR) treatment. Jelena Radjenovic, Mira Petrovic, Damia Barcelo. *Water research* 43 (2009) 831-841
- [87] Membrane bioreactor for treatment of pharmaceutical wastewater containing acetaminophen. Farshid Pajoum Shariati, Mohammad Reza Mehrnia, Bahreh Madadkhah Salmasi, Christelle Wisniewski, Mohammad Hossein Sarrafzadeh, Marc Heran. *Desalination* 250 (2010) 798-800
- [88] Trace Analysis of Pharmaceutical Residues in Wastewater Treatment Plants in Rio de Janeiro, Brazil. Archive of SID. Archive of SID. Aldo Pacheco Ferreira. *Journal of Chemical Health Risks* (2014) 4(1), 1-12
- [89] Biodegradation and removal of pharmaceuticals and personal care products in treatment systems: a review. Kathryn M. Onesios & Jim T. Yu, Edward J. Bouwer. *Biodegradation* (2009) 20: 441-466
- [90] Removal of pharmaceuticals and organic matter from municipal wastewater using two-stage anaerobic fluidized membrane bioreactor. Kasturi Dutta, Ming-Yi Lee, Webber Wei-Po Lai, Chien Hsien Lee, Angela Yu-Chen Lin, Cheng-Fang Lin, Jih-Gaw Lin. *Bioresource Technology* 165 (2014) 42-49
- [91] The treatment of pharmaceutical wastewater using in a submerged membrane bioreactor under different sludge retention times. Yasemin Kaya, Gamze Ersan, Ilda Vergili, Z. Beril Gönder, Gulsum Yilmaz, Nadir Dizge, Coskun Aydiner. *Journal of Membrane Science* 442 (2013) 72-82
- [92] Effects of trophic exposure to dexamethasone and diclofenac in fresh water fish. Izonete Cristina Guiloski, João Luiz Coelho Ribas, Letícia da Silva Pereira, Ana Paula, Perbiche Neves, Helena Cristina, Silva de Assis. *Ecotoxicology and Environmental Safety* 114 (2015) 204-211
- [93] Factory-discharged pharmaceuticals could be a relevant source of aquatic environment contamination: Review of evidence and need for knowledge. Olivier Cardoso, Jean-Marc Porcher, Wilfried Sanchez. *Chemosphere* 115 (2014) 20-30
- [94] Occurrence and fate of bulk organic matter and pharmaceutically active compounds in managed aquifer recharge: A review. Sung Kyu Maeng, Saroj K. Sharma, Karin Lekkerkerker-Teunissen, Gary L. Amy. *Water research* 45 (2011) 3015-3033
- [95] The occurrence of pharmaceuticals, personal care products, endocrine disruptors and illicit drugs in surface water in South Wales, UK. Barbara Kasprzyk-Hordern, Richard M. Dinsdale, Alan J. Guwy. *Water research* 42 (2008) 3498– 3518
- [96] Occurrence and removal of pharmaceuticals and personal care products (PPCPs) in an advanced wastewater reclamation plant. Xin Yang, Riley C. Flowers, Howard S. Weinberg, Philip C. Singer. *Water research* 45 (2011) 5218-5228
- [97] Characterization of Pharmaceuticals and Personal Care products in hospital effluent and waste water influent/effluent by direct-injection LC-MS-MS. Tiago S. Oliveira, Mark Murphy, Nicholas Mendola, Virginia Wong, Doreen Carlson, Linda Waring. *Science of the Total Environment* 518-519 (2015) 459-478

- [98] Occurrence of pharmaceuticals and personal care products and associated environmental risks in the central and lower Yangtze river, China. Chenxi Wu, Xiaolong Huang, Jason D. Witter, Alison L. Spongberg, Kexiong Wang, Ding Wang, Jiantong Liu. *Ecotoxicology and Environmental Safety* 106 (2014) 19-26
- [99] Pharmaceuticals and endocrine disrupting compounds in U.S. drinking water. M.J. Benotti, R.A. Trenholm, B.J. Vanderford, J.C. Holady, B.D. Stanford, S.A. Snyder, *Environ. Sci. Technol.* 43 (2009) 597-603.
- [100] Determination of phenolic endocrine disrupting chemicals and acidic pharmaceuticals in surface water of the Pearl Rivers in South China by gas chromatography-negative chemical ionization-mass spectrometry, J.-L. Zhao, G.-G. Ying, L. Wang, J.-F. Yang, X.-B. Yang, L.-H. Yang, X. *Sci. Total Environ.* 407 (2009) 962-974.
- [101] Occurrence and source apportionment of pharmaceuticals and personal care products in the Beiyun River of Beijing, China. Guohua Dai, Bin Wang, Jun Huang, Rui Dong, Shubo Deng, Gang Yu. *Chemosphere* 119 (2015) 1033-1039
- [102] Pharmaceuticals in the environment: Biodegradation and effects on natural microbial communities. A review. Anna Barra Caracciolo, Edward Topp, Paola Grenni. *Journal of Pharmaceutical and Biomedical Analysis* 106 (2015) 25-36
- [103] Pharmaceutical wastewater treatment by membrane bioreactor process – a case study in southern Taiwan. Chia-Yuan Chang, Jing-Song Chang, Saravanamuthu Vigneswaran, Jaya Kandasamy. *Desalination* 234 (2008) 393-401
- [104] Comparison of removal of pharmaceuticals in MBR and activated sludge systems. Jan Sipma, Begoña Osuna, Neus Collado, Hector Monclús, Giuliana Ferrero, Joaquim Comas, Ignasi Rodriguez-Roda. *Desalination* 250 (2010) 653-659
- [105] Major pharmaceutical residues in wastewater treatment plants and receiving waters in Bangkok, Thailand, and associated ecological risks. S. Tewari, R. Jindal, Y.L. Kho, S. Eo, K. Choi. *Chemosphere* 91 (2013) 697-704
- [106] Thermophilic treatment of bulk drug pharmaceutical industrial wastewaters by using hybrid up flow anaerobic sludge blanket reactor. D. Sreekanth, D. Sivaramakrishna, V. Himabindu, Y. Anjaneyulu. *Bioresource Technology* 100 (2009) 2534-2539
- [107] Influence of residual organic macromolecules produced in biological wastewater treatment processes on removal of pharmaceuticals by NF/RO membranes. Katsuki Kimura, Tomonori Iwase, Shusuke Kita, Yoshimasa Watanabe. *Water research* 43 (2009) 3751-3758
- [108] Removal of pharmaceuticals from wastewater by biological processes, hydrodynamic cavitation and UV treatment. Mojca Zupanc, Tina Kosjek, Martin Petkovšek, Matevz Dular, Boris Kompare, Brane Širok, eljko Blazeka, Ester Heath. *Ultrasonics Sonochemistry* 20 (2013) 1104-1112
- [109] Occurrence and fate of antibiotic, analgesic/anti-inflammatory, and antifungal compounds in five wastewater treatment processes. P. Guerra, M. Kim, A. Shah, M. Alae, S.A. Smyth. *Science of the Total Environment* 473-474 (2014) 235-243

- [110] Occurrence and removal of PPCPs in municipal and hospital wastewaters in Greece. Christina I. Kosma, Dimitra A. Lambropoulou, Triantafyllos A. Albanis. *Journal of Hazardous Materials* 179 (2010) 804-817
- [111] Physiological effects of diclofenac, ibuprofen and propranolol on Baltic Sea blue mussels. Hanna Ericson, Gunnar Thorsén, Linda Kumblad. *Aquatic Toxicology* 99 (2010) 223-231
- [112] Effects of non-steroidal anti-inflammatory drug (NSAID) diclofenac exposure in mussel *Mytilus galloprovincialis*. Maria Gonzalez-Rey, Maria João Bebianno. *Aquatic Toxicology* 148 (2014) 221-230
- [113] Effects of the pharmaceuticals gemfibrozil and diclofenac on the marine mussel (*Mytilus* spp.) and their comparison with standardized toxicity tests. Wiebke Schmidt, Kathleen O'Rourke, Robert Hernan, Brian Quinn. *Marine Pollution Bulletin* 62 (2011) 1389-1395
- [114] Toxicological effects of clofibric acid and diclofenac on plasma thyroid hormones of an Indian major carp, *Cirrhinus mrigala* during short and long-term exposures. Manoharan Saravanan, Jang-Hyun Hur, Narayanasamy Arul, Mathan Ramesh. *Environmental toxicology and pharmacology* 38 (2014) 948-958
- [115] Chronic exposure to diclofenac on two freshwater cladocerans and Japanese medaka. Jinyoung Lee, Kyunghye Ji, Young Lim Kho, Pilje Kim, Kyungho Choi. *Ecotoxicology and Environmental Safety* 74 (2011) 1216-1225
- [116] Toxicity screening of Diclofenac, Propranolol, Sertraline and Simvastatin using *Danio rerio* and *Paracentrotus lividus* embryo bioassays. Silvia Ribeiro, Tiago Torres, Rosário Martins, Miguel M. Santos. *Ecotoxicology and Environmental Safety* 114 (2015) 67-74
- [117] Fish embryo toxicity of carbamazepine, diclofenac and metoprolol. Evert-Jan van den Brandhof, Mark Montforts. *Ecotoxicology and Environmental Safety* 74 (2010) 1862-1866
- [118] The effects of diclofenac on early life stages of common carp (*Cyprinus carpio*). Stanislava Stepanova, Eva Praskova, Lucie Chromcova, Lucie Phalova, Miroslav Prokes, Jana Blahova, Zdenka Svobodova. *Environmental toxicology and pharmacology* 35 (2013) 454-460
- [119] Occurrence of selected pharmaceuticals in the principal sewage treatment plants in Rome (Italy) and in the receiving surface waters. Luisa Patrolecco, Silvio Capri, Nicoletta Ademollo. *Environ Sci Pollut Res* 22 (2015) 5864-5876
- [120] Ecotoxicogenomic assessment of diclofenac toxicity in soil. Guangquan Chen, Michiel W. den Braver, Cornelis A.M. van Gestel, Nico M. van Straalen, Dick Roelofs. *Environmental Pollution* 199 (2015) 253-260
- [121] A solid-phase extraction method for rapidly determining the adsorption coefficient of pharmaceuticals in sewage sludge. Laurence Berthod, Gary Roberts, David C. Whitley, Alan Sharpe, Graham A. Mills. *Water research* 67 (2014) 292-298
- [122] Fate of drugs during wastewater treatment. Bruce Petrie, Ewan J. McAdam, Mark D. Scrimshaw, John N. Lester, Elise Cartmell. *Trends in Analytical chemistry* 49 (2013) 145-159

- [123] Occurrence, transportation, monitoring and treatment of emerging micro-pollutants in waste water — A review from global views. Jia-Qian Jiang, Z. Zhou, V.K. Sharma. *Microchemical Journal* 110 (2013) 292-300
- [124] Active heterotrophic biomass and sludge retention time (SRT) as determining factors for biodegradation kinetics of pharmaceuticals in activated sludge. Marius Majewsky, Tom Gallé, Viviane Yargeau, Klaus Fischer. *Bioresource Technology* 102 (2011) 7415-7421
- [125] Suspended biofilm carrier and activated sludge removal of acidic pharmaceuticals. P. Fala, A. Baillon-Dhumez, H.R. Andersen, A. Ledin, J. la Cour Jansen. *Water research* (2012) 1167-1175
- [126] Biodegradation of ibuprofen, diclofenac and carbamazepine in nitrifying activated sludge under 12 °C temperature conditions. Antonina Kruglova, Pia Ahlgren, Nasti Korhonen, Pirjo Rantanen, Anna Mikola, Riku Vahala. *Science of the Total Environment* 499 (2014) 394-401
- [127] Removal of trace organics by anaerobic membrane bioreactors. Victor M. Monsalvo, James A. McDonald, Stuart J. Khan, Pierre Le-Clech. *Water research* 49 (2014) 103-112
- [128] Influence of pharmaceutical residues on the structure of activated sludge bacterial communities in wastewater treatment bioreactors. Barbara Kraigher, Tina Kosjek, Ester Heath, Boris Kompare, Ines Mandic-Mulec. *Water research* 42 (2008) 4578-4588
- [129] Micropollutant removal by attached and suspended growth in a hybrid biofilm-activated sludge process. P. Fala, P. Longre, J. la Cour Jansen, H. Siegrist, J. Hollender, A. Joss. . *Water research* 47 (2013) 4498-4506
- [130] Mass balance of pharmaceutical and personal care products in a pilot-scale single-sludge system: Influence of T, SRT and recirculation ratio. Sonia Suárez, Rubén Reif, Juan M. Lema, Francisco Omil. *Chemosphere* 89 (2012) 164-171
- [131] Evaluation of a model for the removal of pharmaceuticals, personal care products, and hormones from wastewater. Benjamin D. Blair, Jordan P. Crago, Curtis J. Hedman, Ronan J.F. Treguer, Christopher Magruder, L. Scott Royer, Rebecca D. Klaper. *Science of the Total Environment* 444 (2013) 515-521
- [132] Occurrence and removal of transformation products of PPCPs and illicit drugs in wastewaters: A review. Eleni N. Evgenidou, Ioannis K. Konstantinou, Dimitra A. Lambropoulou. *Science of the Total Environment* 505 (2015) 905-926
- [133] Removal of pharmaceuticals and personal care products in aquatic plant-based systems: A review. Dongqing Zhang, Richard M. Gersberg, Wun Jern Ng, Soon Keat Tan. *Environmental Pollution* 184 (2014) 620-639
- [134] Influence of nitrifying conditions on the biodegradation and sorption of emerging micropollutants. E. Fernandez-Fontaina, F. Omil, J.M. Lema, M. Carballa. *Water research* 46 (2012) 5434-5444
- [135] Influences of solid retention time, nitrification and microbial activity on the attenuation of pharmaceuticals and estrogens in membrane bioreactors. Sung Kyu Maeng, Byeong Gyu Choi, Kyu Tae Lee, Kyung Guen Song. *Water research* 47 (2013) 3151-3162

- [136] Removal of micropollutants by membrane bioreactor under temperature variation. Faisal I. Hai, Karin Tessmer, Luong N. Nguyen, Jinguo Kang, William E. Price, Long D. Nghiem. *Journal of Membrane Science* 383 (2011) 144-151
- [137] Effect of mixed liquor pH on the removal of trace organic contaminants in a membrane bioreactor. Nichanan Tadkaew, Muttucumaru Sivakumar, Stuart J. Khan, James A. McDonald, Long Duc Nghiem. *Bioresource Technology* 101 (2010) 1494-1500
- [138] Removal of bisphenol A and diclofenac by a novel fungal membrane bioreactor operated under non-sterile conditions. Shufan Yang, Faisal I. Hai, Long D. Nghiem, Luong N. Nguyen, Felicity Roddick, William E. Price. *International Biodeterioration & Biodegradation* 85 (2013) 483-490
- [139] Metabolism studies of diclofenac and clofibrac acid in activated sludge bioreactors using liquid chromatography with quadrupole – time-of-flight mass spectrometry. Tina Kosjek, Ester Heath, Sandra Pérez, Mira Petrovic, Damia Barceló. *Journal of Hydrology* 372 (2009) 109-117
- [140] Nitrification activity and community structure of nitrite-oxidizing bacteria in the bioreactors operated with addition of pharmaceuticals. Barbara Kraigher, Ines Mandic-Mulec. *Journal of Hazardous Materials* 188 (2011) 78-84
- [141] Nutrient and trace organic contaminant removal from wastewater of a resort town: Comparison between a pilot and a full scale membrane bioreactor. Hop V. Phan, Faisal I. Hai, James A. McDonald, Stuart J. Khan, Ren Zhang, William E. Price, Andreas Broeckmann, Long D. Nghiem. *International Biodeterioration & Biodegradation* (2015) 1-9
- [142] Effect of hypersaline aniline-containing pharmaceutical wastewater on the structure of activated sludge-derived bacterial community. Junmin Li, Zexin Jin. *Journal of Hazardous Materials* 172 (2009) 432-438
- [143] Membrane bioreactor for pharmaceutically active compounds removal: Effects of carbamazepine on mixed microbial communities implied in the treatment. Michael Aubenneau, Alexandre Tahar, Claude Casellas, Christelle Wisniewski. *Process Biochemistry* 45 (2010) 1826-1831
- [144] Effect of cytostatic drugs on microbial behaviour in membrane bioreactor system. Luis F. Delgado, Sylvie Schetrite, Carlos Gonzalez, Claire Albasi. *Bioresource Technology* 101 (2010) 527-536
- [145] Cytotoxicity micropollutant removal in a crossflow membrane bioreactor. Luis F. Delgado, Virginie Faucet-Marquis, Annie Pfohl-Leszkowicz, Christophe Dorandeu, Benedicte Marion, Sylvie Schetrite, Claire Albasi. *Bioresource Technology* 102 (2011) 4395-4401
- [146] Removal of pharmaceuticals, steroid hormones, phytoestrogens, UV-filters, industrial chemicals and pesticides by *Trametes versicolor*: Role of biosorption and biodegradation. Luong N. Nguyen, Faisal I. Hai, Shufan Yang, Jinguo Kang, Frederic D.L. Leusch, Felicity Roddick, William E. Price, Long D. Nghiem. *International Biodeterioration & Biodegradation* 88 (2014) 169-175
- [147] Influence of the amoxicillin concentration on organics removal and microbial community structure in an anaerobic EGSB reactor treating with antibiotic wastewater. Ling-

Wei Meng, Xiang-kun Li, Ke Wang, Kai-Li Ma, Jie Zhang. *Chemical Engineering Journal* 274 (2015) 94-101

[148] Influence of organic loading on the performance and microbial community structure of an anaerobic stage reactor treating pharmaceutical wastewater. Shreeshivadasan Chelliapan, Thomas Wilby, Ali Yuzir, Paul J. Sallis. *Desalination* 271 (2011) 257-264

[149] Factors affecting the removal of organic micropollutants from wastewater in conventional treatment plants (CTP) and membrane bioreactors (MBR). Magdalena Cirja, Pavel Ivashechkin, Andreas Schäffer, Philippe F. X. Corvini. *Rev Environ Sci Biotechnol* (2008) 7:61-78

[150] The ability of biologically based wastewater treatment systems to remove emerging organic contaminants—a review. Aida Garcia-Rodríguez & Víctor Matamoros, Clàudia Fontàs & Victòria Salvadó. *Environ Sci Pollut Res* (2014) 21: 11708-11728

[151] Effects of selected pharmaceutically active compounds on the ammonia oxidizing bacterium *Nitrosomonas europaea*. Shuyi Wang, Claudia K. Gunsch. *Chemosphere* 82 (2011) 565-572

[152] Pharmaceuticals as emerging organic contaminants in Umgeni River water system, KwaZulu-Natal, South Africa. Foluso O. Agunbiade & Brenda Moodley. *Environ Monit Assess* (2014) 186: 7273-7291

[153] Fate of diclofenac in municipal wastewater treatment plant — A review. Niina Vieno, Mika Sillanpää. *Environment International* 69 (2014) 28-39

[154] A review on the occurrence of micropollutants in the aquatic environment and their fate and removal during wastewater treatment. Yunlong Luo, Wenshan Guo, Huu Hao Ngo, Long Duc Nghiem, Faisal Ibney Hai, Jian Zhang, Shuang Liang, Xiaochang C. Wang. *Science of the Total Environment* 473-474 (2014) 619-641

[155] Impact of certain household micropollutants on bacterial behavior. Toxicity tests/study of extracellular polymeric substances in sludge. Laure Pasquini, Christophe Merlin, Lucille Hassenboehler, Jean-François Munoz, Marie-Noëlle Pons, Tatiana Görner. *Science of the Total Environment* 463-464 (2013) 355-365

[156] Effects of triclosan, diclofenac, and nonylphenol on mesophilic and thermophilic methanogenic activity and on the methanogenic communities. Evangelos C. Symaris, Ioannis A. Fotidis, Athanasios S. Stasinakis, Irimi Angelidaki. *Journal of Hazardous Materials* 291 (2015) 45-51.

(Received 26 June 2015; accepted 12 July 2015)